

72500-14

1-28-2009

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Ms. Kim Davis CC: APC Manager  
Reg West Company, LLC  
For The Scimetrics Ltd. Corporation  
30856 Rocky Road  
Greely, CO 80631-9375

JAN 28 2009

SUBJECT: Application for Pesticide Notification (PRN 98-10)  
Request General Label Change (Container Size)  
EPA Reg. 72500-14  
Application Dated December 29, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 12/29/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Please note that the Agency has issued a Pesticide Registration Notice (PR) 2007-4: Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment Statements." All labeling must be updated by August 17, 2010.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



Non-PRIA Revised Container Sizes Notification



December 29, 2008

Mr. John Hebert, Product Manager 7  
Document Processing Desk – **NOTIF**  
Office of Pesticide Programs – **7504P**  
**U.S. Environmental Protection Agency**  
2777 South Crystal Drive  
One Potomac Yard, Room S-4900  
Arlington, VA 22202

Dear Mr. Hebert:

**Subject:** Scimetrix Ltd. Corporation  
*Kaput*<sup>®</sup> Combo Bait Mini Blocks for Rodents & Fleas  
EPA Reg. No. 72500-14

On behalf of **Scimetrix Ltd. Corporation**, RegWest Company, LLC is submitting the attached documents as notification of revised container sizes for subject product. RegWest will act as sole agent in this endeavor.

Attached are the following:

- 1. Application for Pesticide: Other (Notification)
- 2. Two labels

We are changing the container sizes from:  
0.5, 1, 2, 10 and 20 lbs.

To the following container sizes (approximate number of blocks appears in parentheses):  
4 lbs. (64), 8 lbs. (130) and 18 lbs. (288)

Please contact me at (970) 353-0611 or kim@regwest.com if you require additional information or have any questions.

Regards,  
**RegWest Company, LLC**

Kim Davis, CC, APC, Manager  
Consultant/Agent

Attachments

pc: SLC – Poché, R.

30856 Rocky Road  
Greeley, CO 80631-9375  
970-353-0611  
970-353-0613 Fax  
Kim@regwest.com

4/17

# Kaput<sup>®</sup> Combo Bait Mini Blocks for Rodents & Fleas

Kills Norway Rats, Roof Rats, House Mice and Voles

First dead rats, mice and voles will appear 4 to 5 days after feeding begins.

Rodent Control Blocks

A Bait containing a Rodenticide to kill Norway Roof Rats, Roof Rats, House Mice and Voles and an Insecticide to Aid in the Reduction of Flea Populations on Listed Rodent Hosts.

**Active Ingredients:**

Imidacloprid (CAS Number 138261-41-3) ..... 0.020%

Warfarin (CAS Number 81-81-2) ..... 0.025%

**Other Ingredients** ..... 99.955%

**Total** ..... 100.000%

**NOTIFICATION**

**Keep Out of Reach of Children**

JAN 28 2009

**CAUTION**

**Harmful if Swallowed.**

**See back [side] panel for First Aid and additional Precautionary Statements.**

EPA Reg. No. 72500-14

EPA Est. 72500-CO-1

**Net Weight: 4 lbs. (1.81 kg) About 64 Blocks**

**[8 lbs. (3.63 kg) About 130 Blocks; 18 lbs. (8.16 kg) About 288 Blocks]**

Manufactured by:

**Scimetrics Ltd. Corp.**

P.O. Box 1045

Wellington, CO 80549

(970) 482-1330

mole@scimetricsltd.com

*New Solutions to Old Problems*

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling:

**Do not use this product to control any pests, by any application method or at any use sites, that are not specified on this label.**

Use of this product may aid in the reduction of fleas, including but not necessarily limited to *Xenopsylla cheopis*, that commonly infest listed rodent hosts, if ingesting sufficient quantities of this bait.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets or other non-target animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, other potentially destructive animals or in areas prone to vandalism.
3. Dispose of product container and unused or unconsumed bait as specified by this label.

**RATS AND MICE:**

**Use Restrictions:** This product may be used to kill Norway Rats [(*Rattus norvegicus*)], Roof Rats [(*Rattus rattus*)] and House Mice [(*Mus musculus*)] at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below and to aid in the reduction of their infesting flea populations. Outdoor applications to control rats or house mice must be made in tamper-resistant bait stations. Indoor applications must be made in tamper-resistant bait stations unless children younger than six years-of-age, pets and domestic animals are reliably denied access to the treated area by other means. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

**Urban Areas:** To control Norway rats, roof rats and house mice, this product may be used in and around homes, and industrial, commercial and public buildings; in sewers; in transport vehicles (ships, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations placed along walls or other external parts of buildings (e.g., doorways, ramps, loading docks) where rats or mice might seek to gain entrance.

**Non-urban Areas:** To control Norway rats, roof rats and house mice, this product may be used in and around homes and agricultural buildings. All outdoor placements must be made in tamper-resistant bait stations.

**Selection of Treatment Areas:** Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

**Application Directions:**

**Norway Rats and Roof Rats:** Apply 4 to 16 one-ounce bait blocks per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that rats are feeding on the bait.

**House Mice:** Apply 1 one-ounce bait block per typical placement. Two blocks may be placed at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

**Follow-up (Rats & Mice):** Immediately replace contaminated or spoiled bait. Collect and properly dispose of all visible dead animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

**VOLES:**

**Use Restrictions:** This product may be used to control meadow voles [(*Microtus pennsylvanicus*)], pine voles [(*M. pinetus*)] and montane voles [(*M. montanus*)] around (within 15 feet f) homes, and industrial, commercial, agricultural and public buildings and to aid in the reduction of their infesting flea populations. Do not use this product within the habitat of threatened or endangered *Microtus* spp. voles, threatened or endangered *Peromyscus* spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless

children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in temper-resistant bait stations. All bait applications must be made by hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

**Selection of Treatment Areas:** Determine areas near buildings where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

**Application Directions:** Place 1 one-ounce bait block in each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. If, and only if, children, pets, domestic animals and wildlife reliably are excluded from the treated area, one-ounce blocks may be placed directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board or ground vegetation, if present.

**Follow-up:** Check placements daily. Remove any fouled or spoiled bait and replenish if vole activity and interest persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

**Storage and Disposal**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide**

**Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-refillable container; do not reuse or refill this container. Triple rinse container (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or if allowed by state and local authorities, by burning; if burned, stay out of smoke.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Keep away from humans, domestic animals and pets. Harmful if swallowed because this material may reduce the clotting ability of the blood and cause bleeding. Causes moderate eye irritation. Do not get in eyes or on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating, drinking, chewing gum or using tobacco. Applicators must wear long-sleeved shirt, long pants, socks, shoes and gloves.

**First Aid**

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**If swallowed,** call a poison control center, doctor or 1-800-858-7378 for immediate treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. **If in eyes,** hold eye open and rise slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**Notice to Physician or Veterinarian**

This product reduces the clotting ability of the blood and may cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times. For animal cases, Vitamin K-1 is antidotal at 2-5 mg/kg. Repeated doses may need to be given up to two weeks (based on monitoring of prothrombin times). In severe cases, blood transfusion may be necessary.

**Environmental Hazards**

This product is highly toxic to aquatic invertebrates and toxic to mammals and birds. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply this product directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters.

