



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Mr. lain Weatherston Senior Regulatory Consultant Technology Sciences Group Inc 4061 North 156th Drive Goodyear, AZ 85395

OCT 6 2010

Subject:

Elector PSP Insect Control Agent, EPA Reg. No. 72642-2

Date of Registrant Submission: March 13, 2009

Decision: 416490

Dear Mr. Weatherston

The labeling referred to above, which adds direct poultry application to the label, submitted in connection with registration under the Federal insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

As a reference to the company's website is included on your label, please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance. As an alternative, you may refer consumers to the company's phone number or email address.

If you have any questions regarding this letter, please contact Samantha Hulkower at (703) 603-0683.

Sincerely.

Mark Syarez

Product Manager 13 Insecticide Branch

Registration Division (7505P)

Enclosure:

Copy of label stamped "Accepted"

Front Panel ELANCO*

AH0495-

ELECTOR® PSP INSECT CONTROL AGENT LOGO*

- Control of darkling beetles, hide beetles house flies, stable flies, little house flies (Fannia canicularis) in and around agricultural animal premises
- Control of northern fowl mites (Ornithonyssus sylviarum) infesting poultry.

Keep Out of Reach of Children

CAUTION

Elanco Animal Health, A Division of Eli Lilly and Company Indianapolis, IN 46285 www.elanco.com
Questions or comments call 1-800-428-4441

Net Contents: 8 oz (237 mL) Net Contents: 32 oz (946 mL)

*Elanco®, Elector® PSP Insect Control Agent and the diagonal color bar are trademarks of Eli Lilly and Company.



OCT 6 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under FPA Reg. No. 31.2.641-2

(Label Booklet	Cover-Back Panel)
ELANCO*	AH0495-

ELECTOR® PSP INSECT CONTROL AGENT LOGO*

- Control of darkling beetles, hide beetles house flies, stable flies, little house flies (Fannia canicularis) in and around agricultural animal premises
- Control of northern fowl mites (Ornithonyssus sylviarum) infesting poultry.

Environmental Hazards

This product is highly toxic to molluscs. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning or disposing of equipment washwaters. Apply this product only as specified on this label.

Notice: Shake Well Before Use. Avoid Freezing. Read the entire label before using: Use only according to label directions. See Directions for Use including storage and disposal inside label and the booklet.

EPA Reg. No. 72642-2 EPA Est. No. 37429-GA-02

*Elanco®, Elector® PSP Insect Control Agent and the diagonal color bar are trademarks of Eli Lilly and Company.

Elanco Animal Health
A Division of Eli Lilly and Company
Indianapolis, IN 46285
www.elanco.com
Questions or comments call 1-800-428-4441

Net Contents: 8 oz (237 mL) Net Contents: 32 oz (946 mL)

Peel Back Logo

7

(Label Booklet: Page 1 through end)

ELECTOR® PSP INSECT CONTROL AGENT LOGO*

- Control of darkling beetles, hide beetles house flies, stable flies, little house flies (Fannia canicularis) in and around agricultural animal premises Special and the second
- Control of northern fowl mites (Ornithonyssus sylviarum) infesting poultry.

ENVIRONMENTAL HAZARDS

This product is highly toxic to molluscs. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning or disposing of equipment washwaters. Apply this product only as specified on this label and the second of the sec

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Notice: Shake well before use. Avoid freezing.

MIXING FOR APPLICATIONS THAT REQUIRE DILUTION

This product is intended to be mixed with water and applied as a coarse or low-pressure spray. Fill the tank of the sprayer with water to about ½ of the total spray volume required. Start agitation and add the required amount of product. Continue mixing and agitation while filling the spray tank to the required spray volume. Maintain sufficient agitation during the application to ensure uniformity of the spray. Prepare spray fresh for each use.

GENERAL NOTICE

- This product may be applied as a general surface spray or a crack and crevice application. Applications may be made in and around poultry and livestock premises, including but not limited to poultry houses, beef and dairy barns, feedlots, stables, corrals and housing structures.
- This product may be used as a direct treatment to control Northern Fowl mites infesting poultry.
- Control is enhanced when interior and exterior perimeter applications are made in and around livestock and poultry structures. Normal sanitation practices of the structure consistent with good animal husbandry should be followed to enhance effectiveness of insect control agents.
- Lactating and non-lactating dairy and beef cattle, swine, sheep and poultry may be present at the time of premise treatment. Horses not intended for human consumption may also be present at the time of premise treatment.
- Cover or remove exposed feed and water from the area being treated.
- Do not use in milking rooms.
- Do not apply as a fog or space spray.

APPLICATIONS

TO CONTROL	APPLICATION
House flies Stable flies Little house flies	2 fl. oz. (60 mL) Elector PSP in 10 gallons of water will treat 5,000 – 10,000 ft². Use enough water in a directed spray to contact flies and their resting surfaces. Spray thoroughly, but do not allow run-off to occur.
Darkling beetles Hide beetles	Use enough water to ensure thorough coverage of the area treated, especially litter, along feed and water lines, walls and support beams. Use a sprayer of appropriate design that is equipped to effectively apply low to moderate volumes of spray. Low volume spray: Up to 15 gallons of diluted spray applied to
	20,000 ft ² . o Moderate volume spray: Up to 40 gallons of diluted spray applied to 20,000 ft ² .
	• 2 fl. oz. (60 mL) of product treats 5,000 ft ² (8 fl. oz. of product treats 20,000 ft ²).
	• For severe infestations, 4 fl. oz. of product may be used to treat 5,000 ft ² (16 fl. oz. treats 20,000 ft ²).
Northern fowl mite (Ornithonyssus sylviarum)	 Dilute 3 fl. oz. of product in 10 gallons of water. Apply 1 gallon of coarse spray per 100 birds under high pressure (no less than 100 – 125 psi), directed toward the vent area.

NOTE: One capful of product equals 2 ounces. Rinse cap after use and pour rinsate into spray tank (*This statement only on the 32 fl. oz. label*).

Egg Producing Poultry

Northern Fowl Mite: Re-application may occur every 14 days as needed. Spray birds in cages with a high pressure (100 – 125 psi) spray. Spray should be applied at the rate of 1 gallon per 100 birds. For optimum efficacy, direct spray toward vent area of chicken. Ensure adequate coverage.

Poultry Houses

House Flies, Stable Flies and Little house flies: Treat areas where flies rest: pay attention to side walls, the ends of buildings, upper sides of posts, and crossbeams. Spray in early morning when flies are resting. Use enough water in a directed spray to contact flies and their resting surfaces thoroughly, but do not allow run-off to occur. Do not contaminate feedstuffs, water or watering and feeding equipment with premise treatment. Re-application may occur every 7 to 10 days as needed, or when fly populations reach nuisance levels.

Darkling Beetles and Hide Beetles: Apply to floor area, especially to litter around feed and water lines and to walls and support beams, but do not allow run-off to occur. Application should also be made into cracks and crevices around insulation, and where pests have been seen or can find harborage. Reapply after each grow-out or sanitation procedure. Indoor control can be enhanced by making perimeter treatments to the outside of building foundations to prevent migrating adult beetles.

Livestock Housing

House Flies, Stable Flies and Little house flies: Apply to vertical and overhead surfaces where files may congregate. Feeders, waterers and feedcarts should be covered before application to prevent contamination. Do not contaminate feedstuffs or water with premise treatment. Pay attention to animal areas including fence railings, alleyways, stanchions, pipes, window and doors, and other areas where flies congregate. Spray in early morning when flies are resting. Re-application may occur every 7 to 10 days as needed, or when fly populations reach nuisance levels.

INTEGRATED PEST AND RESISTANCE MANAGEMENT

This product is a member of a unique class of insect control agents (spinosyns) with no pre-existing levels of resistance to any other insect control agents and can be rotated with all other classes, such as synthetic pyrethroids and organophosphates. House flies, stable flies, little house flies and other insects have demonstrated the ability to develop resistance to insect control products. Continual exposure of flies and other insects to a single class of insecticide may lead to the development of resistance to that class of insecticide. The following resistance management strategy will help ensure the prolonged usefulness of this product, as well as other conventional insecticides:

- Apply this product and other insect control agents only according to label directions
- Do not use less than the labeled rate of any insect control product
- Include multiple control tactics (cultural and/or biological controls) within an Integrated Pest Management Program
- Rotate this product with other classes of insecticides (synthetic pyrethroid, organophosphate) during a fly season

For advice concerning integrated pest management practices in relation to specific local conditions, consult resources in resistance management programs, your Cooperative Agricultural Extension Service, or your Elanco Animal Health representative (800-428-4441).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage:	Store in original container only, tightly closed in a secure place. Keep out of reach of children and domestic animals. In case of leak or spill, remove with absorbent materials and dispose of as waste.
Pesticide Disposal:	Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility
Container Handling:	Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then dispose of empty container in a sanitary landfill, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with the use of this product. Lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions, abnormal conditions, presence of other materials, the manner of application, or other factors which are beyond the control of Elanco Animal Health or the seller. To the extent consistent with applicable law all such risks are assumed by the buyer.

LIMITATIONS OF REMEDIES

To the extent consistent with applicable law all the exclusive remedy for losses or damages resulting the exclusive remedy for losses or damages resulting the following from this product (including claims based on contract, negligence, strict liability, or other legal theories); the strict liability or other legal theories is shall be limited to, at Elanco Animal Health's election, one of the following the strict liability is a strict liability.

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

To the extent consistent with applicable law all Elanco Animal Health shall not be liable for losses or damages resulting from the handling or use of this product unless Elanco Animal Health is promptly notified of such loss or damage in writing. In no case shall Elanco Animal Health be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Elanco Animal Health or the seller is authorized to vary or exceed the terms of Warranty Disclaimer of this Limitation of Remedies in any manner.

Elanco Animal Health, A Division of Eli Lilly and Company Indianapolis, IN 46285 www.elanco.com
Questions or comments call 1-800-428-4441

EPA Reg. No.: 72642-2 EPA Est. No.: 37429-GA-02

*Elanco®, Elector® PSP Insect Control Agent, and the diagonal color bar are trademarks of Eli Lilly and Company