

72693-14

05-02-2012

1
18**U.S. ENVIRONMENTAL PROTECTION AGENCY**

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

NOTICE OF PESTICIDE:

- ☒ Registration
☐ Reregistration
Under FIFRA, as amended

EPA Reg. Number:

72693-14Date of
Issuance:**MAY - 2 2012**

Term of Issuance:

Unconditional

Name of Pesticide Product:

Tebucon 3.6F

Name and Address of Registrant (include ZIP Code):

Universal Crop Protection Alliance, LLC
1300 Corporate Center Curve
Eagan, MN 55121

Mailed to:

Ross Gilbert
Pyxis Regulatory Consulting, Inc.
4110 136th Street, NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Page 1 of 2

Signature of Approving Official:

Mary L. Waller, Product Manager (21)
Fungicide Branch/Registration Division/OPP/OCSP (7504P)

Date:

5/2/2012

MAY - 2 2012

2
18


2. Make the following changes to the label:

- a. Change the product registration number to "EPA Reg. No. 72693-14"
- b. On page 2 in the **USER SAFETY RECOMMENDATIONS** box add the following:
 - Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."
- c. On page 3 in the second line of the third paragraph change "...soil or water, is:" to "...soil or water is:"
- d. On page 7 in the list of crops under Dry bulb onion delete "Welch Onion".
- e. On page 8 in the list of crops under Green Onion add "Welsh Onion".
- f. On page 14 in the section **SEED LABELING**:
 1. Enclose the seed tag text in quotation marks and add the following text so that the seed tag reads as follows: "TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice."
 2. Change "180.1001(c) and (d)" to "180.910, 180.920 and 180.950"
- g. On page 15 in the **STORAGE AND DISPOSAL** box, in the instructions for **PESTICIDE DISPOSAL** change "...product must be...." to "...product must be...."
- h. On page 16 in line 2 in the first paragraph of the **CONDITION OF SALE AND WARRANTY** change "...and should be followed carefully." to "...and must be followed carefully."

Submit one copy of the revised final printed label for the record before the product is released for shipment.

Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.


Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure:

Label stamped "Accepted with Comments"

Note to reviewer: [] brackets indicate optional text

Tebucon 3.6F

For control of specified diseases on [asparagus], [barley], [beans], [corn], [cotton], [cucurbit vegetables], [grasses grown for seed], [hops], [dry bulb onion and garlic], [green onions], [leafy Brassica greens], [garden beet roots and tops], [lychee], [okra], [pecan], [peanut], [soybean], [sunflower], [turnip], [and] [wheat].

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-

1H-1,2,4-triazole-1-ethanol 38.7%

OTHER INGREDIENTS: 61.3%

TOTAL: 100.0%

Contains 3.6 pounds tebuconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Refer to the inside of the label booklet for additional Precautionary Statements, First Aid Statements, Worker Protection Statements, Directions for Use, Storage and Disposal Instructions, and the Conditions of Sale and Warranty.

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious or convulsing person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24 Hour Medical Emergency Assistance (Human or Animal) call 1-800-308-1241 or for Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call CHEMTREC at 1-800-424-9300.

Note to Physician: No specific antidote. Treat symptomatically.

Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

EPA Reg. No. 72693-XX

Manufactured For:

Universal Crop Protection Alliance, LLC
1300 Corporate Center Curve
Eagan, MN 55121
(651) 239-1000

EPA Est. No. XXXXX-XX-XXX

Net Contents:

ACCEPTED
with COMMENTS
In EPA Letter Dated:

5/2/2012

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

72693-14

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as bather laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes and socks

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-)], the handler PPE requirements may be reduced or modified as specified in WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into the ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and

5
18

exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

PRODUCT INFORMATION

Spray Volume: Apply Tebucon 3.6F by ground in a minimum of 10 gallons of spray solution per acre or by air in a minimum of 5 gallons of spray solution per acre. Check equipment calibration frequently. For best disease control, complete coverage and uniform application are essential, especially when lower spray volumes are used. Use the appropriate spray volume to ensure complete coverage.

Chemigation: Apply Tebucon 3.6F through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add specified amount of Tebucon 3.6F to the spray tank while filling with water to the desired level. Maintain agitation while mixing. If other materials are added to the spray tank, be sure that the Tebucon 3.6F is thoroughly dispersed before other materials are added. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

Compatibility: To determine the compatibility of Tebucon 3.6F with other mixing partners, use the following procedure: Pour the recommended proportions of the products into a suitable container of

water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information, contact your local Universal Crop Protection Alliance, LLC representative.

Resistance Management Statement

Tebucon 3.6F is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucon 3.6F and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Universal Crop Protection Alliance, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCON 3.6F
Asparagus	Rust (<i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon 3.6F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of Tebucon 3.6F per acre (0.11 lb. ai – 0.17 lb. ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl. oz./acre or 0.51 lb. ai/acre).	
General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tebucon 3.6F with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Barley	Rust (<i>Puccinia</i> spp.)	4 fl. oz. per acre
	Head Blight (<i>Fusarium</i> spp.) – Suppression	
	Notes: Apply Tebucon 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Tebucon 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tebucon 3.6F. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.	

CROP	DISEASE	RATE OF TEBUCON 3.6F
	Application timing directions: Rusts: Apply Tebucon 3.6F at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon 3.6F for Fusarium Head Blight suppression is when stem heads have fully emerged (Feekes 10.5) on 50% of the plants.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Beans (fresh & dry except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 to 6 fl. oz. per acre
	Notes: Apply Tebucon 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Tebucon 3.6F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon 3.6F per acre per crop season. Beans, dry: Tebucon 3.6F may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Tebucon 3.6F per acre per crop season.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours drying time on bean foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Corn (sweet corn, field corn, field corn grown for seed and popcorn)	Rust (<i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	Northern leaf blight (<i>Helminthosporium turcicum</i>)	
	Southern leaf blight (<i>Helminthosporium carbonum</i>)	
	Gray leaf spot (<i>Cercospora zeae-maydis</i>)	
	Notes: Apply Tebucon 3.6F in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pints) of Tebucon 3.6F may be applied per acre per crop season. Sweet corn: Tebucon 3.6F may be applied up to 7 days before harvest of ears or forage and 49 days before the harvest of fodder. Field, seed or popcorn: Tebucon 3.6F may be applied up to 21 days before harvest of forage and 36 days before the harvest of grain or fodder.	
General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should		

8
18

be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on corn foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).
 Restricted-entry interval (REI) for sweet corn = 19 days.
 Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz. per acre
	Notes: Apply Tebucon 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Tebucon 3.6F may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Tebucon 3.6F per acre per crop season.	

General Comments: For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on cotton foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).
 Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl. oz. per acre
	Gummy stem blight – suppression (<i>Didymella bryoniae</i>) (watermelon, squash, pumpkin and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Tebucon 3.6F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Tebucon 3.6F per season.	

CROP	DISEASE	RATE OF TEBUCON 3.6F
Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Dry bulb onion Garlic Great-headed (elephant) garlic Welch Onion Shallot	White rot (<i>Sclerotium cepivorum</i>)	White rot: 20.5 fl. oz. per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (<i>Puccinia dill</i> , <i>Puccinia porii</i>)	4 to 6 fl. oz. per acre
	Purple blotch (<i>Alternaria porii</i>)	
	White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz. Tebucon 3.6F 6 per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl. oz./acre. Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz. Tebucon 3.6F per acre per application. Repeat at 10- to 14-day intervals. Apply Tebucon 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 32.5 fl. oz. of Tebucon 3.6F per acre per season if an in-furrow treatment is made. If Tebucon 3.6F is not applied as an in-furrow treatment then do not apply more than 12 fl. oz. Tebucon 3.6F per acre per season as a foliar spray. Do not apply within 7 days of harvest.	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of spray surfactant may be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering.		

Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).
Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Grasses Grown for Seed	Rust (<i>Puccinia</i> spp.)	4 to 8 fl. oz. per acre
	Apply the specified rate of Tebucon 3.6F as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./acre and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of Tebucon 3.6F when powdery mildew first appears on leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz/acre and shorter spray intervals.	

General Comments: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon 3.6F.

A maximum of 16 fl. oz. (1 pint) may be applied per acre per crop season. Tebucon 3.6F may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Green Onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (<i>Sclerotium cepivorum</i>) Suppression only Rust (<i>Puccinia allii</i> , <i>Puccinia porii</i>) Purple blotch (<i>Alternaria porii</i>)	4 to 6 fl. oz. per acre
For control of diseases, make foliar applications using an interval of 10- to 14-days. Apply Tebucon 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Notes: Do not apply more than 24 fl. oz. of Tebucon 3.6F per acre per season. Do not apply within 7 days of harvest.		

General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of spray surfactant may be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
------	---------	----------------------

Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Tebucon 3.6F may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Tebucon 3.6F per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	
General Comments: For optimum disease control, the lowest labeled rate of spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Leafy & Brassica Greens	Cercospora leaf-spot (<i>Cercospora brassicicola</i>)	3 to 4 fl. oz. per acre
Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna	Powdery mildew (<i>Erysiphe cruciferarum</i>)	
Mustard greens Mustard spinach Rape greens Turnip greens	Alternaria leaf spot (<i>Alternaria brassicicola</i>)	
Notes: Do not apply more than 16 fl. oz. of Tebucon 3.6F per season. Do not apply within 7 days of harvest. Do not apply more often than once every 10 days.		
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restriction: Application to turnip greens is limited to East of the Rockies. Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Garden beet, roots and tops (leaves)	Cercospora leaf-spot (<i>Cercospora beticola</i>)	3 to 7.2 fl. oz. per acre
	Notes: Make applications on 14-day intervals. Do not apply more than 28.8 fl. oz. of Tebucon 3.6F per acre per season. Do not apply within 7 days of harvest.	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled		

rate of a spray surfactant may be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).
Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Lychee	Anthracnose (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl. oz. per acre
	Notes: Begin first application of Tebucon 3.6F as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Tebucon 3.6F per acre per season. Tebucon 3.6F can be applied up to and including the day of harvest (PHI = 0 days).	
General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 2 days		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Okra	Cercospora leaf-spot (<i>Cercospora beticola</i>)	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon 3.6F in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution per air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Tebucon 3.6F per acre per season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
------	---------	----------------------

CROP	DISEASE	RATE OF TEBUCON 3.6F	
Peanut	SOILBORNE: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) FOLIAR: Early leaf spot Late leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptosphaerulina</i>)	7.2 fl. oz. per acre	
	FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventative spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Tebucon 3.6F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest specified rate of a spray surfactant should be tank mixed with Tebucon 3.6F.		
	LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tebucon 3.6F in the first advisory spray in July and continue Tebucon 3.6F applications at 14-day intervals. Applications after August 15 should be tank-mixed with chlorothalonil for resistance management purposes.		
General Comments: For optimum control of the specified soilborne diseases, four consecutive applications of Tebucon 3.6F must be made at 14-day intervals. A maximum of 28.8 fl. oz. of Tebucon 3.6F may be applied per crop season. Tebucon 3.6F may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas. Tebucon 3.6F is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Chlorothalonil may be tank-mixed at the rate of 12 oz of active ingredient with Tebucon 3.6F as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tebucon 3.6F is tank-mixed with chlorothalonil. Mixing or alternating Tebucon 3.6F with other DMI fungicides may lead to resistance. Tebucon 3.6F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i> . Drought conditions will decrease the effectiveness of Tebucon 3.6F against the root and pod rots. Use Tebucon 3.6F in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices. Restricted-entry interval (REI) = 12 hours			
Timing of Tebucon 3.6F			
Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot			
Spray Program	Tebucon 3.6F Application No.	Chlorothalonil Application No.	
7 applications	3, 4, 5 and 6	1, 2 and 7	

CROP	DISEASE	RATE OF TEBUCON 3.6F
Pecan	Brown leaf spot (<i>Sirosporium cliffusium</i>)	4 to 8 fl. oz. per acre
	Downy spot (<i>Mycosphaerella caryigena</i>)	
	Liver spot (<i>Gnomonia caryae</i>)	
	Scab (<i>Cladosporium caryigenum</i>)	
	Vein spot (<i>Gnomonia nerviseda</i>)	
	Zonate leaf spot (<i>Grovesinia pyramidalis</i>)	
Notes: Apply Tebucon 3.6F in a preventative spray schedule beginning at early bud break (young leaves unfolding) and continue applications at 10- to 14-day intervals through the pollination period. Tebucon 3.6F should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for Super-Tin. Do not add a surfactant to the spray solution when using Super-Tin. Apply Tebucon 3.6F in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Tebucon 3.6F to full-size mature trees and 4 to 6 fl. oz. per acre of Tebucon 3.6F to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Tebucon 3.6F may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to grazed treated areas.		
General Comments: For optimum disease control, the lowest labeled rate of spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCON 3.6F
Soybean	Rust (<i>Phakopsora pachyrhizi</i>)	3 to 4 fl. oz. per acre
	Powdery mildew (<i>Microsphaera diffusa</i>)	
Use Directions: Apply Tebucon 3.6F as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications at 10- to 14-day intervals if environmental conditions are favorable for continued disease development. Use of higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray		

15
18

surfactant must be tank-mixed with Tebucon 3.6F. Tebucon 3.6F should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz. per acre per use season.
Restricted-entry interval = 12 hours

CROP	DISEASE	RATE OF TEBUCON 3.6F
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of Tebucon 3.6F at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Tebucon 3.6F per acre per season or within 50 days of harvest.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon 3.6F. Contact your state Extension Service or UPI representative for a list of approved surfactants. Tebucon 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicicola</i>)	4 to 7.2 fl. oz. per acre
	Notes: Apply the specific dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Tebucon 3.6F may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Tebucon 3.6F per acre per crop season.	

General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUCON 3.6F
Wheat	Rusts – leaf, stem and stripe (<i>Puccinia</i> spp.)	4 fl. oz. per acre
	Head blight or scab (<i>Fusarium</i> spp.) - Suppression	
	Notes: Wheat fields should be observed closely for early disease symptoms.	

16
18

	<p>particularly when susceptible varieties are planted and/or under prolonged conditions are favorable for disease development. A maximum of 4 fl. oz. of Tebucon 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tebucon 3.6F. Apply Tebucon 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p>Application timing directions: Rusts: Apply Tebucon 3.6F at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tebucon 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Tebucon 3.6F. Tebucon 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tebucon 3.6F will be resistant to weathering. Tebucon 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>	

<p>SEED TREATMENT – Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.</p>		
<p>SEED LABELING: To meet US Federal Seed Act requirements, all seed treated with Tebucon 3.6F must be labeled:</p> <p style="text-align: center;">TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole.</p> <p>USE PRECAUTION: When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored with an EPA approved dye such as one of the dyes listed in the 40 CFR Section 180.1001(c) and (d) to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.</p>		
DISEASE	RATE FL. Oz./CWT	DIRECTIONS FOR USE
Soilborne and Seedborne Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Tebucon 3.6F. The length of control will vary depending on the rate used.
Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>)	0.27-0.54	

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

17
18

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (five gallons or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

Nonrefillable Container (greater than five gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate

18
18

into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If container is burned, stay out of smoke.

CONDITION OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UNIVERSAL CROP PROTECTION ALLIANCE, LLC, its Supplemental Distributors, or the Seller. All such risks shall be assumed by the Buyer.

UNIVERSAL CROP PROTECTION ALLIANCE, LLC, its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER UNIVERSAL CROP PROTECTION ALLIANCE, LLC NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF UNIVERSAL CROP PROTECTION ALLIANCE, LLC, ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION, THE REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL UNIVERSAL CROP PROTECTION ALLIANCE, LLC, ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. UNIVERSAL CROP PROTECTION ALLIANCE, LLC, its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of UNIVERSAL CROP PROTECTION ALLIANCE, LLC.

No employee or agent of UNIVERSAL CROP PROTECTION ALLIANCE, LLC, its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

[EPA approval date]