
PYRENONE[®] INDUSTRIAL SPRAY E.C.



 A FOOD INDUSTRY INSECT SPRAY FOR USE IN BAKERIES, RESTAURANTS, FOOD PROCESSING PLANTS, INSTITUTIONS, RESIDENCES AND OFFICE BUILDINGS.

02/17/2004

- MAY BE DILUTED WITH EITHER WATER OR OIL.
- AUTHORIZED FOR USE IN ALL DEPARTMENTS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT, POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS.
- **PROVIDES RAPID KILL OF INSECTS WITHOUT OBJECTIONABLE RESIDUES.**
- FOOD ADDITIVE TOLERANCES HAVE BEEN ESTABLISHED FOR THE INSECTICIDE COMPONENTS IN THIS PRODUCT. WHEN THIS PRODUCT IS USED AS DIRECTED, ALLOWABLE TOLERANCE LEVELS WILL NOT BE EXCEEDED.

ACTIVE INGREDIENTS

73049-241

Pyrethrins	1.0 %
Piperonyl Butoxide*	
OTHER INGREDIENTS**	
	100.0 %

*Equivalent to 8% (butylearbityl)(6-propylpiperonyl) ether and 2% related compounds. **Contains petroleum distillate

PRECAUCION AL CONSUMIDOR: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 73049-241

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN

CAUTION

	FIRST AID
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes, Call a poison control center or doctor for treatment advice.
• If swallowed:	 Immediately call poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
	HOT LINE NUMBER
In case of Medical emer	ner or label with you when calling a poison control center or doctor, or going for treatment gencies or health and safety inquiries or in case of fire, leaking or damaged containers, uned by calling 1-877-315-9819.

See Side Panel For Additional Precautions

NET CONTENTS:

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

CAUTION

Avoid contact with skin, eyes and clothing. Prolonged or repeated skin contact may cause allergic retotions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Remove any birds, pets or fish aquaritums before spraying. Vacate premises after treatment and ventilate before reoccupying.

Environmental Hazards

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Confine the use of this product in food processing plants to time periods when the plant is not in operation. Cover or remove food during treatment. Cover all food processing surfaces during treatment or thoroughly wash with an effective cleaning compound followed by a potable water rinse before using.

For maximum effectiveness, a combination of localized application and space treatment is recommended. When diluting with water, prepare only the amount needed for immediate use. Holding a water dilution more than 24 hours is not recommended. Agitate occasionally. For oil dilutions, a deodorized petroleum oil is recommended.

AS A SURFACE SPRAY: For initial clean up of severe insect infestations, dilute at the rate of 1 part concentrate in 4 parts water or oil. This dilution may not be used in edible product areas of Federally inspected meat and poultry plants. In these plants and in other areas for normal infestations, dilute 1 part concentrate in 9 parts water or oil.

To control Ants, Cockroaches, Crickets, Clover Mites, Cheese Mites, Confused Flour Beetles, Cigarette Beetles, Cadelles, Drugstore Beetles, Granary Weevils, Grain Mites, Meal Worms, Rice Weevils, Rust Red Flour Beetles, Silverfish, Spiders, Sawtoothed Grain Beetles, and Spider Beetles: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into cracks and crevices, under pallets, around containers of stored foods, around the base of machinery and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

TO CONTROL *Carpet Beetles*: Spray the edges of carpeting and under carpeting and rugs. Make localized application to floors and baseboards. Spray directly into cracks, crevices, closets and infested areas of shelving.

TO CONTROL *Bedbugs*: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors.

TO CONTROL *Fleas and Brown Dog Ticks*: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and the entire floor and floor coverings. Place fresh bedding in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended.

AS A SPACE SPRAY: Dilute at the rate of 1 part concentrate to 19 parts water or oil. Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems.

For rapid control of Angoumois Grain Moths, Fruit Flies, Flies, Gnats, Hornets, Indianmeal Moths, Mosquitoes, Mediterranean Flour Moths, Skipper Flies, Tobacco Moths and Wasps: Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet. Keep area closed for at least 10 minutes. Ventilate area before reoccupying. For rapid kill of exposed or accessible stages of any of the insects named on this label, dilute 1 part concentrate with from 1 to 9 parts water or oil. Apply with mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at 1 to 10 ounces per 1000 cubic feet of space.

When used in meat and poultry processing plants a dilution of not less than 1 to 1 (50%) is acceptable. It must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns).

OUTDOORS: To control *Brown Dog Ticks* around residential areas, rural fields or wood lots, roadside edges, pathways, and coastal vegetation areas. Dilute 2.5 gallons of material with 50 gallons of water. Apply with a mist blower applicator calibrated to deliver three gallons per acre at 200 psi. Make applications during peak of infestation and repeat if necessary. Apply directly to vegetation where the ticks rest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage And Spill Procedures: Store upright at room temperature. Avoid exposure to excerne temperatures. In case of spil or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste? Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability. CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Valent BioSciences Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: VALENT BIOSCIENCES CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE. THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Valent BioSciences Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Valent BioSciences Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES. INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT. WARRANTY, TORT. NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID. OR AT VALENT BIOSCIENCES CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

OValent BioSciences Corporation, 2003

Pyrenone is a registered trademark of Bayer AG.

Valent BioSciences Corporation Environmental Science Division

870 Technology Way Libertyville, IL 60048

List No. 73049241

02-0090/R1