

73049-305

3/6/2012

1087



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

MAR 6 2012

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Nasser Assaf, Ph.D.  
Valent BioSciences  
870 Technology Way, Suite 100  
Libertyville, IL 60048

**Subject: Amended Reregistration Label Permanone® General Purpose Aqueous Insecticide II  
D413058  
EPA Registration No. 73049-305**

Dear Dr. Assaf:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the referenced product in connection with the **Permethrin RED**, and has concluded that your submission is acceptable.

**PLEASE NOTE:** This product is not being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time.

**PLEASE NOTE:** That the record for this product currently contains the Confidential Statements of Formulation (CSFs) listed below. Any previously dated CSF's are superseded.

- Basic CSF, dated: March 12, 2010

A copy of your label stamped Acceptable is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped after 12 months from the date of this amendment or the next printing of the label whichever occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

If you have any questions about this letter, please contact Linda A. DeLuise at (703) 305-5428 or via e-mail at [deluise.linda@epa.gov](mailto:deluise.linda@epa.gov) or Mr. Richard J. Gebken at (703) 305-6701 or via e-mail at [Gebken.Richard@epa.gov](mailto:Gebken.Richard@epa.gov).

Sincerely yours,

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken  
Product Manager 10  
Insecticide Branch  
Registration Division (7505P)

Enclosures: Label stamped Accepted dated: March 6, 2012

2 of 7


# PERMANONE® GENERAL PURPOSE AQUEOUS INSECTICIDE II

- *For Commercial/Industrial Use Only*
- *[A Ready-To-Use Permanone® Stable Emulsion Insecticide*
- *For Use Outdoors, In Homes And Non-Food Areas Of Food/Feed Processing Establishments, Kennels, Commercial Buildings, Warehouses, Theaters, Office Buildings, Schools, Motels, And Hotels*
- *For Control Of Fleas, Brown Dog Ticks And Lice On Premises And On Dogs*
- *For Control Of Ticks Which May Cause Lyme Disease On Premises*
- *Controls Fleas And Ticks On Dogs For 14 Days]*

**ACTIVE INGREDIENT:**

Permethrin*	0.50%
<b>OTHER INGREDIENTS:</b>	<u>99.50%</u>
	100.00%

\* Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.



**PRECAUCION AL CONSUMIDOR:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**(TO THE USER:** If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 73049-305

EPA EST. NO.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**See Rear Panel For Additional Precautions**

For information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-877-315-9819.

**NET CONTENTS:**

3087

**READ ENTIRE LABEL BEFORE EACH USE  
PET USE APPLIES TO USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS  
Hazards To Humans & Domestic Animals**

**CAUTION**

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**Personal Protective Equipment (PPE):**

Mixers, loaders, applicators and other handlers, (including during mixing/loading, repair and cleaning of equipment and disposal of pesticide) must wear the following:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any water proof material
- Shoes plus socks

In addition to the above PPE, mixers and loaders and/or applicators using high-pressure handwand in an enclosed area must wear at least a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval prefix TC-21C or any N, R, P, or HE filter.

In addition to the above PPE, applicators using hand-held foggers in an enclosed area must wear a half-face or full face or hood style NIOSH approved respirator with a dust/mist filtering cartridge with MSHA/NIOSH approval number prefix TC-21C, or a cartridge or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a cartridge or canister with any N, R, P, or HE filter.

**User Safety Recommendations:**

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Environmental Hazards**

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutter or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**[Containers equal or greater than 5 gallons:]**

[Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.]

4097

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Entry Restrictions:**

- [For surface spray use] Except when applying to [dogs,][livestock] [and clothing], do not enter or allow others to enter treated area until sprays have dried.
- [For space spray use] Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

**Application Restrictions:**

- Except when applying to [dogs,][livestock] [and clothing], do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to [dogs,][livestock] [and clothing], only protected handlers may be in the area during application.

**Use Restrictions:**

- Do not breathe vapors or spray mist.
- Do not apply when food is present.
- Do not use in food areas of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held.
- May be applied in non-food/feed areas. Non-food/feed establishment areas are areas such as garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after canning or bottling).
- In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.
- Remove or cover exposed food and drinking water before application.
- Do not contaminate animal feed or water during application.
- Do not apply as a broadcast or spot-treatment to indoor surfaces at residential sites, including nurseries, daycare centers, schools, hospitals, and nursing homes.
- Do not use in aircraft cabins.
- Do not use on electrical equipment or on surfaces that can be harmed or stained by water.
- Not for plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks or on golf courses or lawns and grounds.
- Not for broadcast use.
- Not for use in outdoor or indoor residential misting systems.
- Remove pets, birds and cover any water inhabited by fish (such as aquariums and ornamental fish ponds) before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.
- *[Alternate text For pet care:* Do not apply to dogs less than 12 weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.]
- *[Alternate text - When used in dairy barns or facilities:* Close milk bulk tank lids to prevent contamination from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.]
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- *[Alternate text – outdoor application:*
  - All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses:

- 1) Treatment to soil or vegetation around structures
  - 2) Applications to lawns, turf, and other vegetation
  - 3) Application to building foundations, up to a maximum height of 3 feet
- Other than application to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications, only.
  - Do not water the treated area to the point of run-off.
  - Do not make applications during rain.]

**SHAKE WELL BEFORE USING****INDOOR APPLICATION**

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

**SURFACE SPRAYING:**

TO CONTROL *Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites* and *Cadellés*: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For *Silverfish*, spray bookcases. For *Ants*, spray trails, nests and points of entry.

TO CONTROL *Carpet Beetles*: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving.

TO CONTROL *Bedbugs*: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors.

TO CONTROL *Fleas (Adults and Larvae)* and *Ticks, Including Ticks Which May Carry Lyme Disease (Adults and Larvae)*: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL *Clothes Moths*: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL *Lice* and *Louse Eggs*: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

**SPACE SPRAYING:**

Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of *Houseflies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths* and *Tobacco Moths*: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

**FOGGING:**

To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before reoccupying.

6007

**FOR DIRECT APPLICATION TO LIVESTOCK:**

To protect cattle (beef and dairy), goats, sheep, hogs and horses from *Horn Flies*, *Face Flies*, *Houseflies*, *Mosquitoes* and *Gnats*: Apply a light mist sufficient to wet the surface of the hair. For *face flies*, spray face and head, but do not spray into eyes.

TO CONTROL *Stable Flies*, *Horse Flies* and *Deer Flies*: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL *Blood-Sucking Lice*: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL *Poultry Lice*: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL *Bedbugs* and *Mites* IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**FOR DIRECT APPLICATION TO DOGS (OVER 12 WEEKS OF AGE):**

TO CONTROL *Fleas*, *Ticks*, and *Lice*:

Hold the container upright. Spray at a distance of 8-12 inches. Starting at the tail, spray over entire hair coat while moving the dispenser rapidly; making sure the coat is thoroughly wetted to the skin. While spraying, fluff the hair so the spray will penetrate to the skin. Make sure spray wets ticks thoroughly. Spray against the lay of the hair, roughing the hair on long-haired dogs. Do not spray into the eyes, face or genitalia. For best results in killing ticks, spray directly on the tick. Allow dog to air dry or blow dry. Do not towel dry. Do not reapply more than once every 14 days.

Use protective gloves or mitts when applying product. DO NOT USE ON CATS OR KITTENS.

**OUTDOOR APPLICATION**

TO CONTROL *Ticks*, Including *Ticks Which May Carry Lyme Disease* (PERIMETER TREATMENT): Use Permanone7 General Purpose Aqueous Insecticide II as a residual treatment to control Ticks including the *Deer Tick* (*Ixodes scapularis*), *Lone Star Tick* (*Amblyomma americanum*), *Brown Dog Tick* (*Rhipicephalus sanguineus*), *American Dog Tick* (*Dermacentor variabilis*) and *Western Black-Legged Tick* (*Ixodes pacificus*).

For each area listed below, apply until surface is wet (approximately 1 gal/750-1000 ft<sup>2</sup>).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6-10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**FOR CONTROL OF ORNAMENTAL GARDEN INSECTS:**

For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies*, *Aphids*, *Exposed Thrips*, *Spiders*, *Red Mites* and *Leafminers*. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

EFFECTIVE WHEN USED AS DIRECTED AGAINST *Whiteflies, Aphids, Spider* and *Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites* and *Armyworms*: When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place away from heat or open flame. Avoid extremes in temperature.

**Pesticide Disposal:** Pesticide, spray mixture or rinse water that cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

#### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Valent BioSciences Corporation. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, VALENT BIOSCIENCES CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Valent BioSciences Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Valent BioSciences Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT VALENT BIOSCIENCES CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

[Or alternate text

Note [Notice]: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, seller shall have no liability for consequential damages.]

[Or Alternate Text

To the extent consistent with applicable law, Purchaser assumes all responsibility for safety and use not in accordance with directions.]

© Valent BioSciences Corporation, 2012  
Permanone is a registered trademark of Bayer AG.

**Valent BioSciences Corporation**  
**Environmental Science Division**  
870 Technology Way  
Libertyville, IL 60048