


PERMANONE® MULTI-PURPOSE 0.25% SPRAY

- *[Contains the Synthetic Pyrethroid Permanone®]*
- *Controls Named Residential, Industrial and Garden Insects*
- *For Use on Dogs and on Premises to Control Fleas and Ticks*
- *An Aqueous Stable Microemulsion*
- *Controls Pests for up to Four Weeks*
- *Kills Lice and Their Eggs*
- *For Use on Bedding, Furniture and Other Inanimate Objects Infested with Lice.*
- *For Use on Livestock and Their Premises*
- *Controls Fleas and Ticks on Dogs for 14 Days!*
- *Not for Use on Humans*



ACCEPTED
 DEC 22 2004
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 73049-306

ACTIVE INGREDIENT:

Permethrin¹ 0.25 %
OTHER INGREDIENTS*: 99.75 %
100.00 %

¹ Cis/trans ratio: Max. 55% (+/-) cis and Min. 45% (+/-) trans.
 *Contains petroleum distillate

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
 (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 73049-306

EPA EST. NO.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If On Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-877-315-9819.	
NOTE TO PHYSICIAN	
Contains petroleum distillate - vomiting may cause aspiration pneumonia.	

See Side Panel For Additional Precautionary Statements

NET CONTENTS:

2/4

**READ ENTIRE LABEL BEFORE EACH USE
FOR PET USES: USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS
Hazards To Humans & Domestic Animals**

CAUTION

Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water.

Physical and Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not use in the edible product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.

In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment. Cover or remove exposed food.

Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. **This product is not for use on humans.** If lice infestations should occur on humans, use a product labeled for use on humans.

NON-FOOD AREAS: Permanone Multi-Purpose 0.25% spray may be applied in non-food/feed areas. Non-food/feed establishment areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after canning or bottling).

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Test product on an inconspicuous area before applying. Spray until surface is slightly wet, but not to the point of run-off. Over application may cause damage. Will not stain surfaces or fabrics where water alone causes no damage. Remove birds and cover fish aquariums before spraying. Prevent re-entry of dogs and cats into the treated area until spray deposits are dry.

Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion.

Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible.

Ants: Spray trails, nests and points of entry. Spray on ants where possible.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment.

Exposed Stages of Confused Flour Beetles, Saw-toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. In the home, dispose of infested stored food products.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse.

TO CONTROL *Clothes Moths* and *Carpet Beetles* on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of *Houseflies*, *Mosquitoes*, *Gnats*, *Wasps*, *Hornets*, *Bees* and *Small Flying Moths*. Remove animals prior to application. Close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of *Stable Flies*, *Horse Flies*, *Deer Flies*, *Face Flies*, *Houseflies*, *Horn Flies*, *Mosquitoes*, and *Gnats*: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL *Blood Sucking Lice*: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair.

TO CONTROL *Poultry Lice*. spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF *Bedbugs* and *Mites* IN POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

***Fleas* and *Ticks (Adults and Larvae)*:** Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. Make a test application to upholstery or drapery fabrics in an inconspicuous place before use.

TO CONTROL *Fleas* and *Ticks* ON DOGS. Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application every two weeks if necessary.

Do not spray directly in/on eyes, mouth or genitalia of pets. Use only on dogs. Do not use on cats or kittens. Do not use on puppies less than twelve (12) weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals. Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

TO KILL *Lice* and *Louse Eggs*: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray, as well as *Whiteflies*, *Aphids*, *Exposed Thrips*, *Spiders*, *Red Mites* and *Leafminers*.

May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, Euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against *Whiteflies*, *Aphids*, *Spider* and *Red Mites*, *Exposed Thrips*, *Lace Bugs*, *Japanese Beetles*, *Pavement Ants*, *Clover Mites* and *Armyworms*. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Over application may cause damage. Contact insects directly whenever possible.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry area away from [inaccessible to] children and pets and away from heat, sparks and open flame. Do not transport or store below 32°F. [Protect from freezing.] [Do not freeze.]

Disposal: Do Not Puncture or Incinerate! *If empty:* Place in trash or offer for recycling if available. *If partly filled:* Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

[This container may be recycled in aerosol recycling centers. At present, there are only a few such centers in the country. Before offering for recycling, empty the can by using the product according to the label.]

4/4

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Valent BioSciences Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: VALENT BIOSCIENCES CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Valent BioSciences Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Valent BioSciences Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT VALENT BIOSCIENCES CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Valent BioSciences Corporation, 2002

Permanone is a registered trademark of Bayer AG.

Valent BioSciences Corporation
Environmental Science Division
870 Technology Way
Libertyville, IL 60048
List No. 73049306

02-0198/R1

[] Indicate alternate text in addition to any subset of the directions for use.