

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Nasser Assaf, Ph.D. Regulatory Manager, ESD Valent BioSciences Corporation 870 Technology Way Libertyville, IL 60048

SEP 2 5 2008

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Dr. Assaf:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated July 31, 2008 and September 2, 2008 for:

EPA Registration 73049-334

Permanone Insect Killer R4

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

\$EPA	United States Environmental Protection Agency Washington, DC 20460		Registra Amenda V Other		OPP Identifier Number	
	Application	n for Pesticide - Sec	tion I			
1. Company/Product Number 73049-334			2. EPA Product Manager George LaRocca		3. Proposed Classification None Restricted	
4. Company/Product (Name)	Company/Product (Name) Permanone Insect Killer R4		PM# 13			
5. Name and Address of App Valent BioSciences C 870 Technology Way Libertyville, IL 60048		(b)(i), my product to: EPA Reg. No	is similar or iden			
. Check it this	13 & NOW GUILTON	Product Name Section - II			<u> </u>	
Notification - Explain Explanation: Use addition This notification of label chan regulations at 40 CFR 156.10 Formula for this product. I ur amended label is not consist and I may be subject to enfor	below. hal page(s) if necessary. (For section age per PR Notice 2007-4. This notificate 0, 156.140, 156.144, 156.146, and 156.1 aderstand that it is a violation of 18 U.S. are twith the requirements of 40 CFR 156 are twith the requirements under secting STORAGE AND DISPOSAL PER P	Other - Exploin II.) ion is consistent with the guida. 156. No other changes have be C. Sec. 1001 to willfully make a 6.10, 156.140, 156.144. 156.14 ions 12 and 14 of FIFRA.	Application. plain below. nce in PR Notice 200 een made to the labe ny false statement to 6, and 156.156, this	eling or the (DEPA. I fur product ma	Confidential Statement of ther understand that if the y be in violation of FIFRA	
		Section - III				
1. Material This Product Wil	Be Packaged In:					
Child-Resistant Packaging Yes No Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt. Container	Water Soluble Packaging Yes No If "Yes" No. per Package wgt Contains		Container Metal Plastic Glass Paper Other (S	specify)	
3. Location of Net Contents		il Container I to 128 ounces	5. Location of La		ins	
6. Manner in Which Label is	5 3]			
		Section - IV	· · · · · · · · · · · · · · · · · · ·			
1. Contact Point Complete	items directly below for identification	of individual to be contacted	, if necessary, to p	rocess this	application.)	
Name Nasser Assaf, Ph.D.		Fitte Regulatory Manag	Regulatory Manager, ESD		Telephone No. (Include Area Code) (847) 968-4844	
	Certificat ments I have made on this form and a	all attachments thereto are tru			6. Date Application Received (Stamped)	

3. Title

5. Date

Nasser Assaf, Ph.D.

NAASSAF

both under applicable law.

2. Signature

4. Typed Name

Regulatory Manager, ESD

September 2, 2008

PERMANONE® INSECT KILLER R4

- FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS.
- FOR LIVESTOCK AND PREMISE USE
- READY-TO-USE FOR CONTROL OF HOUSEHOLD PESTS
- DILUTE FOR USE ON VEGETABLES AND FRUIT AND NUT TREES
- FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES AND ON DOGS
- · CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS
- FOR CONTROL OF PESTS ON LIVESTOCK
- WITH COUNTRY FRESH FRAGRANCE
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF COCKROACHES
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF FLEAS
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF STORED PRODUCT PESTS
- FOR HOUSEHOLD AND DOMESTIC USE ONLY
 - READY TO USE
 - WATER-BASED
 - ORNAMENTAL AND HEDGE TREATMENT
- OUTDOOR PEST CONTROL PRODUCT
- CONTROLS NUMEROUS PESTS OF ORNAMENTAL PLANTS
- CONTROL OF TICKS
- CONTAINS PERMANONE
- KILLS ON CONTACT
- FOR CONTROL OF ANTS
- FOR CONTROL OF GYPSY MOTHS
- KILLS FLEAS AND TICKS

ACTIVE INGREDIENT:

Permethrin ¹	0.25 %
OTHER INGREDIENTS:	
	100 00 %

1 Cis/trans ratio: max. 35% (+/-) cis and min. 65% (+/-) trans.

PRECAUCION AL CONSUMIDOR: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 73049-334

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Side Panel For Additional Precautions.

NET CONTENTS:

In case of *Medical* emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-877-315-9819.

READ ENTIRE LABEL BEFORE EACH USE

PET USE APPLIES TO USE ONLY ON DOGS

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING

Do not use on electrical equipment or on surfaces that can be harmed or stained by water.

INDOOR APPLICATION

HOUSEHOLD PESTS: For spot application only. Do not use as a space spray indoors.

Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SURFACE SPRAYING: FOR UP TO 60-DAY (TWO MONTHS) CONTROL OF: Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms,

Grain Mites and Cadelles: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For **Silverfish**, spray bookcases. For **Ants**, spray trails, nests and points of entry.

TO CONTROL *Carpet Beetles*: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving.

TO CONTROL **Bedbugs**: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors.

TO CONTROL *Fleas (Adults and Larvae) and Ticks (Adults and Larvae)*: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Place fresh bedding in animal quarters following treatment. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL *Clothes Moths*: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL *Lice and Louse Eggs*: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned. Allow all treated particles to dry thoroughly before use.

LIVESTOCK PESTS

Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. Do not contaminate animal feed or water during application.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from *Horn Flies, Face Flies, Houseflies, Mosquitoes* and *Gnats:* Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL **Stable Flies, Horse Flies and Deer Flies**: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL **Blood-sucking Lice**: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL **Bedbugs and Mites IN POULTRY HOUSES**: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS; TO CONTROL *Fleas, Ticks, and Lice*: Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat application every two weeks if necessary. DO NOT USE ON CATS OR KITTENS.

OUTDOOR APPLICATION

TO HELP CONTROL *Ticks* IN OUTDOOR ACTIVITY AREAS: Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. Treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

PESTS ON THE OUTDOOR SURFACE OF BUILDINGS: To temporarily control *Houseflies, Mosquitoes, Cluster Flies, Gnats* and *Flying Moths*. For outdoor use only as an aid in reducing annoyance from these insects.

Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of PERMANONE and 12 fluid ounces of water. Using dilute mixture, spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Flush sprayer with clean water after each use.

ORNAMENTAL FLOWER GARDENS

Ready to use. Affix trigger, pump or other type of sprayer to container for application. Do not transfer to another container.

For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites* and *Leafminers*. May be used on plants as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

ORNAMENTAL TREES AROUND THE HOME: Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of PERMANONE and 12 fluid ounces of water. Using dilute mixture, use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: *Pine Moths, Pine Needle Miners, Pine Beetles* and *Needle Scales* on pine: *Budworms* on conifers; *Oakworms* on oak; *Fall Webworms* on cherry, elm and honeysuckle; *Aphids* on tulip poplar; *Citrus black flies* on non-bearing citrus; *Tussock Moths* on Douglas and grand firs; *Leafminers* on birch; *Elm Leaf Beetles* and *Elm Spanworms* on elms. Flush sprayer with clean water after each use.

FOR HOME VEGETABLE GARDENS: Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of PERMANONE and 12 fluid ounces of water. Spray upper and lower leaf surface as directed. Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. Apply no more than 13 ounces of dilute mixture over 100 linear feet of row (13 ounces of diluted mixture per 100 square feet.) Flush sprayer with clean water after each use.

VEGETABLE TARGET PESTS REMARKS Asparagus Cutworms Spray plants as needed (pre-harvest) to ensure uniform asparagus beetle coverage. Do not make more than 4 applications. Can be applied up to 3 days prior to harvest. Asparagus Larval stages of Treat fern or bush asparagus beetle; (post-harvest) growth after spear tarnished plant harvest when insects bugs, lygus bugs are present. and adult Japanese beetles. Celery - see leafy vegetables Cole Crops: Beet armyworm; Apply every 5 days or Broccoli, Brussels diamond back moth; as needed. Can be Sprouts, Cabbage, imported cabbageapplied up to one day Cauliflower worm; cabbage aphid, prior to harvest. Do cabbage looper not make more than 10 applications on cabbage and 8 applications on the other cole crops. Corn - see sweet corn Cucurbits: Aphids, cabbage Spray plants as needed loopers, cucumber to ensure uniform Balsam pear(bitter coverage. Do not make melon), Chinese beetles (adults), more than 8 applicatcutworms, leafhop-, waxgourd (Chinese ions per crop. Can be melon), pers, leafminers applied up to the day melonworms, Citrus melon, cucumber, Gherkin, pickleworms, plant of harvest. Edible gourds, bugs and squash Melons including bugs. hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, and Persian melon, pumpkin, summer &

Eggplant

winter squash, watermelon

Colorado potato beetles, vegetable leaf miners Spray plants as needed to ensure uniform coverage. Do not make more than 16 applications. Can be applied up to 3 days prior to harvest.

Horseradish

Imported crucifer weevils

Spray as needed. Do not make more than 3 foliar applications. Can be applied up to 22 days prior to harvest.

Leafy Vegetables: amaranth, arrugula, celery, celtuce. chervil, corn salad, chrysanthemum (edible-leaved and garland), cress (garden and upland), dandelion, dick, endive, fennel, lettuce (head & leaf), orach, parsley, purslane (garden and winter), rhubarb, spinach, Swiss chard Melons - see Cucurbits

Alfalfa loopers, aphids, beet army-worms, cabbage loopers, corn earworms, cutworms, European corn borers, fall armyworm, green cloverworms, southern armyworms, tobacco budworms, vegetable leaf miners.

Spray every 5 to 10 days as needed. Do not make more than 7 applications per crop. Can be applied up to 1 day prior to harvest.

Potatoes

Beet armyworms,
Colorado potato
beetles, potato
beetles, potato
leafhoppers, potato
aphids, potato tuberworms,
cabbage
loopers, cutworms,
European corn borers,
potato psyllids,
tarnished plant bugs

Spray plants as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.

Peppers

Vegetable leafminers, cabbage loopers, flea beetles, corn earworms, pepper weevils, European corn borers Spray plants as needed. Do not make more than 8 applications. Can be applied up to 3 days prior to harvest.

Sweet Corn

Corn earworms, European corn borers, fall armyworms, southern armyworms Spray plants every 5 days and when needed. Do not make more than 6 applications per season. Can be applied up to 1 day prior to harvest.

Tomatoes (Do not spray on cherry tomatoes and varieties yielding tomatoes less than one inch in diameter) Beet and southern armyworms, tomato pinworms, tomato fruitworms, hornworms, granulate cutworms, Colorado potato beetles, cabbage loopers Spray plants when needed for uniform coverage. Do not make more than 6 applications per season. Can be applied up to date of harvest.

HOME OWNER GROWN FRUIT AND NUT TREES: Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of PERMANONE and 12 fluid ounces of water. Apply dilute mixture to trees until foliage is dripping wet. Make sure to spray the undersides of leaves and into dense foliage. Flush sprayer with clean water after each use.

(Fruit or Nut) Almonds	Target Pest(s) Navel orange worms, peach twig borers	Remarks Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Plum curculio, red- banded leafrollers, rosyapple aphids, spotted tentiform leafminers, tarnished bugs, white apple leafhoppers, oblique-banded leafrollers, green fuitworms.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
Peaches	Lesser peachtree borers, peach twig borers, plum curculio, plant bugs, green fruit worms, Oriental fruit moths	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears	pear psylla, green fruitworms and codling moths.	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry area away from children and pets.

Container Disposal: If empty: Non-refillable container. Do not reuse or refill this container.

Place in trash or offer for recycling if available.

Trees

If partly filled: Call your local solid waste agency 1-800-CLEANUP for disposal instructions.

Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should <u>must</u> be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Valent BioSciences Corporation. All such risks shall be assumed by the user or buyer.

VALENT BIOSCIENCES CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Valent BioSciences Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Valent BioSciences Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: <u>TO THE EXTENT CONSISTENT WITH APPLICABLE LAW</u>, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT VALENT BIOSCIENCES CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Valent BioSciences Corporation, 2008

Permanone is a registered trademark of Bayer AG.

Valent BioSciences Corporation Environmental Science Division 870 Technology Way Libertyville, IL 60048 List No. 73049334

02-0267/R1