
	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Avenue, N.W. Washington, D.C. 20004	EPA Reg. Number:	Date of Issuance:
		73049- 419	MAY 16 2005
		Term of Issuance: Conditional	
(under FIFRA, as amended)		Pramex TC Plus	
Name and Address of Registrant (include ZIP Code): Valent BioSciences Corporation 870 Technology Way, Suite 100 Libertyville, IL 60048			
<b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.			
On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.  Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.  This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:  1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.  2. Make the labeling changes listed below before you release the product for shipment:  a. Add the phrase "EPA Registration No. 73049-419."  b. Delete the following claims since no data was submitted to support them, "Keeps Termites Away for Up to 5 years on treated areas", "Long Lasting Termite Control Will Not Be Washed Away By The Rain", "Helps reduce flying insects such as mosquitoes which can cause West Nile virus and equine encephalitis, and "Kills deer ticks that may carry Lyme disease."			
Signature of Approving Official: 		Date: May 16, 2005	

c. Revise the precautionary statements as follows: "Causes skin irritation. Do not get on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. causes moderate eye irritation. Avoid contact with eyes. Harmful if swallowed or inhaled. Avoid breathing vapor or spray mist.".

d. Reverse the order of the First Aid statements.

e. Under Directions for use add "Shake Well before Use".

f. Under Subterranean termites delete "and prevent reinfestations" and revise the last statement to read, "A complete termite barrier may require treatment to the inside perimeter of the foundation and to other critical interior areas. For these areas and if termites are found inside of the home consult a licenced pest control operator for treatment.".

g. Add the following to the Note under Subterranean Termites "When treating adjacent structure, check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structures. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is complete.".

h. Under Wood Destroying Insects add the following sentence to the end of the second paragraph, "For these types of application consult a licenced pest control operator for treatment. Also add "Outdoor Use" following the heading Wood Destroying Insects.

i. Change "Controls many garden insect pests" to "Controls many named garden insect pests".

j. Change "Controls many ornamental insect pests" to "Controls many named ornamental insect pests"

3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

MAY 16 2005

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
73049-419

**Pramex® TC Plus**

- FOR OUTDOOR RESIDENTIAL/HOMEOWNER USE ONLY
- [LOW ODOR FORMULA
- EASY TO APPLY
- KILLS TERMITES ON CONTACT
- FOR USE ON AND AROUND BUILDINGS, FOUNDATIONS, PORCHES, WOOD PILES, WOOD FENCES, DECKS & PATIOS
- CONTROLS/KILLS WOOD DESTROYING INSECTS INCLUDING TERMITES, CARPENTER ANTS, CARPENTER BEES, POWDER POST BEETLES AND OTHER LISTED PESTS
- KEEPS TERMITES AWAY FOR UP TO 5 YEARS IN TREATED AREAS
- IT'S EASY TO PROTECT YOUR HOME FROM HOME, LAWN AND GARDEN DAMAGING INSECTS
- CONTROLS HARD-TO-KILL INSECTS FOR UP TO 4 WEEKS
- CREATES AN INSECT BARRIER AROUND YOUR YARD FOR FOUR WEEKS
- LONG LASTING TERMITE CONTROL WILL NOT BE WASHED AWAY BY THE RAIN
- ECONOMICAL - FOR TERMITE CONTROL, 1 QUART MAKES UP TO 5 GALLONS
- ECONOMICAL - FOR TERMITE CONTROL, 1 QUART TREATS UP TO 25 LINEAR FEET
- CONTAINS PRAMEX BRAND PERMETHRIN - A LONG LASTING PYRETHROID INSECTICIDE WHICH PROVIDES CONTACT AND RESIDUAL CONTROL]

**ACTIVE INGREDIENT:**

Permethrin\* ..... 10.0 %  
OTHER INGREDIENTS\*\*..... 90.0 %  
100.0 %

\* Cis/trans ratio: max. 55% (+/-) cis and min. 45% (+/-) trans.

\*\*Contains Petroleum Distillates

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA Reg. No. 73049-

EPA Est. No.

**KEEP OUT OF THE REACH OF CHILDREN  
WARNING**

See Side Panel for Additional Precautions

FIRST AID	
If Swallowed:	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give any liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If Inhaled:	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
If In Eyes:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If On Skin or Clothing:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-877-315-9819.	
NOTE TO PHYSICIAN	
Contains Petroleum Distillate - vomiting may cause aspiration pneumonia.	

**NET CONTENTS:** 1 gallon (3.79 l)

## PRECAUTIONARY STATEMENTS

### Hazards To Humans & Domestic Animals

#### WARNING

Harmful if swallowed or inhaled. Causes skin irritation and moderate eye injury. Do not get on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes. Avoid breathing vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow drift to blooming crops or weeds if bees are visiting the treatment area.

#### Physical or Chemical Hazards

Do not use or store near heat or open flame. Do not apply this product in or on electrical equipment due to possibility of shock hazards.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes. For outdoor use only, do not use in homes. Do not apply to pets.

This is an easy-to-apply product that can be used to control home-invading, wood-destroying, and lawn and garden insects. Apply by using a pump-up pressure sprayer, a hose-end sprayer, or watering can. Always use a clean sprayer. Carefully measure and mix the amount of product with water as directed. Use only in ventilated areas.

When mixing or applying, wear long sleeved shirt, long pants, socks, shoes and chemical resistant gloves. Do not use food utensils or cups for measuring. Spray with the wind to your back. Do not spray on windy days. Do not allow spray to contact food, food utensils or water supplies. Do not allow children or pets to go onto treated area until the treated areas such as grass or soil has dried.

#### HOW TO MIX AND APPLY

Mix the recommended amount of *Pramex TC Plus* per gallon of water and apply in accordance with the directions below. Where appropriate, measure the area to be treated by measuring the length and width then multiply to determine the square footage of the area. Apply *Pramex TC Plus* as a coarse spray with a pump-up, hose-end or other type of hand-held sprayer. When mixing, fill the sprayer with half of the water first and then measure and add the appropriate amount of *Pramex TC Plus* followed by the remaining water. Thorough spray coverage is necessary, but avoid spray run-off.

#### MEASUREMENT CONVERSION CHART.

1 teaspoon = 0.17 fl.oz. = 5 ml.  
1 ounce = 6 teaspoons = 29.6 ml.  
1 gallon = 128 ounces = 3.79 liters

#### SUBTERRANEAN TERMITES:

- For outdoor, homeowner post-construction use only.
- If termites are found within the house or structure, contact a licensed pest control operator for treatment.
- DO NOT use indoors, under slabs or in crawl spaces.
- Do not spray on firewood.

Kills and controls existing subterranean termites and prevents reinfestations. The active ingredient in this product forms a barrier in the soil that prevents termite attack. Diluted *Pramex TC Plus* [product] must be adequately dispersed in the soil to provide a barrier between the wood in the structure and the termite colonies in the soil. Barriers may be established in the soil on the outside perimeter of the house by trenching using a rate of 1 gallon of dilution per 5 linear feet per foot of depth. A complete termite barrier may require treatment to the inside perimeter of the foundation and to other critical interior areas by a licensed pest control operator.

#### Trench Treatment:

**Mixing Instructions:** Add 6.5 fl. oz. of *Pramex TC Plus* [product] to 1 gallon of water (0.5%) for each 5 linear feet [5 feet of trench length]. One quart of *Pramex TC Plus* makes 5 gallons of diluted material and treats 25 linear feet.

- Remove and discard all wood debris while digging trench.
- Poured concrete foundations - Dig a trench next to the foundation 4 inches deep and 6 inches wide or down to the top of the footing, whichever is less.
- Brick & block foundations - Dig a trench 12 inches deep and 6 inches wide. If the footing is more than 12 inches deep, make holes with a crowbar, pipe or steel rod about 1 foot apart that extend from the bottom of the trench to the top of the footing. Make holes closer in hard packed clay soils.
- To avoid shifting your house foundation, never dig below the top of the footing.
- Apply half the volume of diluted product with a sprinkling can (or pump up sprayer) along the length of the trench. Refill trench with soil mixing it with the remaining [other half of] diluted product.

Reapplication for subterranean termites should only be made when there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping, etc. or if termites reappear. Reapplication should be made as a spot treatment to those areas. Annual reapplication of entire premises must be avoided.

#### NOTES:

1. Do not treat soil in the vicinity of wells, cisterns, ponds or other bodies of water. For these situations, contact your County Cooperative Extension Agent or consult a commercial pest control service.
2. Do not apply product until locations of wells and utility lines (gas, water, sewer, cable and electric) are known and identified. Be careful not to apply product into wells or puncture utility lines.
3. Do not treat soil that is water saturated or frozen, or in any conditions where run-off or movement from the treatment area is likely to occur.
4. Do not apply while it is raining.

Desired amount of linear feet to treat	Final Concentration	Amount of Pramex TC Plus	Finished amount of diluted product for treatment
5	0.5%	6.5 fl. oz.	1 gallon
10	0.5%	13 fl. oz.	2 gallons
25	0.5%	1 quart	5 gallons
50	0.5%	2 quarts	10 gallons
100	0.5%	1 gallon	20 gallons

#### WOOD-DESTROYING INSECTS:

**Mixing Instructions:** Mix 6.5 fl. oz. of Pramex TC PLUS per gallon of water (0.5%) to control above ground *Termites* (*Drywood*, *Formosan*), *Carpenter Ants*, *Carpenter Bees*, *Powder Post Beetles*, *False Powder Post Beetles*, *Deathwatch Beetles*, *Old House Borer* and *Ambrosia Beetles*.

Treat localized areas of infested wood. Spray or brush into areas where insects are suspected or known to be, for example holes, spaces between wooden parts of the structure, the space between different types of building materials, such as concrete and wood. Also spray exposed insects. This type of application is not intended to be a substitute for soil treatment mechanical alteration or fumigation to control extensive infestation of wood-infesting insects.

**Fences and Outdoor Wood Surfaces of Buildings:** Apply evenly on wood surfaces. A paintbrush (or pump up sprayer) may be used to ensure uniform application. Allow to dry then repeat to ensure the insecticide soaks into the wood.

**Wooden Posts (fence, deck, mailbox, etc):** Using a sprinkling can, pour ½ gallon of diluted product into the fence post hole and while refilling hole, add another ½ gallon to the fill soil so that all the soil is treated to form a protective barrier.

**Wood Piles and Stored Lumber:** Use a hose-end, pump sprayer or sprinkling can to deliver a coarse drenching spray to the soil beneath and around wood storage areas prior to stacking. Do not treat firewood directly.

**For Carpenter Bees :** Use a suitable sealant to fill carpenter bee holes when there is no evidence of bee activity, usually 24 hours after treating holes.

#### HOSE-END SPRAYER USE DIRECTIONS FOR ORNAMENTAL AND LAWN PEST CONTROL:

Pramex TC Plus can be used in hose end sprayers. Follow instructions on the hose end sprayer adjusting the dial to the desired application rate per gallon.

1. Make sure the hose-end sprayer is firmly connected to the hose.
2. Make sure the sprayer is in the OFF position. Turn water supply on at the faucet. For best results, turn hose faucet to maximum (25-65 psi). Water pressure will build in the hose. To avoid damaging delicate plants water pressure may need to be decreased. Water pressure should be enough to deliver spray in a fan pattern.
3. To apply product, turn the control valve to the ON position. Water and product are now being mixed at the proper rate.
4. Begin spray at farthest section of area to be treated and carefully move backwards avoiding sprayed areas.
5. Spray with wind at your back. Do not spray on windy days.
6. Spray until surface is slightly wet but not to the point of run-off. Over application may cause damage.

7/11

## ORNAMENTAL PLANTS

When using a watering can, pump up or similar type of sprayer, mix *Pramex TC Plus* with water in the following manner:

<i>Pramex TC Plus</i>	Water	Percent
1 ½ teaspoon (7 ml)	1 gallon (3.79 L)	0.02

NOTE: Do not use more *Pramex TC Plus* than recommended, as this may cause some plant injury. Use this dilution unless other instructions are given.

CONTACT AND RESIDUAL SPRAY: Apply a diluted spray of *Pramex TC Plus*. Use sufficient spray to obtain full coverage of all foliage.

HOUSE PLANTS: To protect plants in indoor landscaping, spray directly on insects to kill *Leafrollers*, *Armyworms*, *Cutworms*, *Cabbage Worms* on chrysanthemums; *Fungus Gnats* on velvet plants. Spray as directed above to control *Aphids*, *Armyworms*, *Japanese Beetles*, *Leafminers*, *Leafrollers*, *Mealybugs*, *Scales*, *Spider Mites*, *Spring Cankerworms*, *Exposed Thrips* and *Whiteflies* on the following plants:

Ageratum	English Ivy	Poinsettia
Ardisia	Fuchsia	Portulaca
Aster	Gladiolus	Pothos
Azalea	Grape Ivy	Prayer Plant
Baby's breath	Hypoestes	Purple Passion
Begonia	Ivy	Rabbits foot fern
Birds-nest fern	Juniper	Rose
Bleeding heart	Lily	Schefflera
Carnation	Marigold	Snapdragon
Chrysanthemum	Orchid	Snake Plant
Coleus	Palm	Statice
Crown of Thoms	Pansy	Velvet Plant
Cockscomb	Piperomia	Verbena
Cyclamen	Petunia	Zinnia
Dracaena	Philodendron	
Dumbcane	Piggy-back plant	

Remove infested plants from living areas prior to application. Spray plants outside in an area not likely to be occupied by children or pets (particularly fish). When leaves dry, plants can be returned to original location.

Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours and check for possible injury.

Repeat applications may be necessary to control *Citrus Blackflies* on non-bearing citrus; *Adult Japanese beetles* on roses.

OUTDOOR ORNAMENTALS: Use 1 ½ teaspoon (7 ml) of *Pramex TC Plus* in one gallon of water. (0.02%)

CONTACT AND RESIDUAL SPRAY: Apply a finished spray of *Pramex TC Plus*. Use sufficient spray to obtain full coverage of all foliage. Spray in the late afternoon or evening when the temperature ranges from 50 F to 85 F and when there is little or no wind. Spray at the first indication of insect pests. Repeat treatment at intervals of 4 to 8 days if necessary. *Pramex TC Plus* application can be made up to the day of collection.

Spray as directed above to control *Aphids*, *Cicadas*, *Corn Earworms*, *Fall Cankerworms*, *Inchworms*, *Leafminers*, *Leafrollers*, *Mealybugs*, *Bagworms*, *Spider Mites*, *Exposed Thrips* and *Whiteflies* on the following plants:

Ageratum	Gladiolus	Pea shrub
Arizona Cypress	Gold bells	Petunia
Aster	Hypoestes	Philodendron
Azalea	Ivy	Poinsettia
Begonia	Lilac	Rhododendron
Coleus	Marigold	Rose
Common ninebark	Mock Orange	Snapdragon
Common Snowberry	Nannyberry	Taxus
Elm	Orchid	Wax plants
Exacum	Palm	Weeping fig
Euonymous	Pans	Zinnia
English Ivy		

**FOR ORNAMENTAL TREES GROWN AROUND THE HOME:** Use sufficient spray to obtain full coverage of all foliage. Avoid heavy wetting. Use to control: pine moths, pine needle miners, pine beetles and needle scales on pine; budworms on conifers; oakworms on oak; fall webworms on cherry, elm and honeysuckle; aphids on tulip poplar; citrus black flies on non-bearing citrus; tussock moths on Douglas and grand firs; leafminers on birch; elm leaf beetle and elm spanworms on elms.

#### NUISANCE PESTS IN OUTDOOR AREAS:

Pest	Dilution	Directions
Ants, crickets, earwigs, fleas, ticks (lone star ticks, deer ticks, black legged ticks)	Mix 6.5 fl oz (192 ml) of Pramex TC Plus (this product) in one gallon of water (3.79 L) (1 ½ fl oz (49 ml) per quart (0.95 L) (0.5%)	For localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur but no more than once per week.
Tent Caterpillars	Mix 2.5 fl oz (74 ml) of Pramex TC Plus (this product) in one gallon (3.79 L) of water. (0.2%)	Apply when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents on all sides. A few caterpillars may be seen escaping from the tent, however, they should die within several hours.

#### NUISANCE PESTS ON OUTDOOR SURFACES OF BUILDINGS:

Ants, clover mites, crickets, earwigs, millipedes, sowbugs (pillbugs)	Mix 1 ½ fl. oz. (8.35 ml) in 1 quart of water (0.5%) (Mix 6.5 fl oz (192 ml) of Pramex TC Plus (this product) in one gallon of water (3.79 L) )	To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, patios, garages and other areas where these pests may be found. Repeat applications as reinfestations occur but no more than once per week.
Flies (cluster, house), gnats, flying moths, mosquitoes (including mosquitoes which may carry West Nile Virus or equine encephalitis)	Mix 1 ½ fl. oz. (8.35 ml) in 1 quart of water (0.5%) (Mix 6.5 fl oz (192 ml) of Pramex TC Plus (this product) in one gallon of water (3.79 L) )	As an aid in reducing annoyance from these insects, spray outside surfaces of screens, doors, window frames or wherever these insects may enter the home. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight. Repeat application as necessary but no more than once per week.

#### FOR USE ON LAWNS

For control of *Ants (including fire ants when mound is treated)*, *Armyworms*, *Chinch bugs*, *Crickets*, *Cutworms*, *Japanese Beetle Grubs*, *Grasshoppers*, *Mole Crickets* and *Sod Webworms*, consult the following chart:

PESTS	DILUTION	REMARKS
Armyworms                      Mole crickets Cutworms                      Sod webworm Chinch bugs Japanese beetle grubs	1.2 fl. oz. (7 tsp)/10 gal water/1000 ft <sup>2</sup> (35 ml/38 L water/93m <sup>2</sup> ) (0.009%)	For surface feeders (armyworms, cutworms, chinch bugs, sod webworms), delay watering or mowing for 24 hours. For subsurface feeders (Japanese beetle grubs, mole crickets), irrigate with ½ inch of water immediately after application
Crickets Grasshoppers Ants (including Fire Ants)	3.5 fl. oz. (3 tsp) /3 gal water/1000 ft <sup>2</sup> (15 ml/11.4 L water/93m <sup>2</sup> ) (0.013%)	For best results, delay watering or mowing for 24 hours after treatment. (For fire ants, only aids in control)



9/11

Lawns should be no taller than 3 inches (7.6 cm) at time of application. Slightly more water may be used as long as the amount of product per area is as listed above. For heavy infestations, repeat application after two weeks.

Also for control of *cutworms* on dichondra and Irish moss. Use as directed above for lawns.

#### FOR APPLICATION ON VEGETABLES (For Backyard Vegetable Gardens Only):

Prepare Pramex TC Plus as directed in the following chart. Use 0.25 gallons of finished spray per 110 square feet or one liter per 10.7 square meters of surface area.

Vegetable	Target Pest(s)	Remarks	Recommended Pramex TC Plus Dilution Use Rate Finished Spray			
			tsp/gal 1	ml/gal	ml/liter	percent
Asparagus (pre-harvest)	Cutworm, Asparagus Beetle	Apply for uniform coverage as needed. Do not apply more than 4 applications per crop. Can be applied up to 3 day prior to harvest.	1	5	1 ½	0.015
(post-harvest)	Larval stages of Asparagus Beetle, Tarnished Plant Bug, Lygus Bugs, and Adult Japanese Beetle	Treat ferns or bush growth after spear harvest when insects are present.	1	5	1 ½	0.015
Celery	Vegetable Leaf Miner, Cabbage Looper, Black and Granulate Cutworm, Fall, Beet and Southern Armyworm	Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.	2	10	2 ¾	0.03
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet Armyworm, Diamond Back Moth, Imported Cabbageworm, Cabbage aphid, Cabbage Looper	Apply every 5 days or as needed. Can be applied up to one day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage.	1	5	1 ½	0.015
Eggplant	Colorado Potato Beetle, Vegetable Leaf Miner	Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 prior to harvest.	2	10	2 ¾	0.03
Horseradish (foliar application)	Imported Crucifer Weevil	Apply as needed. No not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.	2	10	2 ¾	0.03
Head Lettuce only (Do not us on leaf lettuce)	Cabbage Looper, Alfalfa Looper, Beet Armyworm, Corn Ear Worm and Tobacco Budworm	Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.	2	10	2 ¾	0.03
Potatoes	Beet Armyworm, Colorado Potato Beetle, Potato Leafhopper, Potato Flea Beetle, Aster Leafhopper, Potato Aphid, Potato Tuberworm, Cabbage Looper, Cutworms, European Corn Borer, Potato Psyllid, Tarnished Plant Bug	Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 day prior to harvest	2	10	2 ¾	0.03

Vegetable	Target Pest(s)	Remarks	Recommended Pramex TC Plus Dilution Use Rate Finished Spray			
			tsp/gal	ml/gal	ml/liter	percent
Peppers	Vegetable Leafminer, Cabbage Looper, Fleabeetle, Corn Earworm, Pepper Weevil, European Corn Borer	Apply as needed, but not more than 8 sprays per season. Can be applied up to 3 days prior to harvest	2	1	2 ¾	0.03
Spinach	Beet Armyworm, Corn Earworm, Cabbage Looper, Cutworms, European Corn Borer, Green Cloverworm, and Leafhoppers	Apply for uniform coverage when needed but not more than 7 applications per season, can be applied up to 7 days prior to harvest.	1	5	1 ½	0.015
Sweet Corn	Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm	Apply every 5 days and when needed but not more than 6 applications per season. Can be applied up to 1 day prior to harvest	2	10	2 ¾	0.03
Tomatoes (Do not apply on cherry tomatoes and varieties yielding less than tomatoes one inch in diameter.)	Beet and Southern Armyworm, Tomato Pinworm Tomato Fruitworm, Horn Worms, Granulate cutworms, Colorado Potato Beetle, Cabbage Looper	Apply for uniform coverage when needed but not more than 6 applications per season. Can be applied up to date of harvest.	2	10	2 ¾	0.03

**FOR TREE FRUIT AND NUT CROPS GROWN AROUND THE HOME** (Homeowner-Backyard use only):

Apply finished spray to trees until foliage is dripping wet.

Trees (Fruit or Nut)	Target Pest(s)	Remarks	Recommended Pramex TC Plus Dilution Use Rate Finished Spray			
			tsp/gal	ml/gal	ml/liter	percent
Almonds	Navel Orange Worm, Peach Twig Borer	Apply when insects appear. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.	3	15	4	0.04
Apples	Plum Curculio, Redbanded Leafroller, Rosyapple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique-banded Leafroller, Green Fruitworm	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.	3	15	4	0.04
Peaches	Lesser Peachtree Borer, Peach Twig Borer, Plum Curculio, Tarnished Plant Bug, Green Fruit Worm, Oriental Fruit Moth	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.	3	15	4	0.04
Pears(dormant through delayed dormant)	Pear Psylla	Apply during the dormant through delayed dormant growth period only. Add if desired, 8 oz. of spray oil to each gallon of Pramex TC Plus. Do not make more than 2 applications per season.	3	15	4	0.04
(summer)	Pear Psylla, Green Fruitworm, and Codling Moth	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.	3	15	4	0.04

#### STORAGE & DISPOSAL

**Storage:** Store in a cool, dry area away from [inaccessible to] children and pets. Do not transport or store below 32°F. [Protect from freezing.] [Do not freeze.]

**Disposal:** If empty: Do not reuse this container. Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions of Sale and Limitation of Warranty and Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions of Sale and Limitation of Warranty and Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Valent BioSciences Corporation. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** VALENT BIOSCIENCES CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Valent BioSciences Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Any warranties having been made, express or implied, are not applicable if this product has been used contrary to label directions, under abnormal conditions, or under conditions not reasonably foreseeable to Valent BioSciences Corporation, and buyer assumes the risk of any such use to the extent permitted by law. Valent BioSciences Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT VALENT BIOSCIENCES CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

This Conditions of Sale and Limitations of Warranty and Liability may not be amended by any oral or written agreement.

©Valent BioSciences Corporation, 2004

Valent BioSciences Corporation  
Environmental Science Division  
870 Technology Way  
Libertyville, IL 60048  
List No. 73049364

02-0177/R1

#### [Alternate Bullet Points:

1. Vegetable, Lawn and Garden Spray Concentrate
2. It's the least you can do for your plants
3. For use on plants
4. For use outdoors around building, homes, in small flower and vegetable gardens and on fruits, nuts and ornamental trees
5. Economical, 8 oz. makes up to 48 gallons
6. Controls/kills home-invading pests
7. Controls/kills garden pests
8. Helps reduce flying insects such as mosquitoes which can cause West Nile virus and Equine Encephalitis
9. Kills deer ticks that may transmit Lyme disease
10. Use on Lawns, gardens and fruit trees

11. Safe on over 60 varieties of plants
12. Controls many garden insect pests
13. Controls many ornamental insect pests
14. Plant pest control
15. Vegetable insect pest control
16. Effective control of hard to kill insect pests
17. Multi-Purpose insect killer
18. Works fast
19. Residual action
20. Long-lasting
21. Quick-acting
22. Kills fast-proven to last
23. You control the pest, we control the spray]