TATES ENVIRONMENTAL PROTECTIO



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Doina Bujor Regulatory Manager Valent BioSciences Corporation 870 Technology Way Libertyville, IL 60048

RE: Product Name: Teknar® HP - Biological Larvicide Aqueous Suspension

EPA Reg. No: 73049-435

Application for Notification Dated: January 25, 2010 to add Alternate Brand Name: VBC-60071 – Biological Larvicide Aqueous Suspension per PR Notice 98-10.

Dear Ms. Bujor,

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticides Registration Notice (PRN) 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PRN 98-10 and it has been determined that the action(s) requested falls within the scope of PRN 98-10. Our records have been duly noted, and the label submitted with this application has been stamped as "Notification, received and accepted" and will be placed accordingly in our records.

Questions concerning this action should be directed to Mary Paden (703) 308-0411 or email at paden mary@epa.gov.

Sincerely,

Sheryl K. Reilly

Sheryl K. Reilly, Ph.D., Chief Microbial Pesticides Branch Biopesticides and Pollution Prevention Division (7511P)

MAY 1 3 2010

CONCURRENCES								
SYMBOL	7511(P)				· · · · · · · · · · · · · · · · · · ·			
SURNAME	Paden .							
DATE.	13May 10.							

Please read instructions on	reverse before ca	ting form.	Form Apr	pro\	OMB No. :	2070-0060). Approvel expires 2-28-95
≎EPA	Environmenta	United States I Protection Ington, DC 204	<u> </u>		Registra Amendr Other		OPP Identifier Number
		Application	on for Pesticide - Sec	tion I			
1. Company/Product Numb 73049-435	er		2. EPA Product Man Sheryl Reilly	nager		3. Pro	posed Classification
4. Company/Product (Name Teknar HP	1)		PM# 90/Microbial Pes	sticides	s Branch/E	3PP	· ·
5. Name and Address of Ap Valent BioSciences 870 Technology Way Libertyville, IL 60048	Corporation y	nde)	, , , , , , , , , , , , , , , , , , ,	is simil	lar or ident	ical in cor	FIFRA Section 3(c)(3) mposition and labeling accepted
			Section - II	F	Reviewer:	mG	Maden
Notification - Explain Explanation: Use addition NOTIFICATION to add an a This notification is consisted labeling or the confidential sepa. I further understand the second sec	n below. onal page(s) if necessar alternate brand. See atta nt with the provisions of f statement of formula of th	ry. (For section ached cover letted PR Notice 98-10 this product. I unot consistent with the section of the sec	Agency let "Me Too" Other - Exp	Applica Applica plain bel stails. 152-46, 18 U.S.O	tion. low. , and no othe C. Sec 1001	er changes to willfully r	nake any false statement to
	<u> </u>		Section - III				· · · · · · · · · · · · · · · · · · ·
1. Material This Product W	ill Be Packaged In:						
Child-Resistant Packaging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt.	No. per container	Water Soluble Packaging Yes No If "Yes" No. per Package wgt Containe	er .	2. Type of	Container Metal Plastic Glass Paper Other (S	pecify)
3. Location of Net Contents	Information	4. Size(s) Ret	tail Container	5. Loc	ation of Lab	el Directio	ns
6. Manner in Which Label is	Container s Affixed to Product	Lithog Paper Stenci	glued iled	<u> </u>	<u> </u>		
		 	Section - IV				· · · · · · · · · · · · · · · · · · ·
1. Contact Point (Complete	e items directly below t	for identificatio	on of individual to be contacted,	if nece	ssary, to pro	ocess this	application.)
Name Maria Pilar Herrero			Title Regulatory Affairs Manager	·		Telephone (847) 968-	No. (Include Area Code)
	ny knowlinglly false or		ation all attachments thereto are true atement may be punishable by fi		ာ rate and con		6. Date Application Received 3 (Stamped)
2. Signature	3		3. Title Regulatory Affairs Manager			0 9 0 0 0 0 0 0 0	a 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
4. Typed Name			5. Date				စီစစ်စုစ်စ ၁
Maria Pilar Herrero			January 25, 2010				

	FIRST AID
If inhaled	 Move person to fresh air.
	 If person is not breathing, call 911 or an
	ambulance, then give artificial respiration,
	preferably by mouth to mouth, if possible.
	 Call a poison control center or doctor for
	further treatment advice.
If on skin	 Take off contaminated clothing.
or clothing	 Rinse skin immediately with plenty of water
	of 15-20 minutes.
	 Call a poison control center or doctor for
	treatment advice.

Have the product container label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-315-9819 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597. Have the product c poison control cente

HONARY STATEMENTS S TO HUMANS (AND DOMESTIC ANIMALS)

Harmful if inhaled or absorbed through the skin. Avoid betafuling space, must, Avoid contact with skin, eyes, or foldhing. Wash throughly with soap and water after handling. Bemove contaminated olothing and wash contaminated before reuse.

Mixer/loaders and applicators not in enclosed cabs or aircraft must wear a dust/mist filtering espirator meeting NUGSH standards of at least N+95, R+95 or P-95. Repeated exposure to high concentrations of microbial proteins can cause altergic sensitization.

PHYSICAL AND CHÉMICAL HAZARDS

Diluted or undiluted VBC-60071 can cause corrosion if left in proflonged contact with aluminum spray system components. Binse spray system with plenty of clean water after use. Care should be taken to prevent contact with aluminum ancraft artises, structural components and control systems. In case of contact, rinse thoroughly with plenty of water. Inspect aluminum aircraft component regularly for signs of corrosion.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply directly to treated firitshed drinking water reservoirs or drinking water receptacles when water is intended for human consumption.

avoid spotting.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction from any equipment and-weather related factors determine the potential for spray drift. The applicator and the treatment coordinator are exponsible for considering all these factors view misking decisions.

Do not apply this product through any impation system unless the labeling on chemigation is followed.

Storage. Store is a cool, less than 86° (36°0) (Joy place. Store is a cool, less than 86°° (36°0) (Joy place. Pestitide Disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal statify.

Container Disposal. Womefillable comtainer. Do not reuse or retill five container. Clean container promptly after emptying. Triple into so collainer (Cast container promptly after emptying. Triple into se so collainer (or equivalent) promptly after emptying. Triple into a so foliaise. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 14 4 tall with water and reaze, Shake for 10 seconds. Pour inseate into application equipment or a mix tank or store missale for later use or disposal part in 10 seconds. Store inseate into application equipment or a mix tank or store missale for later use or disposale this procedure two more times. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, it allowed by state and local authorities, by burning. If burning, 18 burning. If the land is the purpose of the state and local authorities, by burning.

APPLICATION DIRECTIONS
Do not apply when wind speed favors drift beyond the area of treatment.

	Suggested
losquito Habitat	Rate Range*
Such as the following examples):	0.25 - 2 pts/acre
rigation ditches, roadside	
tches, flood water, standing	
pols, woodland pools, snow	
left pools, pastures, catch basins,	
orm water retention areas, tidal	
ater, salt marshes and rice fields.	

In addition, standing water containing mosquito larvae, in fields growing crops such as: Alfalfa, almonds, asparagus, corn, cotton, dates, grapes, peaches and walnuts, may be treated at the

When applying this product to standing water containing mosquito larvae in fields growing crops, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

1 - 2 pts/acre		
'olluted water	such as sewage lagoons,	nimal waste lagoons)

*Use higher rate range in polluted water and when late 3rd and early 4th instar larvae predominate, mosquito populations are high, water is heavily polluted, and/or algae are abundant.

Rate Range	
Blackflies Habitat	Streams

Streams	
Stream water** (=ppm) or 1 minute exposure time.	0.5 - 25 mg/liter
Stream water** (=ppm) or 10 minutes exposure time.	0.05 - 2.5 mg/liter

**Use higher rate range when concentration of organic materials,

**Discharge is a principal factor determining carry of Bit. Use higher rate or increase volume by water dilution in low discharge rivers or streams under low volume (drought) conditions.

00000-04-01

Biological Larvicide

- Aqueous Suspension

	5.6%	OTHER INGREDIENTS	TOTAL100.0%
		•	•
	- :	:	:
	- :	- :	- :
_			
- >⇒	-		•
- 22		•	•
77	ŝ	- :	:
- 0,	.⊆	٠.	
.⊆	~		
, ra	_		
₽	_	•	
S	್ಷ	•	
16		- :	- :
-74	.2		
~ 52	æ		
ē,	S	•	
765	Ö	•	•
ल	.=	- :	•
- 55	_		
	\simeq		
_	₩		
- 55	- :-		•
ĕ	33	•	•
=	=	•	•
Ü	9	- :	
	. ∺	ഗ	- :
= .53		Ε.	
≥ છ	က်	2	
□ ₩	Ö	ш	•
5∵≋	=	$\overline{}$	•
≃ם	, 12	н.	:
≂∵≂	-	≂	
5 5	Ξ.	77	
ラミ	.2	¥	
= ==	ᇴ	=	
пõ	≠.	~	
ACTIVE INGREDIENT: Bacillus thuringiansis, subsp. israelensis, strain SA3A	fermentation solids, spores and insecticidal toxins	-	ઝ
= 📻	Ĕ	=	2
2 %	=	=	_
ron	تو		≃
		_	

Potency: 1200 International toxic units (ITU) per mg (Equivalent to 4.84 billion ITU per gallon, 1.279 billion ITU per fiter)

The percent active ingradient does not indicate product performance and potency measurements are not Federally standardized.

KEEP OUT OF REACH OF CHILDREN

For MEDICAL and TRANSPORT Emergencies ONLY Call 24 Hours A Day 1-877-315-9819. For All Other Information Call 1-800-323-9597.

Notification Accepted

1/3/ Date: 05/

Reviewer: 177

WALENT BIOSCIENCES. 870 Technology Way Libertyville, IL 60048

EPA Reg. No. 73049-435 EPA Est. No. 33762-1A-001

Net Contents:

(NUISANCE FLIES)

For control of nuisance tiles (Psychoda spp., Chironomus spp.) in sewage treatment facilities utilizing trickling filter systems.

APPLICATION DIRECTIONS

Suggested Rate Range*	10-20 mg/liter a (0.833-1.67 ml) per liter of wastewater feed to the filter per 30 minutes.
Nuisance Fly Habitat	Trickling filter system of wastewater treatment plants

*Use high rate for control of *Chitonomus* spp. Apply undiluted with pre-salibrated pump or other device into the wastewater feeding into the filters for a period of 30 minutes. Repeat applications as needed after 2-4 weeks. Control of *Chitonomus* spp. may take up to 2 weeks.

(NUISANCE AQUATIC MIDGES)

For control of Chironomine midges (Chironominae: Chironomini) inhabiting shallow, man-made and natural lakes or ponds.

APPLICATION DIRECTIONS

	ondhesten
Nuisance Midge Habitat	Rate Range*
Shallow Lakes and Ponds per	1 gallon (3,785,5 ml)
sewage oxidation ponds (less than	per acre
acre 6 feet deep)	

*Apply diluted with water in total volume of 5 gallons/acre by plouning or spraying over the surface to be treated with pre-aditivated device. Repeat application as needed after £ 4 weeks. Control of *Chironomine* midges may take up to

GROUND AND AERIAL APPLICATION

VBC-60071 may be applied in conventional ground or aerial application equiplent with tenditible of water sufficient to provide uniform coverage of the target area. The amount of water will depend on weather, spray equipment, and mosquito habitat characteristics. Do not mix more VBC-60071 than can be used in a 72 hour period.

For most ground spraying, apply in 5 - 100 gallons of water per acre using hand pump, airblast, mist blower, etc., spray equipment.

For aerial application, VBC-60071 may be applied either undiuted or diluted with varies. For undiluted applications, apply 0.25 to 2.0 pts. Jacre of VBC-60071 through fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or rotary atomizers.

For diluted application, fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or hydraulic agitation to provide moderate circulation before adding the NBC-5007. VBC-5007 suspends readily in water and will stay suspended over normal application periods. Brief recirculation may be necessary if the supay mixture has sat for several hours or longer. ANOID CONTINUOUS AGITATION OF

Rinse and flush spray equipment thoroughly following each

For blackfly aerial applications, VBC-60071 can be applied untilituded val fixed wing or helicopter arriard equipped with either conventional boom and nozabe systems or open pipes. Rate of application will be determined by the stream discripange and the required amount of VBC-60071 necessary to manifain a 15.–25 ppm concentration in the stream water. VBC-60071 can also be applied diluted with similar spray equipment. Do not mix more VBC-60071 than can be used in a 72 hour period.

SMALL QUANTITY DILUTION RATES Gallons Spray Mixture/Acre (Ounces Needed per Gallon of Spray)

	50 Gal/A	80.0	0.16	0.32	780
	25 Gal/A	0.16	0.32	0.64	1 28
	10 Gal/A	0.4	0.8	9.	33
VBC-60071 Rate in Pints	Per Acre	0.25 (4 oz)	0.5 (8 oz)	1.0 (16 oz)	
ئىدە ∼	5				

CHEMIGATION

Apply this product through flood (basin) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness; or illegal pesticide residues in crop can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

A person knowledgeable of this chemigation system and responsible for soperation, to under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Rice - Flood (Basin) Chemigation

Systems using a gravity flow pesticide dispensing system must meler the pesticide into the water at the head of th field and downstream of a hydrautic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

VBC-60071 is metered or dripped into rice floodwater at application stations positioned at the point of infrodiction (levee cut) of water into each rice field or pan. Two to 3 pints of VBC-60071 are diluted in water to a final volume of 5 gallons. The diluted solution is contained in a 5 gallon container and metered or dispersed into the irrigation water using a constant flow device at the rate of 80 ml per minute. Introduction of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the hours. Floodwater delph should not exceed 10-12 inches to prevent excessive dilution of VBC-60071 which could restriction was a minimum of VBC-60071 which could the period in which the VBC-60071 suspension is being dispersed. Application of VBC-60071 into rice floodwater is not permitted using a pressurated water and pesticide injection system.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance.

30410

04-6338/R1

TEKNAR® HP BIOLOGICAL LARVICIDE AQUEOUS SUSPENSION

Active Ingredient: Bacillus thuringiensis, s fermentation solids, spores and insecticidal		
Other Ingredients		94.49
I tal		100%
Potency: 1200 International Toxic Units (I'ITU per gallon, 1.279 billion ITU per liter)	7 - - 1 -	billion
The percent active ingredient does not indicare not federally standardized.	cate product performance and po	otency measurement
	REACH OF CHILDREN CAUTION	
See side panel for additional precautionary	statements and statement of pra	ctical treatment.
Valent BioSciences Corporation		Lot No.:
870 Technology Way, Suite 100		
Libertyville, IL 60048		
		· .
EPA Reg. No. 73049-435		
EPA Est. No. 33762-IA-1		
Net Contents:		

FIRST AID			
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. 		
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water of 15-20 minutes. Call a poison control center or doctor for treatment advice. 		
HOT LINE NUMBER			

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-315-9819 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS (AND DOMESTIC ANIMALS) CAUTION

Harmful if inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash contaminated clothing before reuse.

Mixer/loaders and applicators not in enclosed cabs or aircraft must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

PHYSICAL AND CHEMICAL HAZARDS

Diluted or undiluted Teknar HP can cause corrosion if left in prolonged contact with aluminum spray system components. Rinse spray system with plenty of clean water after use. Care should be taken to prevent contact with aluminum aircraft surfaces, structural components and control systems. In case of contact, rinse thoroughly with plenty of water. Inspect aluminum aircraft component regularly for signs of corrosion.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling? Do not apply directly to treated finished drinking water reservoirs or drinking water receptacles when water is intended for human consumption.

Do not apply when weather conditions favor drift from treated areas. Do not apply to metallic painted objects, such as automobiles, as spotting may occur. If spray is deposited on metallic painted surfaces, wash immediately with soap and water to avoid spotting.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the treatment coordinator are responsible for considering all these factors when making decisions.

CHEMIGATION

Do not apply this product through any irrigation system unless the labeling on chemigation is followed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a cool, [less than 86° F (30° C)] dry place.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATION DIRECTIONS

Do not apply when wind speed favors drift beyond the area of treatment.

Mosquito Habitat

Suggested Rate Range*

(Such as the following examples): Irrigation ditches, roadside ditches, flood water, standing ponds, woodland pools, snow melt pools, pastures, catch basins, storm water retention areas, tidal water, salt marshes and rice fields.

0.25 - 2 pts/acre

In addition, standing water containing mosquito larvae, in fields growing crops such as: Alfalfa, almonds, asparagus, corn, cotton, dates, grapes, peaches and walnuts, may be treated at the recommended rates.

When applying this product to standing water containing mosquito larvae in fields growing crops, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Polluted water

1 - 2 pts/acre

(such as sewage lagoons, animal waste lagoons).

*Use higher rate range in polluted water and when late 3rd and early 4th instar larvae predominate, mosquito populations are high, water is heavily polluted, and/or algae are abundant.

Blackflies Habitat

Suggested Rate Range

Streams

Stream water** (=ppm) for

0.5 - 25 mg/liter

1 minute exposure time.

Stream water** (=ppm) for

0.05 - 2.5 mg/liter

10 minutes exposure time.

**Use higher rate range when stream contains high concentration of organic; materials, algae, or dense aquatic vegetation.

**Discharge is a principal factor determining carry of Bti. Use higher rate or increase volume by water dilution in low discharge rivers or streams under low volume (drought) conditions.

))))))

(NUISANCE FLIES)

For control of nuisance flies (*Psychoda* spp., *Chironomus* spp.) in sewage treatment facilities utilizing trickling filter systems.

APPLICATION DIRECTIONS

Nuisance Fly Habitat	Suggested Rate Range*
Trickling filter system	10-20 mg/liter a.(0.833-1.67 ml)
of wastewater treatment	per liter of wastewater feed to the
plants	filter per 30 minutes

*Use high rate for control of *Chironomus* spp. Apply undiluted with pre-calibrated pump or other device into the wastewater feeding into the filters for a period of 30 minutes. Repeat applications as needed after 2-4 weeks. Control of *Chironomus* spp. may take up to 2 weeks.

(NUISANCE AQUATIC MIDGES)

For control of *Chironomine* midges (*Chironominae*: *Chironomini*) inhabiting shallow, man-made and natural lakes or ponds.

APPLICATION DIRECTIONS

Nuisance Midge Habitat	Suggested Rate Range*
Shallow Lakes and Ponds	1 gallon (3,785.5 ml) per acre
per sewage oxidation ponds (less than acre	
6 feet deep)	

*Apply diluted with water in total volume of 5 gallons/acre by pouring or spraying over the surface to be treated with pre-calibrated device. Repeat application as needed after 2-4 weeks. Control of *Chironomine* midges may take up to 2 weeks.

Ground and Aerial Application

Teknar HP may be applied in conventional ground or aerial application equipment with quantities of water sufficient to provide uniform coverage of the target area. The amount of water will depend on weather, spray equipment, and mosquito habitat characteristics. Do not mix more Teknar HP than can be used in a 72 hour period.

For most ground spraying, apply in 5 - 100 gallons of water per acre using hand pump, airblast, mist blower, etc., spray equipment.

75049-435/notification/07-21-09/dmb

For aerial application, Teknar HP may be applied either undiluted or diluted with water. For undiluted applications, apply 0.25 to 2.0 pts./acre of Teknar HP through fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or rotary atomizers.

For diluted application, fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical or hydraulic agitation to provide moderate circulation before adding the Teknar HP. Teknar HP suspends readily in water and will stay suspended over normal application periods. Brief recirculation may be necessary if the spray mixture has sat for several hours or longer. AVOID CONTINUOUS AGITATION OF THE SPRAY MIXTURE DURING SPRAYING.

Rinse and flush spray equipment thoroughly following each use.

For blackfly aerial applications, Teknar HP can be applied undiluted via fixed wing or helicopter aircraft equipped with either conventional boom and nozzle systems or open pipes. Rate of application will be determined by the stream discharge and the required amount of Teknar HP necessary to maintain a .5 - 25 ppm concentration in the stream water. Teknar HP can also be applied diluted with similar spray equipment. Do not mix more Teknar HP than can be used in a 72 hour period.

SMALL QUANTITY DILUTION RATES

Gallons Spray Solution/Acre (Ounces Needed per Gallon of Spray)

Teknar HP Rate in Pints

Per acre	10 Gal/A	25 Gal/A	50 Gal/A
0.25 (4 oz)	0.4	0.16	0.08
0.5 (8 oz)	0.8	0.32	0.16
1.0 (16.oz)	1.6	0.64	0.32
2.0 (32 oz)	3.2	1.28	0.64

CHEMIGATION

Apply this product through flood (basin) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in crop can result from non-uniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

A person knowledgeable of this chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Rice - Flood (Basin) Chemigation

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Teknar HP is metered or dripped into rice floodwater at application stations positioned at the point of introduction (levee cut) of water into each rice field or pan. Two to 3 pints of Teknar HP are diluted in water to a final volume of 5 gallons. The diluted solution is contained in a 5 gallon container and metered or dispersed into the irrigation water using a constant flow devise at the rate of 80 ml per minute. Introduction of the solution should begin when 1/3 to 1/2 of the pan or field is covered with floodwater. Delivery of the solution should continue for a period of approximately 4 1/2 hours. Floodwater depth should not exceed 10-12 inches to prevent excessive dilution of Teknar HP which could result in reduced larval kill. Agitation is not required during the period in which the Teknar HP suspension is being dispersed. Application of Teknar HP into rice floodwater is not permitted using a pressurized water and pesticide injection system.

NOTICE TO USER

Seller makes no warranty, expressed or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

© 2009