



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

73049-503

Date of Issuance:

3/4/2015

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

VBC-30396 Plant Growth
 Regulator Water Soluble Granule

Name and Address of Registrant (include ZIP Code):

Valent BioSciences Corporation
 870 Technology Way, Suite 100
 Libertyville, IL 60048

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.
2. Submit storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) data as these data requirements are not satisfied. A one-year study is required to satisfy these data requirements. You have 18 months from the date of this registration to provide these data to the EPA.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

3/4/2015

3. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 73049-503.”
4. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 02/27/2015

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Gina Burnett of my team by phone at (703) 605-0513 or via email at burnett.gina@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

VBC-30396
Plant Growth Regulator
Water Soluble Granule

Active Ingredients

Gibberellin A ₃	40.00% w/w
S-Abscisic Acid.....	3.12% w/w
Other Ingredients.....	56.88% w/w
Total.....	100.00% w/w

Contains a total of 128 grams of Gibberellin A₃ and 9.99 grams of S-Abscisic Acid in 320 grams of product.


KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside booklet for Precautionary Statements.

EPA Registration No. xxxx
 EPA Establishment No. xxxx

Valent BioSciences Corporation
 870 Technology Way, Suite 100
 Libertyville, IL 60048
 1-847-968-4700



FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call toll-free 1-800-892-0099 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-6-Valent.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Caution: Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Waterproof gloves.
- Shoes plus socks.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribe agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours *unless wearing appropriate PPE*.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

GENERAL DIRECTIONS FOR USE

Use only as directed. Read the label thoroughly and make sure it is understood before making applications. Keep out of reach of children.

Application Instructions:

- VBC-30396 water soluble granule contains gibberellic acid and S-abscisic acid which are potent plant growth regulators. When applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, result in undesirable effects. Always consult the local Valent representative or crop specialist in your area for the spray regimen best suited to your conditions.
- Do not apply to plants under pest, nutritional, or water stress.
- When a range of rates is indicated, use the concentration and spray volume indicated locally by the local Valent representative or crop specialist.
- For optimum effectiveness, thorough spray coverage of the target area must be achieved. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Use of a non-ionic surfactant has been shown to increase wetting and uptake of the active ingredients. Discard any unused spray material at the end of each day following local, state or federal law.
- For most efficacious results, use water with a pH of 4.0 to 8.5. Use buffer for water with pH above or below this range.
- Applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption of the active ingredient by the plant, thus optimizing

effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.

- Rainfastness: Re-apply if significant rain occurs within 2 hours of application.
- Avoid drift or accidental application to other crops.
- **Compatibility:** When considering tank mixing with other products, use the following compatibility jar test before mixing a whole tank.

Start with a clear glass or plastic quart jar. Add water from the same water source that will be used for the larger tank mix. Add the pesticides in correct proportions. Mix thoroughly and let stand for a minimum 15 minutes. Heat, separation or gelling are all signs of incompatibility. Before using any mixes that pass the jar tests for compatibility, it is imperative to test the mixture on a designated area as it may result either in phytotoxicity or ineffectiveness. For further information, consult your local Valent representative.

- For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- No pre-harvest interval is required for this product.
- Entry into treated areas is allowed after the restricted entry interval (REI) of 4 hours. Before this time, entry is prohibited unless wearing appropriate PPE (coveralls, waterproof gloves, shoes plus socks).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Keep containers tightly closed when not in use.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal

The product bottle is a non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or mix tank. Fill container ¼ full with water and recap. Shake 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then recycle or dispose of in a sanitary landfill or incinerate, if allowed by state and local authorities. Stay out of smoke if incinerated.

SPRAY GUIDELINES

Apply in sprays of sufficient water volumes to ensure thorough wetting. Tank-mixing with surfactants, fertilizers, and/or other pesticides should not be done unless compatibility and phytotoxicity testing is done first using appropriate methods.

DIRECTIONS FOR CHEMIGATION

Fill the supply tank with the desired amount of water and begin agitation. Agitation should be maintained throughout the mixing and application process. Add the required amount of VBC-30396 to supply tank in order to achieve the final solution rate recommended for the specific crop to be treated. VBC-30396 should be applied at the end of irrigation water application (prior to last complete cycle in moving systems).

CHEMIGATION PRECAUTIONS:

Apply this product only through the following systems: center pivot, lateral move, side/wheel roll, traveler, solid set, big gun or hand move which have overhead sprinklers. Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Prior to application ensure that the chemigation system meets the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year. Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

PASTURES & FORAGE – FIELD USES			
CROP/VARIETY	OBJECTIVE/BENEFIT	USE RATE/ACRE	APPLICATION TIMING
Perennial Forage Grasses	To stimulate dry matter production for grazing, hay, green chop or silage when cool season conditions limit growth rates.	0.3 – 1.0 Ounces product	Spring Application: 1 to 3 applications every 3 to 4 weeks starting at green up after 1 to 2 inches of new shoot growth has emerged. Autumn Application: 1 to 3 applications every 3 to 4 weeks starting when forage growth has slowed due to cool temperatures. Best response occurs when average daily temperatures are between 40° F to 60° F and adequate moisture and nutrition are present.
Annual Forage Grasses	To stimulate dry matter production for grazing, hay, green chop or silage when cool season conditions limit growth rates.	0.3 – 1.0 Ounces product	Apply 1 to 6 applications every 3 to 4 weeks from autumn to early spring during periods of suboptimal growth due to cool temperatures. If applying to over-seeded pasture or newly established pasture, apply only after seedlings are well established. Best response occurs when average daily temperatures are between 40° F to 60° F and adequate moisture and nutrition are present.
Cereal Grains (such as barley, oats, rye, sorghum, wheat, triticale)	To stimulate dry matter production for grazing, hay, green chop or silage when cool season conditions limit growth rates.	0.3 – 1.0 Ounces product	Spring Application: 1 to 3 applications every 3 to 4 weeks starting at green up after 1 to 2 inches of new shoot growth has emerged. Autumn Application: 1 to 3 applications every 3 to 4 weeks starting when forage growth has slowed due to cool temperatures. Application to cereal grains during stem elongation (jointing onwards) can

			<p>result in lodging. Apply during early tillering growth stages prior to stem elongation to avoid lodging.</p> <p>Best response occurs when average daily temperatures are between 40° F to 60° F and adequate moisture and nutrition are present.</p>
<p>Winter Brassicas (such as turnip, kale, rape)</p>	<p>To stimulate dry matter production for grazing, hay, green chop or silage when cool season conditions limit growth rates.</p>	<p>0.3 – 1.0 Ounces product</p>	<p>Spring Application: 1 to 3 applications every 3 to 4 weeks starting at green up after 1 to 2 inches of new shoot growth has emerged.</p> <p>Autumn Application: 1 to 3 applications every 3 to 4 weeks starting when forage growth has slowed due to cool temperatures.</p> <p>Best response occurs when average daily temperatures are between 40° F to 60° F and adequate moisture and nutrition are present.</p>
<p>NOTE:</p> <ul style="list-style-type: none"> • Foliage occasionally and temporarily appears lighter green in color due to accelerated growth rates following application. For best results, ensure soil fertility is adequate to sustain additional pasture growth. • Plants will not respond to treatment without adequate moisture or if under pest and/or nutritional stress. • Once plants are at their maximum growth rate, application of VBC-30396 will not stimulate additional growth. • Plants will not respond when the ground is frozen. • Plants treated at maximum physiological size will not respond with additional growth. 			

FIELD USES			
CROP/VARIETY	OBJECTIVE/BENEFIT	USE RATE/ACRE	APPLICATION TIMING
Corn: Silage, Field	To increase yield and help overcome the effects of heat or drought.	0.3 – 0.6 Ounces product	Apply at V2 - V6
Corn: Popcorn, Sweet corn, Seed corn	To increase yield and help overcome the effects of heat or drought.	0.3 – 0.6 Ounces product	Apply at V2 – V6
<p>Note:</p> <ul style="list-style-type: none"> • Application should be made to corn plants when two leaves (V2) to six leaves (V6) with collars are visible. • Foliage occasionally and temporarily appears lighter green in color due to accelerated growth rates following application. For best results, ensure soil fertility is adequate to sustain additional growth. • Plants will not respond to treatment without adequate moisture or if under pest and/or nutritional stress. • A non-ionic surfactant can improve product efficacy. • VBC-30396 is compatible as a tank-mix partner with Roundup® herbicide on glyphosate resistant corn. Use of VBC-30396 with other tank-mix partners is done solely at the user's risk. • Always consider use recommendations of tank-mix partners when using VBC-30396 mixed with other products. • Do not tank-mix VBC-30396 with 2,4-D or any herbicide containing 2,4-D when applying to corn. • VBC-30396 enhances the effect of some HPPD (Group #27) herbicides and can injure corn when applied post emergent to hybrids with known sensitivity to HPPD herbicides. Users should understand and accept this risk before applying VBC-30396 on corn with HPPD herbicides. 			

USES ON OTHER CROPS			
CROP/VARIETY	OBJECTIVE/BENEFIT	USE RATE/ACRE	APPLICATION TIMING
Cotton	Promote early season growth and increase seedling vigor	0.1 to 0.6 Ounces products	Apply 1 – 2 applications as a foliar broadcast spray during the 3 to 7 leaf/node stage. If applying as a banded spray, reduce rates accordingly. Complete coverage of leaf tissue is essential. Use higher rates when temperatures will likely average 75° F or less during the 14 days following application(s).
Sugarcane	To maintain yields in older plantings, increase bio-mass and stimulate growth before harvest of cane in older production fields (>3 years)	0.1 to 0.6 Ounces products	Apply at 1 st to 5 th internode stage to new plantings or ratoon crop in at least 20 gal/A. Addition of non-ionic surfactant may increase activity.

WARRANTY AND DISCLAIMER STATEMENT

To the fullest extent permitted by law, seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

Roundup® is a registered trademark of Monsanto Company.

VBC-30396 is a registered trademark of Valent BioSciences Corporation.

Products That Work, From People Who Care is a trademark of Valent U.S.A. Corporation.

© 2014 Valent BioSciences Corporation

Registered by:

Valent BioSciences Corporation

870 Technology Way, Suite 100

Libertyville, IL 60048

04-XXXX/RX

Distributed by Valent U.S.A. Corporation.