



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Ms. Jane Rothwell  
Registration Specialist  
Farmsaver.com, LLC  
4515 Fals of Neuse Road., Suite 300  
Raleigh, NC 27609

MAY 30 2008

RE: Notification to update Warranty Statement and alternate marketing language  
EPA Registration Number: 73220-11  
Date of Submission: April 23, 2008

Dear Ms. Rothwell:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated April 23, 2008, for the Quali-Pro Oxadiazon 2G. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

### Application for Pesticide - Section I

1. Company/Product Number 73220-11	2. EPA Product Manager Sherada Hobgood	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Quali-Pro Oxadiazon 2G	PM#	
5. Name and Address of Applicant (Include ZIP Code) Farmsaver.com, LLC 4515 Falls of Neuse Rd., Suite 300 Raleigh, NC 27609 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____  Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.


**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of updated Warranty Statement and alternate marketing language. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

### Section - III

<b>1. Material This Product Will Be Packaged In:</b>			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt.    No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt    No. per container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	
5. Location of Label Directions <input type="checkbox"/> _____		6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____	

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jane Rothwell	Title Registration Specialist	Telephone No. (If include Area Code) 801-881-4400
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Specialist	
4. Typed Name Jane Rothwell	5. Date 4-23-08	

# Quali-Pro<sup>®</sup> Oxadiazon 2G

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1, 3,4-oxadiazoline-5-one]: .....2.0%

**OTHER INGREDIENTS:** .....98.0%

**TOTAL:** .....100.0%

[Optional Text – Guaranteed Fertilizer Analysis]

EPA Reg. No. 73220-11

EPA Est. No.

Manufactured for:  
FarmSaver.com, LLC  
4515 Falls of Neuse Road, Suite 300  
Raleigh, NC 27609

**Net Weight: 50 lbs.**

## KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact PROSAR 1-800-308-5391 for emergency medical treatment information.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. DO NOT get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material

**GENERAL INFORMATION**

Quali-Pro Oxadiazon 2G is a selective, pre-emergent herbicide for control of annual grasses and many broadleaf weeds in commercial turf and on a wide variety of woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

For best results, water-in the product as soon as practical after application.

10-10-10 fertilizer with secondary micronutrients is the carrier for the herbicide. Nutrients will be supplied according to the rates applied, but they are inert to the herbicide and its activity.

**State Specific Restrictions:**

- The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

**Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).**

**Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.**

### COMMERCIAL TURF

For use in established perennial Bluegrass, Bermudagrass, perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses, and lawns. Quali-Pro Oxadiazon 2G controls Crabgrass (*digitaria spp.*), Annual Bluegrass (*Poa annua*), Goosegrass (*Eleusine indica*); and it also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis spp.*), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus offinis*), and Pigweed (*Amaranthus spp.*).

Apply to dry foliage only with a properly calibrated granular applicator at recommended rate in early spring before weed seed germination. If rain is not expected shortly after application, irrigation to remove herbicide granules from foliage will increase effectiveness.

For Crabgrass, Goosegrass and Crowfoot control, apply 100-200 pounds Quali-Pro Oxadiazon 2G per acre (2 to 4 pounds active ingredient) or 2 ¼ to 4 ½ pounds per 1000 sq. ft. Late applications are not recommended and may not be fully effective.

For Annual Bluegrass control apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds/1000 sq. ft. in early spring or late summer prior to seed germination.

For Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass control, apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds per 1000 sq. ft. in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 150 pounds/acre (3 pounds active ingredient) of Quali-Pro Oxadiazon-2G.

Discoloration of Bermudagrass and St. Augustinegrass may result from use of this herbicide, but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

Red Fescue and Bentgrass are susceptible to Quali-Pro Oxadiazon 2G and may be severely damaged. Avoid use of this product on turf containing Red Fescue or Bentgrass except where they are undesirable grasses. Do not apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. Quali-Pro Oxadiazon 2G should not be used on putting greens or tees because of varying cultural practices used.

### WOODY ORNAMENTALS SHRUBS, VINES, AND TREES

#### APPLICATION:

Apply Quali-Pro Oxadiazon 2G anytime during the year, prior to weed seed germination, for weed control. Quali-Pro Oxadiazon 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Quali-Pro Oxadiazon 2G will improve weed control activity. Quali-Pro Oxadiazon 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 2G. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

#### AMOUNT TO USE:

**Ground Application-** Apply Quali-Pro Oxadiazon 2G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 ¼ to 4 ½ pounds per 1000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Quali-Pro Oxadiazon 2G at 100 to 200 pounds (2.0 to 4.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broad-leaf weeds such as:

Annual Broadleaf Weeds	
Bittercress	<i>Cardamine spp.</i>
Carpetweed	<i>Mollugo verticillata</i>
Common Groundsel	<i>Senecio vulgaris</i>
Common Purslane	<i>Portulaca oleracea</i>
*Eveningprimrose	<i>Oenothera spp.</i>
*Galinsoga	<i>Galinsoga spp.</i>
Golden Ragwort	<i>Senecio aureus</i>
Lambsquarters	<i>Chenopodium album</i>

Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>
Petty Spurge	<i>Euphorbia peplus</i>
Redroot Pigweed	<i>Amaranthus Retroflexus</i>
Speedwell	<i>Veronica spp.</i>
Spotted Catsear	<i>Hypochaeris radicata</i>
*Swinecress	<i>Coronopus didymus</i>
Yellow Woodsorrel	<i>Oxalis stricta</i>
<b>Annual Grasses</b>	
Annual Bluegrass	<i>Poa annua</i>
*Barnyardgrass	<i>Echinochloa crus-galli</i>
Crabgrass	<i>Digitaria spp.</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Green Foxtail	<i>Setaria viridis</i>

\*Use 150 to 200 pounds Quali-Pro Oxadiazon 2G (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

Note: For the control of liverwort (*Marchantia spp.*) in California, Goosegrass (*Eleusine Indica*) and Prostrate Spurge (*Euphorbia supina*) use 200 pounds Quali-Pro Oxadiazon 2G (4.0 pounds of active ingredient) per acre.

**CROP TOLERANCE:**

Quali-Pro Oxadiazon 2G is recommended for use on a wide variety of container and field grown woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G may be used at the recommended rates on the following ornamental species:

<b>CONTAINER GROWN</b>	
<b>ESTABLISHED</b>	
<b>Species</b>	<b>Varieties</b>
Agapanthus	Africanus
Andromeda	Japanese
Arborvitae	American, Eastern, Oriental, Aureanana, Globe, Pyramidal, Woodward
Arctostaphylos	Fred Oelher
Azalea	Hino Red, Red, Boudoir, Snowball, Karen, Kaempp, Stuartonian, Mollishybird, Sweetheart Supreme, Salmon Beauty, Caroline Gable, Louise, Stewartstonian, Rosebud
Barberry	Redleaf, Marshall Red, Darwinii, Irwinii, Purpleleaf
Boxwood	Common, Japanese
Callistemon	Bottle-brush
Camellia	Glen #40, Sasanqua, Colonel Fiery
Ceanothus Cedar	Atlas, Portoford (Allumii)
Cotoneaster	Paineyi, Bearberry, Coral Beauty, Greyleaf, Cranberry, Silverleaf, Peking, Bright Bead, Dammeri Skolgholm, Microphylla, Lowfast, Glaucophylla.
Cypress	Monterey
Dogwood	Red osier
Douglas Fir	Pseudotsugamenziesii
Eucalyptus	Sideroxylon, Nicholli
Euonymus	Greenlane, Microphylla Variegata, Emeraldgaety, Dwarf, Gold spot, Golden Green, Silver King, Wintercreeper Vegetus, Ermerald Cushion, Spreading
Forsythia	Lynwood gold
Gardenia	Amy Yashioka, August Beauty, Mystery
Ginko	Biloba
Heath Heather	Mrs. Maxwell
Hedera	English Ivy
Honeysuckle	Purpleleaf, Purpurea
Holly	Dwarf Okinowan, Convexa, Rotundifolia, Hetzi, Helleri, Burfordi, English

	Hellus, Dwarf Yaupon, Nellie R. Stevens, Boxleafod, Green Island, Japanese
Jasmine	Yellow Star, Star
Juniper	Barharbor, Wiltoni, Torulosa, Pfitzeriana Compacti, Pfitzeriana Aurea, Torulosa Variegata, Pfitzeriana Glauca, Gulf Stream, Tam, Armstrongii, Shore Andora, San Jose, Blue Haven, Stricta, Jade River, Broadmoor, Meyeri, Rocket, Plumosa, Royal Beauty, Praecox, Greek, Apiculata Blackburn, Plumosa Compacta, Procumbens, Blue Pfizer
Justica	Brandegeana
Leucothoe	Catesbaei, Axillaris
Myrtle	Compacta Noellii
Magnolia	Dr. Merrilli, Smauel Somers, Grandiflora
Mahonia	Oregon Grape-Holly
Osmanthus	Fortunes
Periwinkle	Vinca Minor
Photinia	Frazerii
Pine	Slash, Japanese Black, Montana, Scotch, Digger, Shore, Thumbergii
Pittosporum	Variegated, Japanese
Podocarpus	Macrophylla
Privet	Japanese, Waxleaf, Glossy, Vicary, Golden Vicary, Curly Leaf
Pyracantha	Victory, Scarlet Firethorn, Wyattii, Graberri
Quercus Raphiolepis	India Hawthorn
Rhododendron	Cunningham, Blue Diamond, Snow, Christmas, Chionoides, 'PJM', Dr. Rutgers, Coral Bells, President Clay
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong
Star Jasmine	Yellow Star Jasmine
Syzygium	Paniculatum
Viburnum	Sandankwa, Laurestinus, David
Xylosma	Congestum

**NEWLY TRANSPLANTED**

<b>Species</b>	<b>Varieties</b>
Andromeda	Japanese
Arbovitae	Glove, Woodward
Azalea	Hindo Red, Louise, Rosebud, Stewartstonian
Barberry	Marshall Red, Darwinii, Irwinii, Purpleleaf
Cedar	Portoford (allumii)
Cotoneaster	Cranberry, Peking, Coral Beauty, Bright Bead, Painei, Dammeri Skolgholm, Microphylla, Glaucohylla
Cypress	Monterey
Dogwood	Red Osier
Euonymus	Greenlane, Emerald Cushion, Spreading
Forsythia	Lynwood Gold
Holly	Rotundifolia, Helleri, English, Japanese
Juniper	Plumosa, Plumosa Compacta, Old Gold, Pfitzeriana Aurea, Torulosa, Armstrongii, Tam, Wiltoni, Procumbens
Leucothoe	Axillaris
Magnolia	Dr. Merrilli
Mahonia	Oregon Grape-Holly
Pine	Slash
Privet	Japanese, Waxleaf, Glossy
Pyracantha	Graberii
Rhododendron	Snow, Blue Diamond, Chionoides, 'PJM', Dr. Rutgers
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong

FIELD GROWN	
ESTABLISHED	
Species	Varieties
Arborvitae	Pyramidal
Ash	Green
Azalea	Hinodi geri
Barberry	Red, Mentor
Birch	Clump, Paper
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, Tom Knudsen, Crimson Robe, R.D. Wheeler
Cotoneaster	Lowfast, Skolgholm
Crabapple	Flowing, Toringo
Dogwood	Redstemmed, Red, Flowering
Euonymus	Burning Bush, Dwarf Burning Bush, Jewell
Forsythia	Showy Boarder, Lynngold, Lynwood Gold, Goldenbell
Holly	Boxleafed, Green Island, Japanese
Honeysuckle	Zabeli
Juniper	Plumosa, Mint Julip, Arcardia, Variegata; Plumosa Compacta, Andora, Wiltoni, Hetzi
Lilac	Persian, Dwarf Korean
Maple	Amur
Oak	Red
Perwinkle	Vinca Minor
Pine	Austrian, Scotch, Norway, Stone, Japanese, Black, Loblolly, Slash, Eastern White
Pittosporum	Undulatum
Privet	California, Japanese
Rhododendron	Rosebud
Rose	Texas Centennial, Sutters Gold, Show Girl
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Taunton Spreading Yew, Hicks
Tsuga	Canadian Hemlock
Viburnum	European Cranberry Bush
Xylosma	
Ground Covers	
Carpobrotus	Edule
Osteo	
Spernum Sedum	Album
Algerian Ivy	
Vinca Major	
English Ivy	
Pachysandra	Terminalis
Ajuga	Reptans
Euonymus	Fortunei Radicans
NEWLY TRANSPLANTED	
Species	Varieties
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, R.D. Wheeler, Tom Knudsen, Crimson Robe
Dogwood	Redstemmed, Red, Flowering
Forsythia	Showy Border, Lynwood Gold, Goldenbell
Hedera	English Ivy
Holly	Green Island, Japanese
Juniper	Plumosa, Andora, Wiltoni, Hetzi



Oak	Red
Periwinkle	Vinca Minor
Pine	White Austrian, Eastern White
Privet	California, Japanese
Rhododendron	Rosebud
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Hicks
Tsuga	Canadian Hemlock

**ADDITIONAL APPLICATIONS**

Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).

Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.

For continued weed control, additional application of Quali-Pro Oxadiazon 2G may be made 60 to 120 days after the previous application in the following situations:

2 REPEAT TREATMENTS					
Container Grown Established		Container Grown Newly Transplanted		Field Grown Established and Newly Transplanted	
Species	Varieties	Species	Varieties	Species	Varieties
Abelia	Grandiflora	Common Boxwood		Holly	Nellie R. Stevens, Dwarf Buford, Buford
Common boxwood		English Ivy			
English Ivy		Juniper	Tam		
Holly	Helleri	Pine	Slash		
Juniper	Tam	Rhododendron	Obtusa, Kurume Formosa		
Pine	Slash				
Rhododendron	Obtusa, Kurume Formosa, Kurume Snow				
3 REPEAT TREATMENTS					
Container Grown Established					
Species		Varieties			
Abelia		Grandiflora			
Holly		Helleri, Rotundifolia			
Rhododendron		Kurume Formosa, Kurume Snow			
Greenhouse Grown Established					
Species		Varieties			
Rose		Sonia, Forever Yours, Coed, Golden Wave			
Forest Nurseries Seedlings- At least 4 weeks old					
Species		Varieties			
Pine		Loblolly, Slash, Eastern White			

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in dry area. Do not store near other pesticides, or seeds.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In case of spill, fire, or leak, contact CHEMTREC at 1-800-424-9300.

**CONDITIONS OF SALE AND WARRANTY**

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of FarmSaver.com, LLC or the seller. All such risks shall be assumed by the buyer.

FarmSaver.com, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE when it is used in accordance with such directions, subject to the inherent risks mentioned above.

FARMSAVER.COM, LLC NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED ON CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. FarmSaver.com, LLC and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

**LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Farmsaver.com, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Farmsaver.com, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Farmsaver.com, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Farmsaver.com, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Farmsaver.com, LLC's election, the replacement of product.

Quali-Pro® is a registered trademark of FarmSaver.com, LLC.

Quali-Pro Oxadiazon 2G (73220-11)(EPA app 1-26-07)(notif to EPA 4-23-08)

# Quali-Pro® Oxadiazon 2G

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1, 3,4-oxadiazoline-5-one]:.....2.0%

**OTHER INGREDIENTS:**.....98.0%

**TOTAL:**.....100.0%

[Optional Text – Guaranteed Fertilizer Analysis]

EPA Reg. No. 73220-11

EPA Est. No.

Manufactured for:  
FarmSaver.com, LLC  
4515 Falls of Neuse Road, Suite 300  
Raleigh, NC 27609

**Net Weight: 50 lbs.**

## KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact PROSAR 1-800-308-5391 for emergency medical treatment information.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. DO NOT get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material

**GENERAL INFORMATION**

Quali-Pro Oxadiazon 2G is a selective, pre-emergent herbicide for control of annual grasses and many broadleaf weeds in commercial turf and on a wide variety of woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

For best results, water-in the product as soon as practical after application.

10-10-10 fertilizer with secondary micronutrients is the carrier for the herbicide. Nutrients will be supplied according to the rates applied, but they are inert to the herbicide and its activity.

**State Specific Restrictions:**

- The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

**Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).**

**Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.**

### COMMERCIAL TURF

For use in established perennial Bluegrass, Bermudagrass, perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses, and lawns. Quali-Pro Oxadiazon 2G controls Crabgrass (*digitaria spp.*), Annual Bluegrass (*Poa annua*), Goosegrass (*Eleusine indica*); and it also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis spp.*), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus offinis*), and Pigweed (*Amaranthus spp.*).

Apply to dry foliage only with a properly calibrated granular applicator at recommended rate in early spring before weed seed germination. If rain is not expected shortly after application, irrigation to remove herbicide granules from foliage will increase effectiveness.

For Crabgrass, Goosegrass and Crowfoot control, apply 100-200 pounds Quali-Pro Oxadiazon 2G per acre (2 to 4 pounds active ingredient) or 2 ¼ to 4 ½ pounds per 1000 sq. ft. Late applications are not recommended and may not be fully effective.

For Annual Bluegrass control apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds/1000 sq. ft. in early spring or late summer prior to seed germination.

For Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass control, apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds per 1000 sq. ft. in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 150 pounds/acre (3 pounds active ingredient) of Quali-Pro Oxadiazon 2G.

Discoloration of Bermudagrass and St. Augustinegrass may result from use of this herbicide, but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

Red Fescue and Bentgrass are susceptible to Quali-Pro Oxadiazon 2G and may be severely damaged. Avoid use of this product on turf containing Red Fescue or Bentgrass except where they are undesirable grasses. Do not apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. Quali-Pro Oxadiazon 2G should not be used on putting greens or tees because of varying cultural practices used.

### WOODY ORNAMENTALS SHRUBS, VINES, AND TREES

#### APPLICATION:

Apply Quali-Pro Oxadiazon 2G anytime during the year, prior to weed seed germination, for weed control. Quali-Pro Oxadiazon 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Quali-Pro Oxadiazon 2G will improve weed control activity. Quali-Pro Oxadiazon 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 2G. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

#### AMOUNT TO USE:

**Ground Application-** Apply Quali-Pro Oxadiazon 2G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 ¼ to 4 ½ pounds per 1000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Quali-Pro Oxadiazon 2G at 100 to 200 pounds (2.0 to 4.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broad-leaf weeds such as:

Annual Broadleaf Weeds	
Bittercress	<i>Cardamine spp.</i>
Carpetweed	<i>Mollugo verticillata</i>
Common Groundsel	<i>Senecio vulgaris</i>
Common Purslane	<i>Portulaca oleracea</i>
*Eveningprimrose	<i>Oenothera spp.</i>
*Galinsoga	<i>Galinsoga spp.</i>
Golden Ragwort	<i>Senecio aureus</i>
Lambsquarters	<i>Chenopodium album</i>

14/27

Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>
Petty Spurge	<i>Euphorbia peplus</i>
Redroot Pigweed	<i>Amaranthus Retroflexus</i>
Speedwell	<i>Veronica spp.</i>
Spotted Catsear	<i>Hypochaeris radicata</i>
*Swinecress	<i>Coronopus didymus</i>
Yellow Woodsorrel	<i>Oxalis stricta</i>
<b>Annual Grasses</b>	
Annual Bluegrass	<i>Poa annua</i>
*Barnyardgrass	<i>Echinochloa crus-galli</i>
Crabgrass	<i>Digitaria spp.</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Green Foxtail	<i>Setaria viridis</i>

\*Use 150 to 200 pounds Quali-Pro Oxadiazon 2G (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

Note: For the control of liverwort (*Marchantia spp.*) in California, Goosegrass (*Eleusine Indica*) and Prostrate Spurge (*Euphorbia supina*) use 200 pounds Quali-Pro Oxadiazon 2G (4.0 pounds of active ingredient) per acre.

**CROP TOLERANCE:**

Quali-Pro Oxadiazon 2G is recommended for use on a wide variety of container and field grown woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G may be used at the recommended rates on the following ornamental species:

<b>CONTAINER GROWN</b>	
<b>ESTABLISHED</b>	
<b>Species</b>	<b>Varieties</b>
Agapanthus	Africanus
Andromeda	Japanese
Arborvitae	American, Eastern, Oriental, Aureanana, Globe, Pyramidal, Woodward
Arctostaphylos	Fred Oelher
Azalea	Hino Red, Red, Boudoir, Snowball, Karen, Kaempp, Stuartonian, Mollishybird, Sweetheart Supreme, Salmon Beauty, Caroline Gable, Louise, Stewartstonian, Rosebud
Barberry	Redleaf, Marshall Red, Darwinii, Irwinii, Purpleleaf
Boxwood	Common, Japanese
Callistemon	Bottle-brush
Camellia	Glen #40, Sasanqua, Colonel Fiery
Ceanothus Cedar	Atlas, Portoford (Allumii)
Cotoneaster	Paineyi, Bearberry, Coral Beauty, Greyleaf, Cranberry, Silverleaf, Peking, Bright Bead, Dammeri Skolgholm, Microphylla, Lowfast, Glaucohylla.
Cypress	Monterey
Dogwood	Red osier
Douglas Fir	Pseudotsugamenziesii
Eucalyptus	Sideroxylon, Nicholli
Euonymus	Greenlane, Microphylla Variegata, Emeraldgaety, Dwarf, Gold spot, Golden Green, Silver King, Wintercreeper Vegetus, Emerald Cushion, Spreading
Forsythia	Lynwood gold
Gardenia	Amy Yashioka, August Beauty, Mystery
Ginko	Biloba
Heath Heather	Mrs. Maxwell
Hedera	English Ivy
Honeysuckle	Purpleleaf, Purpurea
Holly	Dwarf Okinowan, Convexa, Rotundifolia, Hetzi, Helleri, Burfordi, English,

	Hellus, Dwarf Yaupon, Nellie R. Stevens, Boxleafod, Green Island, Japanese
Jasmine	Yellow Star, Star
Juniper	Barharbor, Wiltoni, Torulosa, Pfitzeriana Compacti, Pfitzeriana Aurea, Torulosa Variegata, Pfitzeriana Glauca, Gulf Stream, Tam, Armstrongii, Shore Andora, San Jose, Blue Haven, Stricta, Jade River, Broadmoor, Meyeri, Rocket, Plumosa, Royal Beauty, Praecox, Greek, Apiculata Blackburn, Plumosa Compacta, Procumbens, Blue Pfizer
Justica	Brandegeana
Leucothoe	Catesbaei, Axillaris
Myrtle	Compacta Noellii
Magnolia	Dr. Merrilli, Smauel Somers, Grandiflora
Mahonia	Oregon Grape-Holly
Osmanthus	Fortunes
Periwinkle	Vinca Minor
Photinia	Frazerii
Pine	Slash, Japanese Black, Montana, Scotch, Digger, Shore, Thumbergii
Pittosporum	Variegated, Japanese
Podocarpus	Macrophylla
Privet	Japanese, Waxleaf, Glossy, Vicary, Golden Vicary, Curly Leaf
Pyracantha	Victory, Scarlet Firethorn, Wyattii, Graberri
Quercus Raphiolepis	India Hawthorn
Rhododendron	Cunningham, Blue Diamond, Snow, Christmas, Chionoides, 'PJM', Dr. Rutgers, Coral Bells, President Clay
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong
Star Jasmine	Yellow Star Jasmine
Syzygium	Paniculatum
Viburnum	Sandankwa, Laurestinus, David
Xylosma	Congestum

#### NEWLY TRANSPLANTED

Species	Varieties
Andromeda	Japanese
Arborvitae	Glove, Woodward
Azalea	Hindo Red, Louise, Rosebud, Stewartstonian
Barberry	Marshall Red, Darwinii, Irwinii, Purpleleaf
Cedar	Portoford (allumii)
Cotoneaster	Cranberry, Peking, Coral Beauty, Bright Bead, Painei, Dammeri Skolgholm, Microphylla, Glaucohylla
Cypress	Monterey
Dogwood	Red Osier
Euonymus	Greenlane, Emerald Cushion, Spreading
Forsythia	Lynwood Gold
Holly	Rotundifolia, Helleri, English, Japanese
Juniper	Plumosa, Plumosa Compacta, Old Gold, Pfitzeriana Aurea, Torulosa, Armstrongii, Tam, Wiltoni, Procumbens
Leucothoe	Axillaris
Magnolia	Dr. Merrilli
Mahonia	Oregon Grape-Holly
Pine	Slash
Privet	Japanese, Waxleaf, Glossy
Pyracantha	Graberri
Rhododendron	Snow, Blue Diamond, Chionoides, 'PJM', Dr. Rutgers
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong

FIELD GROWN	
<b>ESTABLISHED</b>	
<b>Species</b>	<b>Varieties</b>
Arborvitae	Pyramidal
Ash	Green
Azalea	Hinodi geri
Barberry	Red, Mentor
Birch	Clump, Paper
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, Tom Knudsen, Crimson Robe, R.D. Wheeler
Cotoneaster	Lowfast, Skolgholm
Crabapple	Flowing, Torringo
Dogwood	Redstemmed, Red, Flowering
Euonymus	Burning Bush, Dwarf Burning Bush, Jewell
Forsythia	Showy Boarder, Lynngold, Lynwood Gold, Goldenbell
Holly	Boxleafed, Green Island, Japanese
Honeysuckle	Zabeli
Juniper	Plumosa, Mint Julip, Arcardia, Variegata; Plumosa Compacta, Andora, Wiltoni, Hetzi
Lilac	Persian, Dwarf Korean
Maple	Amur
Oak	Red
Perwinkle	Vinca Minor
Pine	Austrian, Scotch, Norway, Stone, Japanese, Black, Loblolly, Slash, Eastern White
Pittosporum	Undulatum
Privet	California, Japanese
Rhododendron	Rosebud
Rose	Texas Centennial, Sutters Gold, Show Girl
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Taunton Spreading Yew, Hicks
Tsuga	Canadian Hemlock
Viburnum	European Cranberry Bush
Xylosma	
<b>Ground Covers</b>	
Carpobrotus	Edule
Osteo	
Spermum Sedum	Album
Algerian Ivy	
Vinca Major	
English Ivy	
Pachysandra	Terminalis
Ajuga	Reptans
Euonymus	Fortunei Radicans
<b>NEWLY TRANSPLANTED</b>	
<b>Species</b>	<b>Varieties</b>
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, R.D. Wheeler, Tom Knudsen, Crimson Robe
Dogwood	Redstemmed, Red, Flowering
Forsythia	Showy Border, Lynwood Gold, Goldenbell
Hedera	English Ivy
Holly	Green Island, Japanese
Juniper	Plumosa, Andora, Wiltoni, Hetzi



Oak	Red
Periwinkle	Vinca Minor
Pine	White Austrian, Eastern White
Privet	California, Japanese
Rhododendron	Rosebud
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Hicks
Tsuga	Canadian Hemlock

**ADDITIONAL APPLICATIONS**

Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).

Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.

For continued weed control, additional application of Quali-Pro Oxadiazon 2G may be made 60 to 120 days after the previous application in the following situations:

**2 REPEAT TREATMENTS**

Container Grown Established		Container Grown Newly Transplanted		Field Grown Established and Newly Transplanted	
Species	Varieties	Species	Varieties	Species	Varieties
Abelia	Grandiflora	Common Boxwood		Holly	Nellie R. Stevens, Dwarf Buford, Buford
Common boxwood		English Ivy			
English Ivy		Juniper	Tam		
Holly	Helleri	Pine	Slash		
Juniper	Tam	Rhododendron	Obtusa, Kurume Formosa		
Pine	Slash				
Rhododendron	Obtusa, Kurume Formosa, Kurume Snow				

**3 REPEAT TREATMENTS**

Container Grown Established	
Species	Varieties
Abelia	Grandiflora
Holly	Helleri, Rotundifolia
Rhododendron	Kurume Formosa, Kurume Snow
Greenhouse Grown Established	
Species	Varieties
Rose	Sonia, Forever Yours, Coed, Golden Wave
Forest Nurseries Seedlings- At least 4 weeks old	
Species	Varieties
Pine	Loblolly, Slash, Eastern White

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in dry area. Do not store near other pesticides, or seeds.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In case of spill, fire, or leak, contact CHEMTREC at 1-800-424-9300.

18/  
27

**LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Farmsaver.com, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Farmsaver.com, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Farmsaver.com, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Farmsaver.com, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Farmsaver.com, LLC's election, the replacement of product.

Quali-Pro® is a registered trademark of FarmSaver.com, LLC.

Quali-Pro Oxadiazon 2G (73220-11)(EPA app 1-26-07)(notif to EPA 4-23-08)

Quali-Pro Oxadiazon 2G (73220-11)(EPA app 1-26-07)(notif to EPA 4-23-08)

# Quali-Pro<sup>®</sup> Oxadiazon 2G

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1, 3,4-oxadiazoline-5-one]:.....2.0%

**OTHER INGREDIENTS:**.....98.0%

**TOTAL:**.....100.0%

[Optional Text – Guaranteed Fertilizer Analysis]

EPA Reg. No. 73220-11

EPA Est. No.

Manufactured for:  
FarmSaver.com, LLC  
4515 Falls of Neuse Road, Suite 300  
Raleigh, NC 27609

**Net Weight: 50 lbs.**

## KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact PROSAR 1-800-308-5391 for emergency medical treatment information.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes substantial but temporary eye injury. DO NOT get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

#### Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material

### GENERAL INFORMATION

Quali-Pro Oxadiazon 2G is a selective, pre-emergent herbicide for control of annual grasses and many broadleaf weeds in commercial turf and on a wide variety of woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

For best results, water-in the product as soon as practical after application.

10-10-10 fertilizer with secondary micronutrients is the carrier for the herbicide. Nutrients will be supplied according to the rates applied, but they are inert to the herbicide and its activity.

#### State Specific Restrictions:

- The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

**Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).**

**Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.**

### COMMERCIAL TURF

For use in established perennial Bluegrass, Bermudagrass, perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses, and lawns. Quali-Pro Oxadiazon 2G controls Crabgrass (*digitaria spp.*), Annual Bluegrass (*Poa annua*), Goosegrass (*Eleusine indica*); and it also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis spp.*), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus offinis*), and Pigweed (*Amaranthus spp.*).

Apply to dry foliage only with a properly calibrated granular applicator at recommended rate in early spring before weed seed germination. If rain is not expected shortly after application, irrigation to remove herbicide granules from foliage will increase effectiveness.

For Crabgrass, Goosegrass and Crowfoot control, apply 100-200 pounds Quali-Pro Oxadiazon 2G per acre (2 to 4 pounds active ingredient) or 2 ¼ to 4 ½ pounds per 1000 sq. ft. Late applications are not recommended and may not be fully effective.

For Annual Bluegrass control apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds/1000 sq. ft. in early spring or late summer prior to seed germination.

For Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass control, apply 200 pounds/acre Quali-Pro Oxadiazon 2G (4 pounds active ingredient) or 4 ½ pounds per 1000 sq. ft. in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 150 pounds/acre (3 pounds active ingredient) of Quali-Pro Oxadiazon 2G.

Discoloration of Bermudagrass and St. Augustinegrass may result from use of this herbicide, but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

Red Fescue and Bentgrass are susceptible to Quali-Pro Oxadiazon 2G and may be severely damaged. Avoid use of this product on turf containing Red Fescue or Bentgrass except where they are undesirable grasses. Do not apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. Quali-Pro Oxadiazon 2G should not be used on putting greens or tees because of varying cultural practices used.

### WOODY ORNAMENTALS SHRUBS, VINES, AND TREES

#### APPLICATION:

Apply Quali-Pro Oxadiazon 2G anytime during the year, prior to weed seed germination, for weed control. Quali-Pro Oxadiazon 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Quali-Pro Oxadiazon 2G will improve weed control activity. Quali-Pro Oxadiazon 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 2G. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

#### AMOUNT TO USE:

**Ground Application-** Apply Quali-Pro Oxadiazon 2G broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 ¼ to 4 ½ pounds per 1000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Quali-Pro Oxadiazon 2G at 100 to 200 pounds (2.0 to 4.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broad-leaf weeds such as:

Annual Broadleaf Weeds	
Bittercress	<i>Cardamine spp.</i>
Carpetweed	<i>Mollugo verticillata</i>
Common Groundsel	<i>Senecio vulgaris</i>
Common Purslane	<i>Portulaca oleracea</i>
*Eveningprimrose	<i>Oenothera spp.</i>
*Galinsoga	<i>Galinsoga spp.</i>
Golden Ragwort	<i>Senecio aureus</i>
Lambsquarters	<i>Chenopodium album</i>

Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>
Petty Spurge	<i>Euphorbia peplus</i>
Redroot Pigweed	<i>Amaranthus Retroflexus</i>
Speedwell	<i>Veronica spp.</i>
Spotted Catsear	<i>Hypochaeris radicata</i>
*Swinecress	<i>Coronopus didymus</i>
Yellow Woodsorrel	<i>Oxalis stricta</i>
<b>Annual Grasses</b>	
Annual Bluegrass	<i>Poa annua</i>
*Barnyardgrass	<i>Echinochloa crus-galli</i>
Crabgrass	<i>Digitaria spp.</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Green Foxtail	<i>Setaria viridis</i>

\*Use 150 to 200 pounds Quali-Pro Oxadiazon 2G (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

Note: For the control of liverwort (*Marchantia spp.*) in California, Goosegrass (*Eleusine Indica*) and Prostrate Spurge (*Euphorbia supina*) use 200 pounds Quali-Pro Oxadiazon 2G (4.0 pounds of active ingredient) per acre.

**CROP TOLERANCE:**

Quali-Pro Oxadiazon 2G is recommended for use on a wide variety of container and field grown woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G may be used at the recommended rates on the following ornamental species:

<b>CONTAINER GROWN</b>	
<b>ESTABLISHED</b>	
<b>Species</b>	<b>Varieties</b>
Agapanthus	Africanus
Andromeda	Japanese
Arborvitae	American, Eastern, Oriental, Aureanana, Globe, Pyramidal, Woodward
Arctostaphylos	Fred Oelher
Azalea	Hino Red, Red, Boudoir, Snowball, Karen, Kaempp, Stuartonian, Mollishybird, Sweetheart Supreme, Salmon Beauty, Caroline Gable, Louise, Stewartstonian, Rosebud
Barberry	Redleaf, Marshall Red, Darwinii, Irwinii, Purpleleaf
Boxwood	Common, Japanese
Callistemon	Bottle-brush
Camellia	Glen #40, Sasanqua, Colonel Fiery
Ceanothus Cedar	Atlas, Portoford (Allumii)
Cotoneaster	Paineyi, Bearberry, Coral Beauty, Greyleaf, Cranberry, Silverleaf, Peking, Bright Bead, Dammeri Skolgholm, Microphylla, Lowfast, Glaucophylla.
Cypress	Monterey
Dogwood	Red osier
Douglas Fir	Pseudotsugamenziesii
Eucalyptus	Sideroxylon, Nicholli
Euonymus	Greenlane, Microphylla Variegata, Emeraldgaety, Dwarf, Gold spot, Golden Green, Silver King, Wintercreeper Vegetus, Ermerald Cushion, Spreading
Forsythia	Lynwood gold
Gardenia	Amy Yashioka, August Beauty, Mystery
Ginko	Biloba
Heath Heather	Mrs. Maxwell
Hedera	English Ivy
Honeysuckle	Purpleleaf, Purpurea
Holly	Dwarf Okinowan, Convexa, Rotundifolia, Hetzi, Helleri, Burfordi, English,

	Hellus, Dwarf Yaupon, Nellie R. Stevens, Boxleafod, Green Island, Japanese
Jasmine	Yellow Star, Star
Juniper	Barharbor, Wiltoni, Torulosa, Pfitzeriana Compacti, Pfitzeriana Aurea, Torulosa Variegata, Pfitzeriana Glauca, Gulf Stream, Tam, Armstrongii, Shore Andora, San Jose, Blue Haven, Stricta, Jade River, Broadmoor, Meyeri, Rocket, Plumosa, Royal Beauty, Praecox, Greek, Apiculata Blackburn, Plumosa Compacta, Procumbens, Blue Pfizer
Justica	Brandegeana
Leucothoe	Catesbaei, Axillaris
Myrtle	Compacta Noellii
Magnolia	Dr. Merrilli, Smauel Somers, Grandiflora
Mahonia	Oregon Grape-Holly
Osmanthus	Fortunes
Periwinkle	Vinca Minor
Photinia	Frazerii
Pine	Slash, Japanese Black, Montana, Scotch, Digger, Shore, Thumbergii
Pittosporum	Variegated, Japanese
Podocarpus	Macrophylla
Privet	Japanese, Waxleaf, Glossy, Vicary, Golden Vicary, Curly Leaf
Pyracantha	Victory, Scarlet Firethorn, Wyatti, Graberri
Quercus Raphiolepis	India Hawthorn
Rhododendron	Cunningham, Blue Diamond, Snow, Christmas, Chionoides, 'PJM', Dr. Rutgers, Coral Bells, President Clay
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong
Star Jasmine	Yellow Star Jasmine
Syzygium	Paniculatum
Viburnum	Sandankwa, Laurestinus, David
Xylosma	Congestum

**NEWLY TRANSPLANTED**

Species	Varieties
Andromeda	Japanese
Arborvitae	Glove, Woodward
Azalea	Hindo Red, Louise, Rosebud, Stewartstonian
Barberry	Marshall Red, Darwinii, Irwinii, Purpleleaf
Cedar	Portoford (allumii)
Cotoneaster	Cranberry, Peking, Coral Beauty, Bright Bead, Paineysi, Dammeri Skolgholm, Microphylla, Glaucohylla
Cypress	Monterey
Dogwood	Red Osier
Euonymus	Greenlane, Emerald Cushion, Spreading
Forsythia	Lynwood Gold
Holly	Rotundifolia, Helli, English, Japanese
Juniper	Plumosa, Plumosa Compacta, Old Gold, Pfitzeriana Aurea, Torulosa, Armstrongii, Tam, Wiltoni, Procumbens
Leucothoe	Axillaris
Magnolia	Dr. Merrilli
Mahonia	Oregon Grape-Holly
Pine	Slash
Privet	Japanese, Waxleaf, Glossy
Pyracantha	Graberri
Rhododendron	Snow, Blue Diamond, Chionoides, 'PJM', Dr. Rutgers
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong

FIELD GROWN	
ESTABLISHED	
Species	Varieties
Arborvitae	Pyramidal
Ash	Green
Azalea	Hinodi geri
Barberry	Red, Mentor
Birch	Clump, Paper
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, Tom Knudsen, Crimson Robe, R.D. Wheeler
Cotoneaster	Lowfast, Skolgholm
Crabapple	Flowing, Toringo
Dogwood	Redstemmed, Red, Flowering
Euonymus	Burning Bush, Dwarf Burning Bush, Jewell
Forsythia	Showy Boarder, Lynngold, Lynwood Gold, Goldenbell
Holly	Boxleafed, Green Island, Japanese
Honeysuckle	Zabeli
Juniper	Plumosa, Mint Julip, Arcardia, Variegata; Plumosa Compacta, Andora, Wiltoni, Hetzi
Lilac	Persian, Dwarf Korean
Maple	Amur
Oak	Red
Perwinkle	Vinca Minor
Pine	Austrian, Scotch, Norway, Stone, Japanese, Black, Loblolly, Slash, Eastern White
Pittosporum	Undulatum
Privet	California, Japanese
Rhododendron	Rosebud
Rose	Texas Centennial, Sutters Gold, Show Girl
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Taunton Spreading Yew, Hicks
Tsuga	Canadian Hemlock
Viburnum	European Cranberry Bush
Xylosma	
Ground Covers	
Carpobrotus	Edule
Osteo	
Spermum Sedum	Album
Algerian Ivy	
Vinca Major	
English Ivy	
Pachysandra	Terminalis
Ajuga	Reptans
Euonymus	Fortunei Radicans
NEWLY TRANSPLANTED	
Species	Varieties
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, R.D. Wheeler, Tom Knudsen, Crimson Robe
Dogwood	Redstemmed, Red, Flowering
Forsythia	Showy Border, Lynwood Gold, Goldenbell
Hedera	English Ivy
Holly	Green Island, Japanese
Juniper	Plumosa, Andora, Wiltoni, Hetzi



Oak	Red
Periwinkle	Vinca Minor
Pine	White Austrian, Eastern White
Privet	California, Japanese
Rhododendron	Rosebud
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Hicks
Tsuga	Canadian Hemlock

**ADDITIONAL APPLICATIONS**

Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year) except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).

Do not apply more than 200 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 4 lbs. a.i./A) in a single application.

For continued weed control, additional application of Quali-Pro Oxadiazon 2G may be made 60 to 120 days after the previous application in the following situations:

2 REPEAT TREATMENTS					
Container Grown Established		Container Grown Newly Transplanted		Field Grown Established and Newly Transplanted	
Species	Varieties	Species	Varieties	Species	Varieties
Abelia	Grandiflora	Common Boxwood		Holly	Nellie R. Stevens, Dwarf Buford, Buford
Common boxwood		English Ivy			
English Ivy		Juniper	Tam		
Holly	Helleri	Pine	Slash		
Juniper	Tam	Rhododendron	Obtusa, Kurume Formosa		
Pine	Slash				
Rhododendron	Obtusa, Kurume Formosa, Kurume Snow				
3 REPEAT TREATMENTS					
Container Grown Established					
Species			Varieties		
Abelia			Grandiflora		
Holly			Helleri, Rotundifolia		
Rhododendron			Kurume Formosa, Kurume Snow		
Greenhouse Grown Established					
Species			Varieties		
Rose			Sonia, Forever Yours, Coed, Golden Wave		
Forest Nurseries Seedlings- At least 4 weeks old					
Species			Varieties		
Pine			Loblolly, Slash, Eastern White		

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in dry area. Do not store near other pesticides, or seeds.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In case of spill, fire, or leak, contact CHEMTREC at 1-800-424-9300.

**LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Farmsaver.com, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Farmsaver.com, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Farmsaver.com, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Farmsaver.com, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Farmsaver.com, LLC's election, the replacement of product.

Quali-Pro® is a registered trademark of FarmSaver.com, LLC.

Quali-Pro Oxadiazon 2G (73220-11)(EPA app 1-26-07)(notif to EPA 4-23-08)

