ACTINOVATE [®] SOLUBLE (ALTERNATE BRAND NAME: ACTINOVATE® SP) MASTER LABEL

Sublabel A	Turf,	Greenhouse,	Nursery and	Landscape Use
------------	-------	-------------	-------------	---------------

Sublabel B Agriculture Production Use

Sublabel C Residential Use

Optional Label Claims

ACTIVE INGREDIENT:

Streptomyces lydicus*

......00.0371%

OTHER INGREDIENTS:

.....99.9629%

Information regarding the contents and levels of metals in this product is available on the Intenet at http://www.aapfco.org/metals.htm

KEEP OUT OF REACH OF CHILDRE N

CAUTION

See back panel for additional precautionary statements.

US Patent Number: 5,403,584

EPA Reg. No.: 73314-1

EPA Establishment No.: 73314-TX-001


Manufactured by:

Natural Industries, Inc.

6223 Theall Road

Houston, Texas 77066

Questions? (888) 261-4731



^{*}End-use product contains not less than 1 X 10⁷ colony forming units per gram Streptomyces (vdicus WYEC 108

Sublabel A: Turf, Greenhouse, Nursery, and Landscape Use

ACTINOVATE [®] SOLUBLE (ALTERNATE BRAND NAME: ACTINOVATE® SP)

	CAUTION
	KEEP OUT OF REACH OF CHILDREN
Information regarding the contents	and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm
*End-use product contains not less	than 1 X 10 ⁷ colony forming units per gram Streptomyces lydicus WYEC 108
OTHER INGREDIENTS: .	99.9629%
Streptomyces lydicus* .	
ACTIVE INGREDIENT:	

See back panel for additional precautionary statements.

US Patent Number: 5,403,584

EPA Reg. No.: 73314-1

EPA Establishment No.: 73314-TX-001

Manufactured by:

Natural Industries, Inc.

6223 Theall Road

Houston, Texas 77066

Questions? (888) 261-4731

Net Contents: 18-oz

PRECAUTIONARY STATEMENTS

Hazard to humans and domestic animals: Caution. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE): Applicators and other handlers must wear long-sleeved shirt and long pants, and shoes plus socks. Mixer/loaders and applicators must wear a dust/mist-filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow manufacturer's instructions for cleaning /, maintaining PPE. Keep and wash PPE separately from other laundry.

User Safety Requirements: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. If gloves are worn, wash the outside of gloves before removing. Users should remove clothing immediately if pesticide gets inside. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

For foliar application, such as foliar greenhouse, nursery, landscape and turfgrass, do not allow worker entry into treated areas during the restricted entry interval (REI) of one (1) hour or until solution has dried. Exception: If the product is soil incorporated (such as landscape, ornamental bulbs, seed coating and drench) the Worker Protection Standard, under certain circumstances, allows workers to enter treated area if there is no contact with anything that has been treated.

PPE requirements for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is: coveralls, waterproof gloves and shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of the product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

General Information:

Actinovate® Soluble is a biological fungicide for the suppression/control of root rot and damping-off fungi and the suppression/control of foliar fungal pathogens. When used as a soil drench, soil borne fungi controlled include Fusarium, Rhizoctonia, Pythium, Phytophthora, Phytomatotricum, Aphanomyces, Monosporascus, Armillaria, Sclerotinia, Postia, Verticillium, Geotrichum, and other root-decay fungi. The active ingredient in Actinovate® Soluble colonizes the root system and protects it from harmful fungi. When used as a spray, Actinovate® Soluble effectively suppresses/controls foliar diseases such as Powdery and Downy Mildew, Greasy Spot, Botrytis. Sclerotinia, Alternaria, and others.

When applied to the soil Actinovate® Soluble also breaks down minerals and micronutrients making them more available to plants resulting in increased size and vitality. Plants and turf treated with Actinovate® SP as a soil drench will become hardier, more vigorous and have a robust and protected root system.

APPLICATION DIRECTIONS:

Compatibility

Actinovate® Soluble is completely soluble and does not require agitation to keep suspended in a solution. Actinovate® Soluble is compatible with chemical fungicides, insecticides and fertilizers. If tank mixes are desired, observe all directions, precautions and limitations on labeling of all products used. Actinovate® Soluble can be tank mixed with all chemical fungicides, insecticides, and fertilizers unless otherwise restricted. Avoid tank mixing with bactericides. Avoid applying bactericides and soil furnigants to the areas treated with Actinovate® Soluble. Consult manufacturer for compatibility questions.

Application Timing

Apply Actinovate® Soluble throughout the growing season from early spring to late fall on all types of turf, ornamental, and greenhouse vegetable crops.

Note: Actinovate® SP contains live spores of a microbe that should be used prior to disease onset to obtain best results. Actinovate® SP becomes active in soil or on the plant foliage when the temperatures are above 45° F. and is not effective when temperatures remain cold. Actinovate® SP can be applied to sterilized or furnigated soil but it must be applied after sterilization or furnigation.

Application Uses:

Actinovate® Soluble is a biological fungicide for use in soil and seed coating applications (drench and in furrow) and greenhouse/nursery foliar applications in or on all raw agricultural commodities, food and fiber crops, (greenhouse use only), ornamentals, landscape plants, including tree seedlings for transplanting to the field.

GREENHOUSE, NURSERY OR INTERIORSCAPE DRENCH

Mix 4-6 oz. of Actinovate® Soluble in 100 gallons of water to create solution. Apply solution as a drench to plants/growing media at a rate of 1 gallon per cubic foot of growing media (this equates to enough solution to saturate soil without creating run-off).

For smaller quantities: Use I teaspoon of Actinovate SP per 2 gallons of water to create solution and apply as above.

Actinovate® SP can be applied through low pressure watering nozzles such as fan nozzles, through overhead boom type sprayers or sprinklers, hydroponics systems, injectors, flood benches or other drench watering systems. Actinovate® Soluble is compatible with chemical fungicides, insecticides, and fertilizers as well as other biological products.

GREENHOUSE, NURSERY, LANDSCAPE AND INTERORSCAPE FOLIAR SPRAYS

For preventative suppression/control of powdery mildew, downy mildew, *Botrytis*, Greasy Spot, *Sclerotinia*, and *Alternaria* and other foliar diseases on indoor greenhouse crops.

Apply 6-12 oz Actinovate® Soluble per acre. Dissolve Actinovate® Soluble in 50-100 gallons of water and apply to foliage and blossoms every 7 to 14 days depending on disease pressure. Crop size, spray equipment and local practice will determine the volume of water needed. Spray to wet, but avoid run-off. Actinovate® SP can be applied using hand-held backpack or ground spray equipment. Clean application equipment before use of this product, and use prepared sprays within 4 hours of preparation.

For best results use a non-ionic spreader-sticker in conjunction with application. Consult manufacturer or sales representative for specific product recommendations.

Cutting or Bare Rooted Transplant Dip:

Dip cuttings or transplants in Actinovate® Soluble dry powder or in a solution of 18-oz Actinovate® Soluble and 5 gallons water. For larger trees or woody ornamental, dissolve 18-oz of Actinovate® Soluble in 50 gallons of water and let bare root plants soak in the solution for 4-12 hours prior to planting. Plant treated cuttings or transplants in potting mix or soil in the usual manner.

6 2 23

Ornamental Bulb Crops:

Bulb Soak: Soak bulbs in solution of Actinovate® Soluble at 6-18 oz per 100 lbs. of bulbs. Thoroughly cover all surfaces of bulbs with solution

for 1 hour prior to planting).

Soil Drench: Apply to soil through irrigation or as an in-furrow seed spray at a rate of 6-12 oz per acre.

Bulb Dusting: Prior to Planting Or Shipping: Evenly dust bulbs at a rate of 2-6 oz per 100-lb of bulbs

TURF GRASS:

Mix Actinovate® Soluble with appropriate amount of water (100-125 gallons per acre). Apply at a rate of 54-oz of Actinovate® Soluble per acre

for initial application or problem areas. Apply maintenance applications of 18 oz. per acre every 4 to 8 weeks during the growing season. For

best results use a wetting agent to best move the solution to the root zone of the turf. Consult manufacturer for product recommendations.

For smaller application areas:

For initial application or problem area use 3 tablespoons (36 grams) of Actinovate® Soluble dissolved in 5 gallons of water to create a solution.

Apply solution to soil at a rate of 5 gallons per 1,000 sq. ft. of turf grass.

Reapply every 6-8 weeks as a maintenance application. Use 1 tablespoon (12 grams) of Actinovate® Soluble dissolved in 5 gallons of water to

create a solution. Apply solution to soil at a rate of 5 gallons per 1,000 sq. ft. of turf.

LANDSCAPE

For Transplants, Installations and Established Plants: Dissolve 1 teaspoon per 2 gallons of water to create a solution. Apply solution to soil

around plant's root system until soil is saturated without creating a run-off. Apply to plant before, during or after transplant.

For foliar spray dissolve 1-2 teaspoons per 2 gallons of water and apply in order to acquire thorough uniform coverage. Volume of solution

needed will vary with crop type and growth.

GREENHOUSE AND NURSERYCHEMIGATION

Apply Actinovate® Soluble at 4-6 oz per 100 gallons of water. Actinovate® Soluble may be applied through 1) overhead boom and mist-type

systems, sprinklers such as impact or micro-sprinklers 2) pressurized drench (flood) or drip (trickle) systems 3) micro irrigation such as spaghetti-

tube or individual tube irrigation, 3) hand-held calibrated irrigation equipment such as hand-held wand with injector and 5) ebb and flow systems,

Do not apply this product through any other type of irrigation system.

Plant injury or lack of effectiveness can result from non-uniform distribution of treated water. Questions concerning calibration should be

directed to your State Extension Service Specialist, the equipment manufacturer or other expert.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for

public water systems are in place.

Actinovate Soluble EPA # 73314-1

Master Label

Last Updated 11-17-06

hrb

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

If you have any questions regarding calibrations, contact equipment manufacturers, state extension specialists or other experts.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a dry, cool place out of direct sunlight and away from heat sources. Keep from overheating or freezing. Optimum storage temperature is 40° F to 85° F.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty package into application equipment. Then dispose of empty package in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Limited Warranty / Disclaimer

Manufacturer warrants that this product is suited for the labeled use when applied according to label directions. Manufacturer makes no warranty of merchantability. There are no warranties that extend beyond the description on this label and in no event shall manufacturer be liable for any consequential damages.

Actinovate Soluble EPA # 73314-1

Master Label

Version 7.1

Last Updated 11-17-06
hrb

Manufactured By:

Natural Industries, Inc.

6223 Theall Rd.

Houston, TX 77066 .

Questions? (888) 261-4731

www.naturalindustnes.com

BEST IF USED BY:

LOT NUMBER:

SUBLABEL B: AGRICULTURE PRODUCTION USE

ACTINOVATE ® SOLUBLE (ALTERNATE BRAND NAME: ACTINOVATE® AG)

ACTIVE INGREDIENT:	
Streptomyces lydicus*	00.0371%
OTHER INGREDIENTS:	99.9629%
End-use product contains not	less than 1 X 10 ⁷ colony forming units per gram Streptomyces lydicus WYEC 108

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

US Patent Number: 5,403,584

EPA Reg. No.: 73314-1

EPA Establishment No.: 73314-TX-001

Manufactured by:

Natural Industries, Inc.

6223 Theall Road

Houston, Texas 77066

Questions? (888) 261-4731

Net Contents: 18-oz

PRECAUTIONARY STATEMENTS

Hazard to humans and domestic animals: Caution. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Actinovate Soluble EPA # 73314-1

Personal Protective Equipment (PPE): Applicators and other handlers must wear long-sleeved shirt and long pants, and shoes plus socks. Mixer/loaders and applicators must wear a dust/mist-filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization. Follow manufacturer's instructions for cleaning / maintaining PPE. Keep and wash PPE separately from other laundry.

User Safety Requirements: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. If gloves are worn, wash the outside of gloves before removing. Users should remove clothing immediately if pesticide gets inside. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

For foliar application do not allow worker entry into treated areas during the restricted entry interval (REI) of one (1) hour or until solution has dried. Exception: If the product is soil incorporated (such as landscape, ornamental bulbs, seed coating and drench) the Worker Protection Standard, under certain circumstances, allows workers to enter treated area if there is no contact with anything that has been treated.

PPE requirements for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is: coveralls, waterproof gloves and shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of the product that are not within the scope of the Worker Protection Standard for agricultural

pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or

greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

General Information:

Actinovate® Soluble is a biological fungicide for the suppression/control of root rot and damping-off fungi and the suppression/control of foliar

fungal pathogens. When used as a soil drench, soil borne fungi controlled include Fusarium, Rhizoctonia, Pythium, Phytophthora,

Phytomatotricum, Aphanomyces, Monosporascus, Armillaria, Sclerotinia, Postia, Verticillium, Geotrichum, and other root-decay fungi. The

active ingredient in Actinovate® Soluble colonizes the root system and protects it from harmful fungi. When used as a spray, Actinovate®

Soluble effectively suppresses/controls foliar diseases such as Powdery and Downy Mildew, Botrytis, Monilinia, Anthracnose, Greasy Spot,

Sclerotinia, Alternaria, Erwinia and others.

When applied to the soil Actinovate® Soluble also breaks down minerals and micronutrients making them more available to plants resulting in

increased size and vitality. Plants and lawns treated with Actinovate® AG as a soil drench will become hardier, more vigorous and have a robust

and protected root system.

INTEGRATED PEST MANAGEMENT (IPM)

Integrate Actinovate® Soluble into an overall disease and pest management strategy whenever fungicide use is necessary. Follow practices

known to reduce disease development. Consult local agricultural authorities for specific IPM strategies developed for your crop(s) and location.

USE RATE DETERMINATION -Agricultural Use

Carefully read and follow all label directions, use rates and restrictions. Actinovate® Soluble should be applied prior to or in the early stages of

disease development. For proper foliar application, determine the number of acre to be treated, the recommended label use rate and select the

appropriate gallonage to give thorough uniform coverage of all plant parts to be protected.

For proper soil application, determine the number of acre to be treated, the recommended label use rate and select the appropriate gallonage to

give good saturation of the soil in order for the product to establish itself on the root system. For best results apply product solution to damp soil.

Prepare only the amount of spray or soil drench solution to treat the measured area. Accurate spray equipment calibration is essential prior to use.

PREHARVEST INTERVAL - Agricultural Use

Actinovate® Soluble can be applied up to and including the day of harvest

Actinovate Soluble EPA # 73314-1 Master Label

APPLICATION DIRECTIONS:

Compatibility

Actinovate® Soluble is completely soluble and does not require agitation to keep suspended in a solution. Actinovate® Soluble is compatible

with chemical fungicides, insecticides and fertilizers. If tank mixes are desired, observe all directions, precautions and limitations on labeling of

all products used. Actinovate® Soluble can be tank mixed with all chemical fungicides, insecticides, and fertilizers unless otherwise restricted.

Avoid tank mixing with bactericides. Avoid applying bactericides and soil furnigants to the areas treated with Actinovate® Soluble. Consult

manufacturer for compatibility questions.

Application Timing

Apply Actinovate® Soluble throughout the growing season from early spring to late fall on all types of production agriculture crops

Note: Actinovate® Soluble contains live spores of a microbe that should be used prior to disease onset to obtain best results. Actinovate® Soluble

becomes active in soil or on the plant foliage when the temperatures are above 45° F. and is not effective when temperatures remain cold.

Actinovate® Soluble can be applied to sterilized or furnigated soil but it must be applied after sterilization or furnigation.

Application Uses:

Actinovate® Soluble is a biological fungicide for use in soil applications (drench and in furrow) and foliar applications in or on all raw

agricultural commodities, food and fiber crops, and other agricultural production crops.

AGRICULTURE PRODUCTION

For soil treatment for the control of Pythium, Rhizoctonia, Phytopthora, Fusarium, Verticillium, and other root decay fungi.

For foliar treatment of Powdery Mildew, Botrytis, Schlerotinia, Monilinia, Greasy Spot, Alternaria, Downy Mildew and other foliar fungi.

Soil Treatment At Planting:

Use at planting, seeding, or transplant. Apply 3-12 oz of Actinovate Soluble, dissolve in appropriate amount of water, per acre.

Soil Treatment Through Irrigation:

Actinovate® Soluble may be used in drip, overhead or other type of irrigation system at any stage of the plant growth as a soil treatment. Use 3-12 oz of Actinovate®

Soluble, dissolved in enough water, per acre.

Foliar Treatment

Use 3-12 oz of Actinovate® Soluble in 20-150gallons of water per acre. Apply initial application prior to onset of disease season. Reapply every 7-14 days

depending on disease pressure and environmental conditions. For best results use a spreader-sticker (adjuvant) in conjunction with product application.

Crops on which Actinovate® Soluble may be used

CROPS

Soil Drench Rate

Foliar Spray Rate

Agronomic Field and Row Crops:

3-oz per acre at rooting

3-12 ounces per acre.

buckwheat, com (all types), cotton, canola, safflower,

Reapply every 7-14 days

sunflower, succulent and dry peas, peanuts, soybeans

For best results use

and other agronomic field and row crops

spreader-sticker.

Alfalfa Hay and Forage:

Alfalfa, clover, vetch, trefoil

Cucurbit Vegetables:

3-12 oz per acre

3-12 ounces per acre.

cucumbers, melons, gourds, squash, cantaloupe, and other curcubits

Reapply every 7-14 days

For best results use

Fruiting Vegetables:

with spreader-sticker.

Eggplant, sweet peppers,

hot peppers, tomatoes, tomatillos and other fruiting vegetables

Herb, Spices and Mints:

sage, rosemary, thyme,

peppermint, dill, basil. Oregano and other herbs and spices

Leafy Vegetables and Cole Crops:

Broccoli, brussel sprouts, cabbage, cauliflower, celery, collards,

endive, kale, kohlrabi, lettuce, mustard greens

parsley, spinach and other leafy vegetable crops

Legume and Vegetable Crops:

snap and dry beans, leutils, succulent and dry peas, peanuts,

soybeans and other legume/vegetable crops

Small Grains:

rye, sorghum, millet and other small grains

Root/Tuber and Bulb Crops:

Garlic, onions, white potatoes, sweet potatoes, carrot, ginger, ginseng,

horseradish, turnip, beets, radish and other root/tuber/bulb crops

Berry Crops

Blueberries, blackberry, raspberry, loganberry, huckleberry,

gooseberry, elderberry, currant, caneberry and other berry crops

Strawberry

Actinovate Soluble EPA # 73314-1

Master Label Version 7.1 Last Updated 11-17-06 hrb

Asparagas
Citrus
Orange, grapefruit, lemon, tangerine, tangelo, lime,
pummelo and other citrus crops
Corn
Popcora, seed corn, sweet corn and other corn crops
Grape
Wine grapes, table grapes, raisons and other grape crops
Hops
Pome Fruit
Apple, crabapple, pear, quince, mayhaw and other pome fruit
Stone Fruit
Apricot, cherry, nectarine, peach, plum, prune and other stone fruit
Tree Nuts
Almood, pistachio, pecan, walnut, filberts and other tree nuts
Tropical Fruits
Avocado, mango, papaya and other tropical fruits
Bananas/ Plantains
Watercress
Mushrooms

CHEMIGATION

Apply 3-12 ounces of Actinovate® Soluble in 20-200 gallons of water per acre. Actinovate® Soluble may be applied through 1) overhead boom and mist-type systems, sprinklers such as impact or micro-sprinklers 2) pressurized drench (flood) or drip (trickle) systems 3) micro irrigation such as spaghetti-tube or individual tube irrigation, 3) hand-held calibrated irrigation equipment such as hand-held wand with injector and 5) ebb and flow systems, 6) sprinkler and drip irrigation systems including central pivot, lateral move, end tow, side wheel roll, traveler, solid set or hand move systems.

Do not apply this product through any other type of irrigation system.

Actinovate Soluble EPA # 73314-1

Master Label

Version 7.1

Last Updated 11-17-06
hrb

Plant injury or lack of effectiveness can result from non-uniform distribution of treated water. Questions concerning calibration should be

directed to your State Extension Service Specialist, the equipment manufacturer or other expert.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for

public water systems are in place.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to

prevent water source contamination from back flow.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of a responsible person, shall shut

the system down and make necessary adjustments should the need arise.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the

injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the

injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is

either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor

stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure

decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed

of materials that are compatible with pesticides and capable of being fitted with a system interlock.

If you have any questions regarding calibrations, contact equipment manufacturers, state extension specialists or other experts.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a dry, cool place out of direct sunlight and away from heat sources. Keep from overheating or freezing. Optimum

storage temperature is 40° F to 85° F.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Actinovate Soluble EPA # 73314-1

Master Label

Container Disposal: Completely empty package into application equipment. Then dispose of empty package in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Limited Warranty / Disclaimer

Manufacturer warrants that this product is suited for the labeled use when applied according to label directions. Manufacturer makes no warranty of merchantability. There are no warranties that extend beyond the description on this label and in no event shall manufacturer be liable for any consequential damages.

Manufactured By:

Natural Industries, Inc.

6223 Theall Rd.

Houston, TX 77066

Questions? (888) 261-4731

www.naturalindustries.com

BEST IF USED BY:

LOT NUMBER:

Sublabel C: Residential Use

ACTINOVATE ® SOLUBLE (ALTERNATE BRAND NAME: ACTINOVATE® LAWN & GARDEN)

ACTIVE INGREDIENT:	
Streptomyces lydicus*	00.0371%
OTHER INGREDIENTS:	99.9629%
End-use product contains not le	ess than 1 X 10 ⁷ colony forming units per gram Streptomyces lydicus WYEC 108

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements.

US Patent Number: 5,403,584

EPA Reg. No.: 73314-1

EPA Establishment No.: 73314-TX-001

Manufactured by:

Natural Industries, Inc.

6223 Theall Road

Houston, Texas 77066

Questions? (888) 261-4731

Net Contents: 18-oz

FIRST AID				
If inhaled: • Move person to fresh air.				
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. 			
	 Call a poison control center or doctor for further treatment advice. 			
Have the product container or	label with you when calling a poison control center or doctor, or going for treatment.			

PRECAUTIONARY STATEMENTS

Hazard to humans and domestic animals: Caution. Harmful if inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

General Information:

Actinovate® Soluble is a biological fungicide for the suppression/control of root rot and damping-off fungi and the suppression/control of foliar fungal pathogens. The active ingredient in Actinovate® Soluble is a patented bacterium that, when applied, grows around the root system (when soil drenched) and foliage of the plant (when sprayed on). Actinovate® Lawn and Garden uses several novel modes of antifungal action to protect plants and lawns. Actinovate® Soluble may be used on all vegetables, fruits, and nuts including tomatoes, peppers, melons, carrots, broccoli, lettuce, onions, apples, pears, and walnuts in addition to annual and perennial bedding plants and flowers, roses, potted flowers, foliage plants, trees, shrubs and lawns located in residential homes, residential greenhouses and residential and commercial landscapes. Actinovate® Soluble can be applied up to and including the day of harvest.

In addition, when applied to the soil, Actinovate® Soluble also breaks down minerals and micronutrients making them more available to plants resulting in increased size and vitality. Plants and turf treated with Actinovate® Soluble as a soil drench will become hardier, more vigorous and have a robust and protected root system.

Diseases Controlled/Suppressed

Soil Diseases (When applied as a drench):

Foliar Diseases (When applied as a spray)

*Root Rot and damping off in ornamentals & vegetables

*Powdery and downy mildew

(Pyhium, Phytophthora, Rhizoctonia, Fusarium et.al)

*Grey Mold (Botrytis)

*Turf Brownpatch (Rhizoctonia)

*Black Spot (Diplocarpon rosae)

*Turf Dollarspot (Sclerotinia)

*Leaf Spots and Rusts

*Turf Take-all patch (Gaeumannomyces graminis)

*Fire Blight (Erwinia)

*And others

*And others...

APPLICATION DIRECTIONS:

Compatibility

Actinovate® Soluble is completely soluble and does not require continuous mixing to keep suspended in a solution. Actinovate® Soluble is compatible with all fungicides, insecticides and fertilizers, unless otherwise restricted.. If tank mixes are desired, observe all directions, precautions and limitations on labeling of all products used. Avoid tank mixing with bactericides. Avoid applying bactericides to the areas treated with Actinovate® Soluble.

DIRECTIONS FOR USE:

SOIL DRENCH

Dissolve I teaspoon (4-grams) of Actinovate Soluble per 2 gallons of water to create solution.

Apply solution to pot, base of plant, or roots of plant by watering until soil is completely saturated without creating run-off. One cup of solution usually treats about one 6" pot or it's equivalent.

For best results apply to damp soil and/or apply in conjunction with a wetting agent. Pre-dampened soil will allow the Actinovate® Soluble microorganism to work to the root system of the plant much more easily.

FOLIAR SPRAY

Dissolve 1 teaspoon (4-grams) of Actinovate Soluble per 2 gallons of water to create solution.

Spray leaves, stems, and new shoots to runoff providing complete coverage of entire plant. For best results, apply product prior to disease development or at the first sign of infection. Repeat at 7-day intervals to protect new foliage. Under conditions of heavy rainfall, it may be necessary to reapply the product after the rain has stopped. Do not water foliage within 4 hours of application. Use a pump bottle, handheld pump, backpack or similar type of spray equipment.

For best results use a spreader-sticker (a product designed to break the surface tension of water and evenly spread it over the surface of the foliage) in conjunction with application. Consult your garden center or dealer for specific product recommendations. (Examples of spreader-stickers include Weather Shield®, Agri-2® and yucca extract)

Actinovate Soluble EPA # 73314-1 Master Label Version 7.1 Last Updated 11-17-06 hrb

LAWN TREATMENT:

Dissolve Actinovate® Soluble in the appropriate amount of water to create a solution. Apply solution to soil at a rate of 5 gallons per 1,000 sq. ft. Use 3 tablespoons of Actinovate® Soluble per 1,000 sq ft for initial application or problem areas. Use 1 tablespoon per 1,000 sq. ft for maintenance every 4-8 weeks. For best results use a wetting agent, such as yucca extract, Coco-Wet®, or Mega Wet®, in conjunction with Actinovate® Soluble in order to help move product to the root system of the lawn. Consult your garden center or dealer for specific product recommendations.

LAWN APPLICATION CHART:

Amount of Actinovate® Soluble For Initial Application/Problem Area	Amount of Actinovate® Soluble For Maintenance Application	Amount of Water	Area To Treat	Application
3 tablespoons (36 grams) .	l tablespoon (12 grams)	5 gallons		Use watering device or hose end sprayer to apply until soil is saturated. Reapply at maintenance rate every 4-8 weeks.

HOSE-END SPRAYER SETTINGS FOR LAWNS

Initial Application/Problem Area	Maintenance Application	Water to Add to Sprayer Jar	Dial Setting	Area Treated
3 tablespoons (36 grams) of Actinovate® Soluble	l tablespoon (12 grams) of Actinovate® Soluble	32-oz	1:20 (1-oz per 20-oz) 32-oz per 5 gallons	1,000 sq ft
15 tablespoons (180 grams) of Actinovate® Soluble	5 tablespoon (60 grams) of Actinovate® Soluble	32-oz	1:100 (1-oz per 100-oz) 32-oz per 25 gallons	5,000 sq ft

STORAGE AND DISPOSAL

STORAGE: Store in a dry, cool area inaccessible to children. Store in original container only. Keep container sealed when not in use. CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Limited Warranty / Disclaimer

Manufacturer warrants that this product is suited for the labeled use when applied according to label directions. Manufacturer makes no warranty of merchantability. There are no warranties that extend beyond the description on this label and in no event shall manufacturer be liable for any consequential damages.

Manufactured By:

Natural Industries, Inc.

6223 Theall Rd.

Houston, TX 77066

Questions? (888) 261-4731

www.naturalindustries.com

BEST IF USED BY:

LOT NUMBER:

Label version: 7.0

OPTIONAL LABEL CLAIMS

- Biological Fungus Control.
- Controls soil-borne plant diseases such as *Pythium, Rhizoctonia, Phytophthora, Verticillium, Fusarium* and others.
- Suppresses and Controls Foliar Disease such as Powdery Mildew, Downy Mildew, Greasy Spot, *Botrytis, Sclerotinia, Monilinia, Alternaria, Erwinia,* and others.
- 100% Soluble. Will not clog machinery.
- Enhances plant vitality.
- Encourages larger root systems.
- For Greenhouse, Nursery, Interiorscapes, Agriculture and Turf
- For turf diseases such as BrownPatch, Dollar Spot and Take-all Patch
- Use on Roses, Vegetables, Fruits, Flowering Plants, Trees and Shrubs
- Fungicide that attacks harmful garden diseases
- Controls foliar diseases such as Powdery Mildew, Rust, Grey Mold, Black Spot and more
- Controls damping off and other root rot diseases
- Aids in Turf Recovery
- For Home, Garden and Lawn Care Use
- Concentrated Formula
- Treats Up to 5,000 sq. ft. of lawn or 550 6" potted plants
- Easy To Use