



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 10, 2021

Mandy Slack
Sr. Analyst, Regulatory Affairs
The Scotts Company
14111 Scottslawn Road
Marysville, OH 43041

Subject: Registration Review Label Amendments for Atrazine Incorporating Mitigation Measures from the Interim Decision and the Technical Registrants' Commitments for the Endangered Species Act (ESA) Biological Evaluation
Product Name: SWISS FARMS CENTIPEDE WEED & FEED
EPA Registration Number: 73327-1
Application Date: 11/03/2020
Decision Number: 567707

Dear Ms. Slack:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Atrazine Interim Decision and with the technical registrants' commitments for the ESA Biological Evaluation. The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only

Page 2 of 2
EPA Reg. No. 73327-1
Decision No. 567707

distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Anitha Kisanga at Kisanga.Anitha@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a long horizontal flourish extending to the right.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

NOTE:

EPA Reg. No. 73327-1

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

[FRONT]

For Outdoor Residential Use Only

Swiss Farms Centipede Weed & Feed

EPA Reg. No. 73327-1

TREATS/COVERS 5,000 Sq Ft

ACTIVE INGREDIENT:

Atrazine 1.102%

Related Compounds 0.023%

OTHER INGREDIENTS 98.875%


TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back / side / top | bottom [panel / label] for additional Precautionary Statements.

NET WT. 20 lbs: [9.07 kg]/ XX lbs. [XX kg]



Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material.

Do not apply granular lawn products by hand. Avoid contact with hands and skin. Broadcast applications must NOT be made using hand-held devices such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments.

FIRST AID	
If Swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a Poison Control Center or doctor for further treatment advice.
<p>Have the product container label with you when calling a Poison Control Center or doctor, or going for treatment. Note to Physicians: Emergency information call 1-877-465-5161.</p>	

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to well drained soils where the water table (groundwater) is close to the surface and where these soils are very permeable (i.e. well drained). Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

NON-TARGET ORGANISM ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on St. Augustine [including Floratam], Carpetgrass, Centipede and Zoysia Lawns ONLY

Application Restrictions

- Do not apply this product in a way that will contact any person or pet either directly or through drift.
- Keep people and pets out of area during application.
- This product must be WATERED IN IMMEDIATELY after application.
- Do not enter or allow others (including children and pets) to enter the treatment areas (except those involved in the watering) until watering-in is complete and the surface is dry.
- Do not apply by hand or hand-held device such as hand-held rotary applicator.
- Do not apply more than 1 lb of product per 250 sq. ft. (172 lbs of product per acre), which is equivalent to 2 pounds of active ingredient per acre, per application.
- Do not make more than 2 applications per year.
- Use on roadsides; Conservation Reserve Program (CRP) land; conifers, including Christmas Tree plantings; timber; forestry; and, Miscanthus and other perennial bioenergy crops is prohibited.
- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).

Use Precautions

- Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.
- Not for use on or under trees, shrubs, bedding plants, flowers or garden plants. Do not apply under the branches (root zone area) of shrubs, trees, flowers, etc.
- Do not apply to newly seeded Zoysia, centipede, or carpetgrass lawns. Wait until they have required at least 2 mowings.
- Do not apply to flowers, bedding plants, ornamental trees, or shrubs.

General Use Tips

1. Determine size of area to be treated - measure length and width in feet, then multiply to determine coverage area in sq. ft. (length x width). **Next, subtract non-lawn areas.**
2. Use the spreader setting chart to determine the appropriate setting.
3. Treat as directed - thorough coverage is important.
4. For best results, water soil until damp.

When To Apply

- Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.
- Apply anytime broadleaf weeds are growing to get control.
- Apply in the fall to control henbit, chickweed, and other winter weeds.
- If complete control is not obtained, a second application may be necessary 2 months after the first.
- Apply on Centipede grasses after greening has occurred.

How To Apply

1. Calibrate your spreader according to the spreader manufacturer's directions. Adjust settings as needed to apply the recommended product.
2. Using a drop-type or rotary spreader, apply granules evenly over grass, avoiding streaking, skips or overlaps. Avoid using spreaders that apply product in narrow or concentrated bands.
3. Check calibration of spreader frequently to ensure spreader is functioning properly and applying product evenly.

NOTE:

EPA Reg. No. 73327-1

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

SPREADER SETTINGS

This product also contains a high quality slow-release fertilizer. This product will green your lawn in a few days and will continue feeding up to [2] [3] [4] months. For best results apply with a fertilizer spreader at the setting listed [below] [above] [at right] [at left]. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°F) water the lawn to rinse particles off the grass.

ROTARY SPREADERS		DROP SPREADERS	
Republic EZ	X	Republic EZ	X
Ortho	X	Ortho	X
Quaker	X	Precision	X
Scotts Easy Green	X	Scotts (PF)	X
Scotts Speedy Green	X	Scotts Even Green	X

Calibrate your spreader with each application. Adjust to apply at the rate of 1 lb. of product per 250 sq. ft. This 20 lb. bag covers 5,000 sq. ft. of lawn area. Do not make more than 2 applications per year.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult owner's manual. NEVER APPLY BY HAND.

This product contains iron that aids in the greening process but WILL stain sidewalks and driveways. Sweep off any product that may have drifted onto these areas.

Typically, a lawn will utilize 1 to 4 lbs. of nitrogen per 1,000 sq. ft. per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX lbs. of nitrogen per 1,000 sq. ft. Consult your local County Extension office for questions regarding a fertilizer program suitable for your geographic region.

[Spreader Tips:]

[To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn. (Graphic of pattern)]

[Always apply with a rotary or drop spreader to distribute the granules evenly.]

WEEDS CONTROLLED

This product controls I suppresses the following list of growing weeds.

Annual Bluegrass (Poa Annua)	Dichondra	Purslane
Annual Sowthistle	Dollarweed	Ragweed
Black Medic	Florida Betony	Red Sorrel
Blue Toadflax	Florida Pusley	Rescuegrass
Buckhorn Plantain	Fragrant Cudweed (Rabbit Tobacco)	Shepherd's-purse
Burclover	Henbit	Spotted Spurge
Carolina Geranium	Knotweed	Spurweed (lawn burweed)
Chickweed, Common	Lambsquarters	Stinging Nettle
Chickweed, Mouseear	Lespedeza	Venus Looking Glass
Clover	Lippia (matchweed)	Wild Mustard
Common Dandelion	Narrowleaf Vetch	
Crabgrass	Oxalis (woodsorrel)	
Cudweed	Parsley-Piert	
Cutleaf Evening Primrose	Piqweed	

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a clean, dry place. Reseal opened bag [by folding top down and securing.]

PESTICIDE AND/CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

OPTIONAL CLAIMS - Claims may appear on any or all label panel(s)

[Pesticidal Claims]

- Apply Anytime Broadleaf Weeds Are Actively Growing
- Controls Many More Weeds Listed on the [Back] [Side] of this Bag
- Controls Weeds as Listed while Feeding Your Lawn with a Quick Greening Long Lasting Fertilizer
- Enjoy a Beautiful Weed-Free Lawn
- Kills Weeds [While Feeding Your Lawn].
- Kills Dollarweed & 35 other major lawn weeds
- Kills Over 35 Types of Lawn Weeds
- Promotes Thick Green Growth and Controls Dollarweed, Dandelion, Henbit and Other Listed Weeds (see back/ side for a complete list of weeds)

[Fertilizer Claims]

- A Formula for St. Augustine Lawns
- Contains (X) % Fertilizer Nutrient for Extra Greening
- Contains (X) % Fertilizer Nutrient to Improve Deficient St. Augustine Lawns
- Delivers a Long Lasting Feeding without Burning for a Stronger Greener Lawn
- Feeds for up to [XX months] (at 72°F)
- Greens quickly and Feeds Up to 2 months
- Greens in 72 hours
- Improves deficient St. Augustine Lawns
- Quick Greening & Extended Feeding for up to 8 weeks
- Timed-Release Feeding for up to 8 Weeks
- Timed-Release Nitrogen for Extended Feeding

[General Benefit Claims]

[Use Tip Box]

["To Do" List]

- Measure lawn
- Apply at listed rate
- Water in as indicated]

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

[Turf Note]

- CENTIPEDE GRASS IS DIFFERENT FROM OTHER TURFGRASSES. It requires very little fertilizer. In fact, turf experts say that too much fertilizer can hurt centipede grass. With this in mind, we have formulated Swiss Farms Centipede Weed & Feed to give your Centipede grass the light feeding it needs to grow thick, lush and sturdy. Our exclusive timed-release nitrogen provides a slow, gradual feeding that keeps your grass greener longer and helps build a stronger, hardier root system. Extra iron helps keep the grass green without too much nitrogen.
- [Brand name] products with POLYON® timed-release nitrogen work differently than other slow-release products. POLYON® timed-release nitrogen releases in response to soil temperature rather than moisture level. Other slow release products release according to the amount of moisture present in the soil... the more moisture the quicker the nutrients release... putting your grass at risk from burning. Because [brand name] products contain POLYON® timed-release nitrogen, your grass is fed for up to XX months [at 72° F].
- [Brand name] products with POLYON® timed-release nitrogen feed as the ground temperature warms in the spring and summer, and slows down as the soil cools in the fall, thus supplying nitrogen according to the requirements of your grass.
- Timed-Release Fertilizer - Polyon® If it's not Green It's not Polyon®
- Swiss Farms timed-release nitrogen is formulated to give your grass the primary nutrients it needs to grow thick, lush and sturdy GREEN grass. Our [exclusive] timed-release nitrogen provides a slow, gradual feeding that keeps your grass greener longer and helps build a stronger, hardier root system.
- Why does your lawn need Swiss Farms Weed & Feed? Swiss Farms Weed & Feed is formulated to kill lawn weeds and give your grass [the primary] [many of the] nutrients it needs for thick growth. Our timed-release formula keeps your grass greener longer and helps build strong, hardy root systems.

GUARANTEED ANALYSIS

Total Nitrogen (N).....	XX.X.%
X.X% Ammoniacal Nitrogen X.X% Urea Nitrogen*	
Available Phosphate (P205)	XX.X%
Soluble Potash (K20).....	XX.X%

*Contains XXX% Slowly available Urea Nitrogen from polymer-coated urea

Derived From: Polymer-coated Urea, Diammonium Phosphate and Muriate of Potash.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>.

NOTE:

EPA Reg. No. 73327-1

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by/ denotes and/or options.

Optional Florida Fertilizer Application Language

Florida Fertilizer Applications: [Do not exceed/May apply up to] 4 annual feedings [4 lbs. N/1000 sq. ft. per year] with the following exceptions: On centipedegrass do not exceed 2 feedings [2 lbs. N/1000 sq. ft. per year] in the north and 3 feedings [3 lbs. N/1000 sq. ft. per year] in the central and south. On zoysiagrass do not exceed 3 feedings [3 lbs. N/1000 sq. ft. per year] in the north.

North Florida is defined as north of a line extending from Cedar Key to Daytona Beach. Central Florida is defined as south of the line extending from Cedar Key to Daytona Beach to a line extending from Tampa to Vero Beach. South Florida includes the remaining southern portion of the state.

In Florida, 4 feedings may exceed the Nitrogen limits on certain grass types in some [geographic] areas.

Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices [for feeding your lawn.]

Check with your county or city government to determine if there are local regulations for fertilizer use.

Notice: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Swiss Farms Products, Inc. No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail you a refund check promptly.

Swiss Farms Products, Inc.
3993 Howard Hughes Pky, Suite 250
Las Vegas, NV 89169-6754

Questions/Information Consumer helpline
1-800-543-TURF (8873)

©20XX Licensed by Swiss Farms Products, Inc. World rights reserved

Made in U.S.A.

EPA Reg. No. 73327-1 EPA Est. No. XXX-XX-X

[Superscript used is first letter of run code on end of container]