



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

February 10, 2022

Rebecca M. Horton  
Agent  
Ambrands  
1000 Parkwood Circle, Ste. 700  
Atlanta, GA 30339

Subject: Registration Review Label Mitigation for Imazaquin  
Product Name: IMAGE HERBICIDE CONSUMER CONCENTRATE  
EPA Registration Number: 73342-4  
Application Date: 2/3/2020  
Decision Number: 559340

Dear Ms. Horton:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Imazaquin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2  
EPA Reg. No. 73342-4  
Decision No. 559340

If you have any questions about this letter, please contact Quinn Gavin by phone at 202-566-2284, or via email at [gavin.quinn@epa.gov](mailto:gavin.quinn@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

# IMAGE® Herbicide Consumer Concentrate

[Additional Brand Name(s): IMAGE® Ready-to-Spray Herbicide Consumer Concentrate]

## Optional Marketing Statements:

Kills Nutsedge

Nutsedge Killer

Kills Really Tough<sup>†</sup> [Hard-to-Kill<sup>†</sup>] Weeds [in Lawns]

<sup>†</sup>including dandelion, dollar weed, crabgrass and others listed

[Also] Kills Dollarweed, Wild Onion, Sandbur, Annual Bluegrass

Can be used to treat listed weeds in dormant Bermudagrass

Kills tough<sup>†</sup> [hard-to-kill<sup>†</sup>] weeds in dormant Bermudagrass

<sup>†</sup>including dandelion, dollar weed, crabgrass and others listed

Kills warm and [&] cold season weeds

For use on Bermudagrass, Zoysiagrass, St. Augustinegrass and other listed grasses

Kills Yellow and Purple Nutsedge

Kills [nutsedge], [chickweed], [white clover], [dandelion], [kyllinga][Poa annua][annual bluegrass] and other listed weeds

**ONE PINT TREATS [COVERS] 4,000 SQUARE FEET**

**24 FL. OZ. TREATS [COVERS] 6,000 SQUARE FEET**

**32 FL. OZ. TREATS [COVERS] 8,000 SQUARE FEET**

## **A Selective Herbicide for Use in Established:**

Bermudagrass

St. Augustinegrass

Buffalograss

Zoysiagrass

Centipedegrass

And Selected Landscape Ornamentals and Groundcovers

## **CONTROLS NUTSEDGE Wild Onion and Sandbur**

## Optional Marketing Statements (when packaged in RTS applicator):

Connect and Spray Applicator


Treats 8,000 sq ft.

## Optional Marketing Statements when packaged in 2.5 fl oz single-cartridge for use in [AMDRO Spray delivery system sprayer]:

Contains XXX cartridges

Each 2.5 fl.oz. cartridge makes 1 Gal. of spray

Can be used to spot treat listed weeds in dormant Bermudagrass



[Optional graphic:



]

**ACTIVE INGREDIENT:**

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-3-quinolinecarboxylic acid)\*..... 3.3%

**OTHER INGREDIENTS** ..... 96.7%

**TOTAL** ..... 100.0%

\*Equivalent to 3.14% 2-[4,5-dihydro-4-methyl-4(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-3-quinolinecarboxylic acid. (1 pint contains 0.0375 pounds [lbs.] of active ingredient as the free acid)

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Back [Inside/Side/Other Panel] for Additional Precautionary Statements and First Aid.  
(See inside for additional Precautionary Statements, First Aid, and complete Directions for Use.)

**Net Contents:** \_\_\_\_\_

{BACK/SIDE/OTHER PANEL[S]}

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, shoes and socks when applying this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

**FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call (800) 265-0761 for emergency medical information.



**If inhaled** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on skin** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in eyes** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**ENVIRONMENTAL HAZARDS**

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**NON-TARGET ORGANISM ADVISORY STATEMENT:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

**GROUNDWATER ADVISORY STATEMENT:** This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**SURFACE WATER ADVISORY STATEMENT:** This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of imazaquin from

EPA Reg No. 73342-4

EPA Est. No. 5905-GA-01<sub>A</sub>, 2217-KS-02<sub>B</sub>, 59144-GA-1<sub>C</sub>,  
71106-GA-003<sub>D</sub>, 74750-GA-1<sub>E</sub>, 70815-GA-001<sub>F</sub>

Oct. 5, 2020 - revised

Page 3 of 13 Pages

runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown on golf courses or lawns and grounds.

Observe all cautions and limitations on this label. Keep children and pets out of the treated area until sprays have dried. The use of IMAGE Consumer Concentrate not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

### Storage & Disposal

**Pesticide Storage:** KEEP FROM FREEZING. Do not store below 32°F. Store in a cool, dry place inaccessible to children or pets.

**Pesticide Disposal:** *If partially filled* — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**Container Disposal:** *If empty* — Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

## PRODUCT INFORMATION

IMAGE Consumer Concentrate selectively kills (controls) tough<sup>†</sup> [hard-to-kill] weeds. Kills purple and yellow nutsedge, wild onion and garlic, dollarweed<sup>†</sup>, annual bluegrass<sup>†</sup>, and others as listed. It can be used to kill (control) weeds in bermudagrass, St. Augustinegrass, buffalograss, centipedegrass, or zoysiagrass lawn and also in certain landscape ornamentals and groundcovers (see KILL (CONTROL) OF WEEDS IN ORNAMENTALS). IMAGE is mainly absorbed by the roots of sprayed weeds and is translocated throughout the weed to provide complete kill. Complete kill should occur in 3-5 weeks.



Do not exceed labeled application rate of 4 fl. oz. per 1000 sq. ft. (0.375 lb ai per acre) per application. Repeat applications may be necessary, but, do not exceed 8 fl. oz. per 1000 sq. ft. (0.75 lb. ai per acre) annually.

{Note to EPA: to include graphics demonstrating weed kill}



1. IMAGE is absorbed into the weed directly after application
2. Weed is yellow and wilted in 1-2 weeks
3. Weed is dead in 3-5 weeks]

### MIXING INSTRUCTIONS

APPLICATION METHOD		USE RATE
<b>TRIGGER OR PUMP-UP SPRAYER</b> 	Spot Weed Treatment	Mix 2.5 fl. oz. (5 Tablespoons) in 1 gal. of water. [2 teaspoons in 16 fl. oz.]. Spray to wet.
	Large Area Treatment	Mix 0.8 fl. oz. (1-1/2 Tbs.) in 1 gallon of water to cover 200 sq. ft.
		<b>USE RATE</b>
<b>HOSE-END SPRAYERS</b> 		Use at the rate of 24 fl. oz. Image to cover 6,000 sq. ft. of lawn. Apply in sufficient water (i.e. 5 gallons per 1000 sq. ft.) to insure adequate coverage.
For ORTHO® Dial 'n Spray®, pour 24 fl. oz. Image into jar, set metering dial at 4 tsp. and uniformly apply until jar is empty over 6,000 sq. ft. Ortho® and Dial 'n Spray® are registered trademarks of OMS Investments, LLC		
		<b>USE RATE</b>
[AMDRO Spray Delivery System Name]		Apply as a spot treatment to wet the weeds to the point of runoff. [One 2.5 fl. oz. Image cartridge covers 625 sq. ft. of lawn.]



{Optional Hose-end Ready Spray}

Apply at the rate of 24 oz. Image to cover 6,000 sq. ft. of lawn

(1 quart Image to cover 8000 sq. ft.)

1. Make sure knob control lever is in the OFF position. Shake well and attach to hose.
2. Turn water on at faucet, aim nozzle toward lawn and turn control lever to ON to begin spraying. The product will automatically siphon and mix with water.

3. When finished, turn control lever to OFF position. Turn faucet off.

Discharge residual water pressure by turning control lever ON and OFF again.



(Alternate Hose-end Sprayer Directions for applicable packaging)

Apply at the rate of 24 oz. Image to cover 6,000 sq. ft. of lawn

(1 quart Image to cover 8000 sq. ft.)

1. Make sure the handle is in the upright position and locked **OFF** position. Shake container well and attach spray nozzle to hose.
2. Turn water on at faucet, aim nozzle toward lawn, bend tab back, and rotate handle clockwise towards the hose connection in the **ON** position to begin spraying. The product will automatically siphon and mix with water.
3. When finished turn the handle counterclockwise until the handle is in the upright position and locked position. Turn faucet off. Discharge residual water pressure by turning the handle ON and OFF again. Unscrew spray nozzle from hose.

(Alternate Hose-end Sprayer Directions for K2K sprayer packaging)

#### HOW TO APPLY

APPLY AT THE RATE OF 1 QUART OF THIS PRODUCT TO COVER 8,000 SQ.FT. OF LAWN.

1. Measure the entire area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over the entire measured area.

Amount of IMAGE® RTS HERBICIDE CONSUMER CONCENTRATE	Provides Coverage of Area Size:
[1 fl oz]	[250 ft <sup>2</sup> ]
[2 fl oz]	[500 ft <sup>2</sup> ]
4 fl oz	1000 ft <sup>2</sup>
[6 fl oz]	[1500 ft <sup>2</sup> ]
[8 fl oz]	[2000 ft <sup>2</sup> ]
[XX fl oz]	[XXXX ft <sup>2</sup> ]

{Note to reviewer: Values for less than or greater than 1000 ft<sup>2</sup> may be added to deleted, depending on package size}





The [AMDRO Spray Delivery System Name] is a patent pending innovative home lawn, landscape, and perimeter tank spray system. Spot treat weeds with this easy to use system.

1. Unscrew the tank cap and fill the tank with water up to the one gallon indicator. [The tank holds one gallon of water.]
2. Remove the [sprayer][wand][applicator] from the tank holster and unfold the [sprayer][wand][applicator].
3. Select the [cartridge][bottle] [pesticide] [herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
4. Remove the cap from the [cartridge][bottle][herbicide]. [Each [cartridge][bottle] contains 2.5 fl. oz. of [product][liquid].][Each 2.5 fl.oz. [cartridge][bottle] of concentrate makes one gallon of [liquid][ready to use] spray.]
5. Insert the [cartridge][bottle] [pesticide] [herbicide] into the top of the sprayer and twist until tight.
6. Press the trigger to spot spray [weeds] as listed on the product label.
7. When finished spot treating remove the [cartridge][bottle] [pesticide] [herbicide]. Replace the cap on the [cartridge][bottle] [pesticide] [herbicide]. Follow label Storage and Disposal instructions for product [pesticide][herbicide] [cartridge] [bottle].
8. After spot spraying rinse the [sprayer][wand][applicator] according to the RINSE: USE DIRECTIONS.

#### **RINSE: USE DIRECTIONS**

1. Remove the [cartridge][bottle][pesticide][herbicide]. Ensure that there is water in the one gallon tank.
2. Press and hold the [trigger][sprayer] for 10 seconds to flush the [sprayer][wand][applicator] with clean water from the tank.

#### **CHANGE [CARTRIDGE][BOTTLE]: USE DIRECTIONS**

1. Remove the [cartridge][bottle][herbicide] currently loaded in the [AMDRO Spray Delivery System Name].
2. Follow label Storage and Disposal instructions for product [pesticide] [herbicide] [cartridge] [bottle].
3. Press and hold the [trigger][sprayer] for 10 seconds to flush the [sprayer][wand][applicator] with clean water from the tank.
4. Select a new [cartridge][bottle][pesticide][herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
5. Insert the [cartridge][bottle][pesticide][herbicide] into the top of the sprayer and twist until tight.
6. Press the trigger to spot spray [weeds] as listed on the product label.
7. When finished spot treating remove the [cartridge][bottle][pesticide][herbicide]. Replace the cap on the [cartridge][bottle][pesticide][herbicide]. Follow label Storage and Disposal instructions for product [pesticide] [herbicide] [cartridge] [bottle].
8. After spot spraying, rinse the [sprayer][wand][applicator] according to the RINSE Use Directions. Store the [AMDRO Spray Delivery System Name] OR RINSE and change [cartridge][bottle].

**NOTE: IF OUTPUT SPRAY IS DIMINISHED OR SPUTTERING, DO THE FOLLOWING:**

1. Check that there is water remaining in the one gallon tank. If not, refill the tank with water.
2. Check that there is liquid remaining in the [cartridge][bottle]. If not, change to a new [cartridge][bottle].

[Two [cartridges][bottles] can be stored on the back of the [AMDRO Spray Delivery System Name]].

**[Optional:]CHANGING BATTERIES:**

If sprayer does not operate or spray performance declines significantly, change the four AA alkaline batteries in the sprayer handle as follows:

1. Remove the [cartridge][bottle].
2. Slide the battery cover open in the underside of sprayer handle.
3. Allow battery cover to swing open.
4. Remove old batteries.
5. Replace with four AA alkaline batteries. Make sure the positive and negative ends match the direction indicated on the inside of the battery door.
6. Close battery door and slide the door until latch is closed.


[AMDRO® Spray Delivery System Name] Helpful Hints

- Remove yellow tab and insure batteries are inserted correctly
- Fill the [AMDRO® Spray Delivery System Name] with one gallon of water prior to loading the cartridge
- Do not discard red caps-keep for storage
- Do not engage the trigger when the wand is folded
- Twist nozzle to switch from mist to stream
- Do not store [AMDRO® Spray Delivery System Name] in excessive heat or cold
- Remove [AMDRO® Spray Delivery System Name] cartridge. Spray water for 10 seconds before switching cartridge or storing
- Never put chemical into the tank. Only water can be put into the tank
- Read the entire label for the selected [AMDRO® Spray Delivery System Name] cartridge prior to use. Observe all precautions and follow directions carefully

**APPLICATION INSTRUCTIONS**


- Use only on well-established, non-stressed, labeled lawngrasses with a dense and uniform stand. TEMPORARY YELLOWING MAY OCCUR.
- Water in within 1-7 days after application to wash the IMAGE into the root zone where the greatest weed control activity occurs.
- For difficult to kill weeds, a second application can be made 6 weeks after the initial application.
- Applied as directed, this product provides 0.375 pound a.i. per acre, per application. Do not exceed 2 applications (0.75 pound a.i. per acre) per year.
- For best results, do not mow immediately before or for 48 hours after application.
- IMAGE may injure tall fescue. Do not use on tall fescue when it is used as a desirable turfgrass.
- Wait at least 1½ months after treatment before reseeding, winter overseeding, or plugging lawns.

- Not for use on newly planted or plugged lawns – nor on winter overseeded lawns, including ryegrass, when the overseeded grass is desirable.
- Avoid application to soil where annual bedding plants may be planted the following year, or plant injury may result.
- Try to avoid overlaps when spraying. Apply only when wind is no more than 10 mph.
- Avoid drift onto vegetables, flowers and unlabeled ornamental shrubs and plants as injury may result.
- Do not graze or feed treated grass clippings. Do not apply in vegetable gardens.
- Do not apply in greenhouses.

KILL [CONTROL] OF SUMMER WEEDS		
To kill	yellow nutsedge	purple nutsedge
	annual sedge <sup>†</sup>	green kyllinga
	rice flatsedge	globe sedge
	dollarweed <sup>†</sup>	violets <sup>†</sup>
	Virginia buttonweed <sup>†</sup>	field sandbur
	crabgrass <sup>†</sup>	bahiagrass <sup>†</sup>
	perennial ryegrass	tall fescue
	common chickweed	dandelion <sup>†</sup>
	white clover	

Do not apply during the cooler winter months when turf and weeds are not actively growing. Do not apply just prior to or during transition or during periods of very slow turf growth or severe discoloration could occur.

<sup>†</sup>More than one application may be needed for tough, hard-to-kill weeds.

KILL [CONTROL] OF WINTER WEEDS		
To kill	annual bluegrass <sup>†</sup>	hairy bittercress
	<i>(Poa annua)</i>	lawn burweed
	tall fescue	red sorrel
	common chickweed	wild onion
	cutleaf evening primrose	perennial ryegrass
	dovetail geranium	buttercup
	knawel	cudweed
	purple deadnettle	Carolina geranium
	wild garlic	henbit
	black medic	parsley-piert
	mouseear chickweed	white clover
	dandelion <sup>†</sup>	

Time applications to follow the first killing frost. IMAGE may be applied prior to or soon after emergence of the weeds listed with the exception of wild onion and wild garlic which must be emerged at application time for control. DO NOT USE ON ST. AUGUSTINEGRASS FOR WINTER WEED CONTROL or other labeled turfgrasses that do not go dormant.

<sup>†</sup>More than one application may be needed for tough, hard-to-kill weeds.

#### KILL (CONTROL) OF WEEDS IN ORNAMENTALS AND GROUND COVERS

IMAGE is effective in killing (controlling) the listed weeds in established landscape plantings. Apply IMAGE spray solution to the area where weeds occur. Applications must only be made over the top of or around the *listed* ornamentals. APPLICATION TO THE FOLIAGE OR ROOT ZONE OF UNLABELED ORNAMENTALS, PERENNIALS, OR ANNUAL BEDDING PLANTS MAY RESULT IN SEVERE PLANT INJURY. Use only on well-established, non-stressed plants. Ornamentals which are known to be severely injured by IMAGE include azalea, ligustrum (privet), viburnum, pieris, abelia and roses.

IMAGE CAN BE APPLIED TO THE FOLLOWING ORNAMENTALS:

Indian Hawthorne	<i>(Raphiolepis indica)</i>
Dwarf Yaupon Holly 'Schillings Dwarf'	<i>(Ilex vomitoria)</i>
Blue Pfitzer Juniper 'Pfitzeriana Glauca'	<i>(Juniperus chinensis)</i>
Helleri Holly 'Helleri'	<i>(Ilex crenata)</i>
Red Tip Photinia	<i>(Photinia X fraseri)</i>
Yucca	<i>(Yucca pendula)</i>
Wax Myrtle	<i>(Myrica cerifera)</i>
Burford Holly 'Burfordii'	<i>(Ilex cornuta)</i>
Gardenia 'Miami Supreme'	<i>(Gardenia jasminoides)</i>
Flame of the Woods	<i>(Ixora coccinea)</i>
Confederate Jasmine	<i>(Trachelospermum jasminoides)</i>
Parsons Juniper 'Parsoni'	<i>(Juniperus conferta)</i>
Blue Pacific Juniper 'Blue Pacific'	<i>(Juniperus conferta)</i>
Blue Rug Juniper 'Wiltonii'	<i>(Juniperus horizontalis)</i>
Red Crape Myrtle 'Red'	<i>(Lagerstroemia indica)</i>

Asiatic Jasmine  
'Minima' *(Trachelospermum asiaticum)*

---

Giant Liriope  
'Evergreen Giant' *(Liriope muscari)*

---

Variegated Liriope  
'Silvery Sunproof' *(Liriope muscari)*

---

Pachysandra *(Pachysandra terminalis)*

---

Mondograss *(Ophiopogon japonicus)*

---

Hosta  
'Lancifolia' *(Hosta sp.)*

---

Society Garlic *(Tulbaghia violacea)*

---

[In case of an emergency endangering life or property involving this product, call day or night, Area Code 800-265-0761.]

[For more information, please visit our website at [www.imageforweeds.com](http://www.imageforweeds.com).]

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call [xxx-xxx-xxxx] [the National Pesticide Information Center at 1-800-858-7378].

Image, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.



Manufactured for:

Ambrands  
1000 Parkwood Circle, Suite 700, Atlanta, GA 30339  
1-800-265-0761  
[gardendivision@central.com]

{Batch Code to be inserted at production.}

[Made in USA logo



{Small package, 2.5 fl oz cartridge label for single-cartridge delivery system use}

**IMAGE herbicide consumer concentrate**

NET CONTENTS 2.5 fl oz (73.93 ml)

**ACTIVE INGREDIENT:**

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-3-quinolinecarboxylic acid)\*..... 3.3%

**OTHER INGREDIENTS** ..... 96.7%

**TOTAL** ..... 100.0%

\*Equivalent to 3.14% 2-[4,5-dihydro-4-methyl-4(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-3-quinolinecarboxylic acid.(1 pint contains 0.0375 pounds of active ingredient as the free acid)

**Keep out of reach of children**

**CAUTION**

See accompanying box and insert for Additional Precautionary Statements, First Aid and Complete Directions for Use.

AMBRANDS, 1000 Parkwood Circle, Suite 700, Atlanta, GA 30339

EPA Reg. No. 73342-4

EPA Est. No. \_\_\_\_\_