8/14/2009

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1/2

AUG 1 4 2009

Richard W. Fresh, Director Regulatory Affairs Suterra LLC 213 SW Columbia St. Bend, OR 97702-1013

Subject: Puffer CM-O (EPA Reg. No 73479-9)

Label Amendment

Application Dated: May 27, 2009

Decision Number: 414700

Dear Mr. Fresh:

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

- 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- 2. Submit three (3) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(b). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

If you have any questions contact John Fournier at 703-308-0169 or by email at: fournier.john@epa.gov. A stamped copy of the label is enclosed for your records.

Sincerely,

Linda A. Hollis, Chief

Biochemical Pesticides Branch

CONCURRENCES icides and Pollution						
SYMBOL ▶ 75// F		Prevention Division	on (7511P)			
SURNAME Forum	sure OV					
DATE MANG	09					

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swellowed, inheled or absorbed through the skin. Causes moderate eye and skin initiation, Avoid breathing vepor, Avoid contact with eyes, skin or clothing. Whish thoroughly with seep and water after handling and before eating, dinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

IF INHALED: Move person to fresh air, If person is not breathing call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible, Call a Poison Control Center or doctor

If ON SKIN: Take off contaminated clothing, Rinse skin immediately with plenty of water for 15-20 minutes, Call a Poison Control Cente

IF IH EYES: Hold eye(s) open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and continue rinsing. Call a Poison Control Center or doctor for advice.

IF SWALLOWED: Call a Poleon Control Center or doctor immed stely for treatment advice. Heve person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Polson Control Center or doctor. Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment, You may also contact 1-866-326-6737 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment

PHYSICAL OR CHEMICAL HAZARDS

Plammeble, Contents under pressure, Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause burating.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal, PESTICIDE STORAGE: Store container in cool, dry area away from heat or open flame, PESTICIDE DISPOSAL: Westes resulting from the use of this product may be disposed of on site or at an approved disposal fecility.



CONTAINER DISPOSAL: Nonrefiltable container. Do not reuse or refill this container. Remove from cabinet, wrap empty container in newspaper and discard in trash or at an approved waste disposal facility. Do not puncture or incinerate. If can is empty offer for recycling.

WARRANTY

Substra, LLC warrants that this meteoral conforms to the chemical description on the lebel and is reseconably fit for the purposes stated in the Directions for Use. To the extent consistent with monitoritie law. Buterra neither makes nor authorizes any agent or representative to make, any approxima may always in networ makes nor autonoxima and gover of representatives to make, any other westanty of literase or of makes, any other westanty of literase nor incommon and other material. To the extent consistent with applicable law, Seterra's maximum fieldity for breach of this ventum prised into succeed the purchase price of this product. To the extent processistent with applicable law, layer and user schooledge and assume all risks and liabilities reaching from the handling, storage and use of this material contrary to label instructions.

CheckMate[®] Puffer CM-O



1 [

FOR ORGANIC PRODUCTION

Lasts up to 200 Days For the Control of Codling Moth in Apple. Pear and Walnut Orchards

CheckMate® Puffer CM-O contains a behavior modifying blochemical (pheromone) that disrupts the mating behavior of codling moth, Cydia pomonella. Use in apple, pear, walnut and other orchards where the codling moth is a pest. CheckMate Puffer CM-O is used with the CheckMate Puffer Aerosol Cabinet, an automatic metered dispenser, so that one puff of pheromone is delivered every 15 minutes throughout the night to disrupt the noclumal mating behavior of codling moths.

Active Ingredient*					
8,10 dodecadien-1-ol, (E,E)	18.05 %				
Other Ingredients	81.95 %				
Total	100:00 %				
*Containe 60.33 a.a.i. ner annieter:					

7.22 mg a.i. per 40 microliter puff KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back panel for First Aid and Precautionary States

Manufactured for:

puffCM-O#6 01/2009

Suterra°

Suterra LLC 213 SW Columbia St. Bend, Oregon U.S.A. 97702 Phone: (866) 326-6737 Fax: (541) 388-3705

EPA Registration No.: 73479-9 EPA Establishment No. 9444-LA-1 Net Contents: 13.5 oz (384 grams) Batch Code #: See bottom of can

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

> Do not apply more than 150 grams of active ingredient per acre per year.

METHOD OF APPLICATION: For use on apple, peer and walnut orchards and other crops where the codling moth is a pest. In the spring, start applications prior to moth emergence and continue throughout the crop's susceptible period; typically 80 to 180 days. Use a cabinet to automatically dispense the CheckMate Puffer CM-O canister. One puff of product is delivered every 15 minutes for a period of 12 hours, starting at 5 PM.

Plecing cabinets in orchards and replacing canisters are to be done when cabinets are inactive, typically during daylight hours between 5 AM and 5 PM: To avoid accidental spraying ensure that canister nozzles are pointed away during the performance of these tasks.

One Checkliste Puffer CM-O canister will last approximately 200 days. Where effective disruption of codling moth mating requires longer application periods, replace used canisters before the 200 day period is reached,

Product Piecement: Suspend a filled cabinet from a limb in the upper third of the tree canopy. Place cabinet where spray is directed away from follage; fruits or nuts, with 3-foot clearance, if possible

For best results, use in orchards greater than or equal to 40 acres. Use in orchards less than 40 acres will lead to less than optimal results unless properly managed. Use in combination with other effective meting disruption products around the entire orchard perimeter for codling moth when adjacent orchard(s) is not simultaneously treated with codling moth meting disruption.

Orchards Greater than or Equal to 40 Acres: For best results piece cabinets around the orchard perimeter or in a grid pattern to achieve a density of at least 1 puffer canister per acre; use up to 2 puffer canisters per acre in heavily infested orchards. In situations with a prevailing wind place additional canisters along the upwind edge to achieve uniform pheromone distribution within the interior of the orchard.

Orchards Smaller than 40 Acres; Improved effectiveness is achieved when adjacent orchard(s) is simultaneously treated with codling moth meting disruption, and the resulting total treated area is 40 acres or greater. Improved effectiveness is also achieved by placing puffer cantelers at a density of up to 2 CheckMate Puffer CM-O cantelers per acre. Place cabinels around the orchard perimeter or in a grid pattern, In situations with a prevailing wind place additional canieters along the upwind edge to achieve uniform pheromone distribution within the inte-

APPLICATION NOTES: For best results apply at the time or shortly before the codling moths begin to emerge in the spring. This product only affects adult male moths and will have no effect on female moths. eggs or larvae. If application occurs following biofix (the first date moths are found consistently in monitoring traps) or during the growing season, the field must be treated with insecticides that will effectively control hatching larvae until egg laying by previously mated female moths has coased to occur, in moderate to high insect pressure situations, supplemental insecticide applications during the season may be necessary to provide adequate protection. These supplemental insecticide applications must be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control young larvae. Monitor insect infestation with traps and by visual inspection of plants. Re-apply as needed, based on monitoring results and field scouting.

Rodenticide pesticide regi Ę