SEP 3 0 2008

Mr. Richard Fresh Suterra, LLC. 213 SW Columbia Street Bend, OR 97702

Subject: Puffer OFM-O

EPA Registration No. 73479-10

Label Amendment to update container disposal per PRN 2007-4, addition of steel

recycle symbol, and add batch code number.

Application Dated: June 16, 2008

Dear Mr. Fresh:

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(5), is acceptable provided that you:

- 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- 2. Submit five (5) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. If you have any questions contact Andrew Bryceland at 703-305-6928 or by e-mail at: bryceland.andrew@epa.gov. A stamped copy of the label is enclosed for your records.

Sincerely,

Linda A. Hollis, Chief

Biochemical Pesticides Branch

Biopesticides and Pollution

Prevention Division (7511P)

Enclosure			CONCURRENCES					
SYMBOL >	5118				,			
SURNAME O	bryceland							
DATE 0	1/4/08							

EPA Form 1320-1A'(1/90)

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

Caution: May be harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye and skin imitation. Avoid contact with eyes, skin or clothing. Avoid breathing vapor. Wash thorough's with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

#### FIRST AID STATEMENT

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vorniting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes, Call a Poison Control Center or doctor for advice.

IF IN EYES: Hold eye(s) open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and continue rinsing. Call a Poison Control Center or doctor for advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-866-326-6737 for emergency medical treatment information.

### **ENVIRONMENTAL HAZARDS**

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or finsate.

### PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. PESTICIDE STORAGE: Store container in cool, dry area away from heat or open flame. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.



CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Remove from cabinet, wrap empty container in newspaper and discard in trash or at an approved waste disposal facility. Do not puncture or incinerate. If can is empty offer for recycling.

### WARRANTY

Suterra, LLC warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use. To the extent consistent with applicable law, Suterra neither makes nor authorizes any agent or representative to make, any other warranty of times or of merchantability, guarantee or representation, express or implied, concerning this material. To the extent consistent with applicable law, Suterra's maximum fability for breach of this warranty shall not exceed the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and labilities resulting from the handling, storage and use of this material contrary to tabel instructions.

# CheckMate® Puffer OFM-O

FOR ORGANIC PRODUCTION

Lasts up to 200 Days
For the Control of Oriental Fruit Moth in
Almond, Apple, Apricot, Cherry, Nectarine,
Peach, Pear, Plum, Prune and Quince Orchards

CheckMate® Puffer OFM-O is an aerosol canister containing behavior modifying biochemicals (pheromones) that disrupt the mating behavior of oriental fruit moth, *Grapholita molesta*, it is intended for use in almond, apple, apricot, cherry, nectarine, peach, pear, plum, prune, and quince orchards and other areas where the oriental fruit moth is a pest.

CheckMate Puffer OFM-O is designed for use with the CheckMate Puffer Aerosol Cabinet, an automatic metered dispenser, so that one puff of pheromone is delivered every 15 minutes throughout the night to disrupt the noctumal mating behavior of oriental fruit moths.

### Active Ingredients\*

(Z)-8-Dodecenyl Acetate	11.63	%
(E)-8-Dodecenyl Acetate	. 0.75	%
(Z)-8-Dodecenol	. 0.12	%
Other Ingredients	87,50	%
Total	00.00	%

\*Contains 48.004 g a.i. per canister; 4.81 mg a.i. per 40 microliter puff

### C Δ I TION

See side/back panel for First Aid and Precautionary Statements
Manufactured for:

# Suterra®

Suterra LLC 213 SW Columbia St. Bend, Oregon U.S.A. 97702 Phone: (866) 326-6737 Fax: (541) 388-3705

EPA Registration No.: 73479-10 EPA Establishment No. 9444-LA-1 Net Contents: 13.5 oz. (384 grams) Ratch Code #: See bottom of can.

puffOFM-O#5 08/2008

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

Do not apply more than 150 grams of active ingredient per acre per year.

METHOD OF APPLICATION: For use in almond, apple, apricot, cherry, nectarine, peach, pear, plum, prune and quince orchards and other areas where the oriental fruit moth is a pest. In the spring, start applications prior to moth emergence and continue throughout the crop's susceptible period, typically 80 to 180 days. Use a cabinet to automatically dispense the CheckMate Puffer OFM-O canister. One puff of product is delivered every 15 minutes for a period of 12 hours, starting at 5 PM.

Placing cabinets in orchards and replacing canisters are to be done when cabinets are inactive, typically during daylight hours between SAM and 5 PM. To avoid accidental spraying, ensure that canister nozzles are pointed away during the performance of these tasks.

One CheckMate Puffer OFM-O canister will last approximately 200 days. Where effective disruption of oriental fruit moth mating requires longer application periods, replace used canisters before the 200 day period is reached.

<u>Product Placement</u>; Suspend a filled cabinet from a free limb close to the center of the tree. Ideally, placement height will be approximately 2/3 of orchard height ±10%. Do not place cabinets where foliage, nuls or fruits can be directly sprayed. Where feasible a 3-foot clearance is requested.

Orthards Smaller than 40 Acres; Do not use CheckMate Puffer OFM-O in orchards less than 40 acres unless adjacent acreage is also being treated with CheckMate Puffer OFM-O and total acreage block being treated is 40 acres or more.

40-Acre Orchard: Place cabinets around the orchard perimeter or in a grid pattern to achieve a density of two (2) puffer canisters per acre. In situations with a prevailing wind place additional canisters along the upwind edge to achieve uniform pheromone distribution within the interior of the orchard.

Orchards Greater than 40 Acres: Place cabinets around the orchard perimeter or in a grid pattern to achieve a density of up to two (2) puffer canisters per acre. In situations with a prevailing wind place additional canisters along the upwind edge to achieve uniform pheromone distribution within the interior of the orchard.

APPLICATION NOTES: For best results apply at the time or shortly before the oriental fruit moths begin to emerge in the spring. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix (the first date moths are found consistently in monitoring traps) or during the growing season, the field must be treated with insecticides that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. In moderate to high insect pressure situations, supplemental insecticide applications must be made based on trap monitoring, field scouting, and appropriate degree-day models for the local growing area and must be timed to control young larvae. Monitor insect infestation with traps and by visual inspection of plants. Reapply, as needed, based on monitoring results and field scouting.

## **ACCEPTED**

SEP 3 0 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 73479-10

E CC