

73600-3

6/4/2010

1098



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

June 4, 2010

Mr. Chuck Koopmann
Ridley Us Holdings, Inc.
P.O. Box 8500
Mankato, MN 56002

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Mr. Koopmann:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated April 29, 2010 for the following product(s):

✓ **Sweetlix 16:8 Free Choice Mineral with Rabon Oral Larvicide**
EPA Reg. No. 73600-3

VMS Rabon 7.76 Oral Larvicide Premis
EPA Reg. No. 73600-4

VMS Pest-A-Side Block with Rabon
EPA Reg. No. 73600-5

VMS EnProAI Fly Control Block with Rabon Oral Larvicide
EPA Reg. No. 73600-6

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact Regina Foushee'-Smith at 703-605-0780.

2098

Sincerely,

A handwritten signature in black ink, appearing to read "Richard Gebken". The signature is written in a cursive style with a long horizontal stroke at the end.

Richard J. Gebken
Product Manager
Insecticide Branch
Registration Division (7505P)

JUN 4 - 2010

(optional text according to species targeted)

Sweetlix 16:8 Free Choice Mineral with Rabon Oral Larvicide

For control of fecal flies in manure of treated cattle and horses. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle. Prevents development of house flies and stable flies in the manure of treated horses.

A free choice supplement block for cattle and/or horses.

ACTIVE INGREDIENT:

Tetrachlorvinphos: (Z)-2-Chloro-1-(2,4,5-trichlorophenyl) vinyl demethyl phosphate*	0.494%
Other ingredients**	99.506%

*Each lb. contains 2.24 g Rabon®

**Refers only to ingredients which are not larvicidal

GUARANTEED ANALYSIS

Will vary according to species requirements, geographic area and regional feeding requirements.

FEED INGREDIENTS**

Molasses products, roughage products, grain products, plant protein products, processed grain by-products, mineral products, vitamins, food additives permitted in animal feed/drinking water, GRAS ingredients, fats/oils, animal products, animal protein products***, miscellaneous products***, chemical preservatives, amino acids & related products, fermentation products, distillers products, enzymes, brewers products, marine products, special purpose products

** AAFCO approved feed ingredients may vary depending upon nutritional requirements in different species and geographic regions of the country.

***Contains no prohibited animal-by-products in accordance with FDA regulation 21 CFR 589.2000.


EPA No: 73600-3

EPA Est No: 73600-XX-01 or 74604-KY-001

Net weight: 25 to 50 lbs.

KEEP OUT OF REACH OF CHILDREN

CAUTION



6048

FIRST AID	
IF SWALLOWED:	-Call a poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to by a poison control center or doctor.
IF ON SKIN:	-Take off contaminated clothing. -Rinse skin with plenty of water for 15-20 minutes. -Call poison control center or doctor for treatment advice.
IF IN EYES:	-Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. -Call a poison control center or physician for treatment advice.
HOT LINE NUMBER	
In case of emergency, contact 507-388-9400.	
NOTE TO PHYSICIANS AND VETERINARIANS	
Symptomalogy: Signs of organophosphate toxicity are those of cholinergic over stimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes 3 categories; musarinic blocking agents (atropine sulfata), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry).	

PRECAUTIONALY STATEMENTS

CAUTION

Harmful is swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Not to be used on horses destined for slaughter.

Precautions For Horse Use: This product contains tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, consult your veterinarian immediately. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses

(Product Name) should be fed continuously on a free choice basis to cattle and/or horses to provide minerals, vitamins and Rabon® daily. Feed as the only FREE CHOICE source of salt, other minerals or vitamins. Rabon oral larvicide should be fed at a level to provide 70 mg Rabon per 100 lb. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). Feed supplies (pasture or other forages) and feeder locations should be adequate to insure proper mineral consumption. Consumption in accordance with intake levels listed below will deliver 70 mg Rabon per 100 lb bodyweight per day:

ANIMAL WEIGHT (lbs)	SUPPLEMENT DAILY INTAKE		DAILY INTAKE OF RABON (mg)
	(oz)	(lbs)	
500	2.5	0.15625	350
600	3.0	0.18750	420
700	3.5	0.21875	490
800	4.0	0.25000	560
900	4.5	0.28125	630
1000	5.0	0.31250	700
1100	5.5	0.34375	770
1200	6.0	0.37500	840

If animals consume more or less than these levels, feeders should be moved further from or closer to the general resting or watering areas. If animals still fail to consume the correct amounts, discontinue feeding this free choice supplement.

Provide fresh, clean water to all animals at all times.

Follow feeding directions. Consumption of selenium should not exceed 3 mg per head daily.

Fly Control Program

(Product Name) prevents the development of fecal flies in the manure of treated cattle and horses, but it is not effective against existing adult flies. It may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and

8098

stable flies which can breed not only in animal manure but also in other decaying matter on the premises. In order to achieve optimum fly control, (Product Name) should be used in conjunction with other good management and sanitation practices. Start feeding (Product Name) early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. (Product Name) may be safely fed to cattle up to slaughter and to pregnant and lactating mares.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of the smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

For additional information, please contact your Sweetlix representative.

Rabon[®] is a registered trademark of KMG Chemicals, Houston, TX.

Manufactured by
Ridley US Holdings, Inc.
P.O. Box 8500
Mankato, MN 56002

Contact us at 1-87SWEETLIX
www.sweetlix.com

