

73600-6

6/24/2003

1/2

VMS

UNSTOCKING VITAMIN SUPPLEMENT

ENPROAL[®] FLY CONTROL BLOCK with RABON[®] Oral Larvicide A Free Choice Block for Cattle and Horses

ACCEPTED
JUN 24 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide Registered under EPA Reg. No. 73600-6

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle and to prevent the development of house flies and stable flies in the manure of treated horses.

ACTIVE INGREDIENTS:	BY WEIGHT:
Tetrachlorvinphos: (Z) 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate*	0.30%
OTHER INGREDIENTS**	99.70%
TOTAL	100.00%

*Contains 1.35g of Rabon per pound
**Refers only to ingredients which are not larvicidal

KEEP OUT OF REACH OF CHILDREN
CAUTION

See reverse for additional precautionary statements and for feeding directions.

GUARANTEED ANALYSIS

Crude Protein (Min)	7.50%
Crude Fat (Min)	0.20%
Crude Fiber (Max)	3.00%
Calcium (Min)	2.00%
Calcium (Max)	2.50%
Phosphorus (Min)	1.00%
Salt (Min)	5.00%
Salt (Max)	6.00%
Magnesium (Min)	3.00%
Potassium (Min)	2.40%
Copper (Min)	400 ppm
Selenium	12 ppm
Zinc (Min)	1,750 ppm
Vitamin A (Min)	58,000 IU/lb.
Vitamin D-3 (Min)	10,000 IU/lb.
Vitamin E (Min)	25 IU/lb.

FEED INGREDIENTS

Cane Molasses, Cottonseed Meal, Magnesium Oxide, Salt, Defluorinated Phosphate, Magnesium Sulfate, Potassium Sulfate, Zinc Oxide, Iron Carbonate, Manganous Oxide, Sodium Selenite, Calcium Carbonate, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

EPA No: 73600-6
ESP Est No: 73600-AL-01

Manufactured By:
Sweetlix LLC
Salt Lake City, UT 84111

NET WEIGHT 50 LB.
(22.7 KG)

41530

05/03

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give by mouth and do not induce vomiting.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention.
IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIANS AND VETERINARIANS

Symptomatology: Signs of organophosphate toxicity are those of cholinergic over stimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes 3 categories; musannic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxine chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry). In case of emergency, contact Sweetlix LLC (800) 325-1486.

PRECAUTIONARY STATEMENTS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals. Personal Protective Equipment (PPE): long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This compound is toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Not to be used on horses destined for slaughter.

1. Feed 1 block per 5 head of cattle or horses. This allows all animals equal access to blocks. Feed blocks at the rate of 0.88 oz. per 100 lb. of body weight per day. This intake will supply 0.07g. of larvicide per 100 lb. of body weight per day (2.1 grams per 100 lbs. body weight per month).
2. Place blocks where animals congregate. Locate blocks through-out pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Cattle will consume about 8.8 oz. per head daily. When animals consume the block at the recommended rate, it will supply 3 mg. of supplemental selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE CHOICE source of salt, other minerals and vitamins.
7. VMS ENPROAL Fly Control Block with Rabon® Oral Larvicide prevents the development of fecal flies in the manure of treated cattle and horses, but it is not effective against existing adult flies. It may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, VMS ENPROAL Fly Control Block with Rabon Oral Larvicide should be used in conjunction with other good management and sanitation practices. Start feeding VMS ENPROAL Fly Control Block with Rabon Oral Larvicide early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. VMS ENPROAL Fly Control Block with Rabon Oral Larvicide may be fed to cattle up to slaughter and to lactating cows without withholding the milk from market either during or after treatment. VMS ENPROAL Fly Control Block with Rabon Oral Larvicide may be safely fed to pregnant and lactating mares.
8. Follow feeding directions. Consumption of selenium should not exceed 3 mg per head per day.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

For additional information, please contact your Sweetlix representative.

Rabon® is a registered trademark of KMG Chemicals, Houston, TX