

74152-4

01/12/2005

1/3

MegaGro
GROWTH STIMULATOR
CONCENTRATE
FOR MAXIMUM GROWTH OF
INDOOR AND OUTDOOR
PLANTS, VEGETABLES, FLOWERS, AND FRUITS
IN AND AROUND THE HOME

ACTIVE INGREDIENT:

Gibberellic Acid 0.025%

OTHER INGREDIENTS: **99.975%**

TOTAL **100.000%**

GUARANTEED ANALYSIS (Fertilizer)

Sulfur (S) combined..... 15.00%

Derived from Liquid Ammonium Thiosulfate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR FIRST AID AND PRECAUTIONARY STATEMENTS.

NET CONTENTS: 8 Fl. Oz./237 mL

NET WEIGHT: 0.625 LBS/0.28 KG

MANUFACTURED IN U.S.A FOR:

W.A. Industries, Inc
700 W. Van Buren St.
Chicago, IL 60607
EPA EST. NO. 081148-IL
EPA REG. NO. 74152-U

ACCEPTED

JAN 12 2005

**Under the Federal Insecticides,
Fungicides, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 74152-4**

READ ENTIRE LABEL BEFORE USE

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing dust or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

**IF ON SKIN
OR
CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF
SWALLOWED**

- Call a poison control center or doctor immediately for further treatment advice..
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents) call the National Pesticide Information Center at 1-800-858-7378.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purposes stated on the label when used in accordance with the directions concerning this product. There are no other warranties expressed or implied, including a warranty of fitness for a particular purpose. Because the time, place, rate of application and other conditions of use are beyond the seller's control, seller's liability from handling, storage, and use of this product is limited to replacement of product or refund of purchase price.

DIRECTION FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE: Spray as a fine mist at 14 day intervals during growing season. Spray until solution runs from leaf surface. 5 capfuls = 1 oz.

OUTDOOR/INDOOR FLOWERING SHRUBS, ORNAMENTAL PLANTS: Apply at 2 capfuls per 32 oz of water at 14-day intervals during the growing season. (DO NOT USE ON FERNS)

LEAFY VEGETABLES, MELONS, PUMPKINS, SQUASH, TOMATOES, PEPPERS, CABBAGE, BROCCOLI, CAULIFLOWER, STRAWBERRIES: Apply at 1 capful per gallon (128 oz) of water at 14-day intervals during the growing season.. Spray until solution runs from leaf surface. For peppers, tomatoes and strawberries apply 7 days after planting. For leafy vegetables, cabbage, broccoli, and cauliflower apply at 2-3 inch stage. For melons, pumpkins, squash apply at bunching.

CITRUS: Apply at 1 capful per gallon (128 oz) of water at 14-day intervals during the growing season. Cease application 2 weeks prior to harvest.

GRAPES, BLUEBERRIES: Apply at 3 capfuls per gallon (128 oz) of water, as fruit begins to form.

COMPATIBILITY: Product is generally compatible with pesticides and fertilizers used commonly in vegetable and fruit growing. Do not mix this product with lime or other highly alkaline materials. If mixing this product with other products, this product should always be the last ingredient added to the mixture.

STORAGE AND DISPOSAL: Store in original container away from extremes of temperature, excessive moisture and humidity. Avoid storage near oxidizers, strong acids or strong alkalies. Do not use copper, zinc, tin, or their alloys in storage or handling systems for this product. Do not reuse bottle. Rinse thoroughly before discarding in trash.