



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 6, 2025

Matthew Granahan
mgranahan@helmagro.com
HELM AGRO US, INC.

Subject: Label Amendment – Add a description of the action here
Registration Review Label Amendment – Incorporating Mitigation Measures
from the Registration Review Interim Decision for cycloate
Product Name: RO-NEET HERBICIDE
EPA Registration Number: 74530-16
EPA Receipt Date: 08/19/2021
Action Case Number: 00472614

Dear Matthew Granahan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The Agency, in accordance with FIFRA, as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the cycloate Interim Decision. The Agency has concluded that your submission is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. The next label printing of this product must use this labeling unless subsequent changes have been approved. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Lydia Crawford at 202-566-2575 or at crawford.lydia@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Lydia Crawford". The signature is fluid and cursive, with the first and last names being more prominent.

Lydia Crawford, Biologist
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



CYCLOATE

GROUP

8

HERBICIDE

Ro-Neet[®] Herbicide

[ABNs: Rovic & Rovic Herbicide]

Liquid Herbicide for Weed Control in Sugar Beets, Garden Beets, and Spinach

ACTIVE INGREDIENT:

Cycloate (S-ethyl cyclohexylethylthiocarbamate)	73.9%
OTHER INGREDIENTS	26.1%
TOTAL	100.0%

Contains petroleum distillate.

Contains 6 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

**SEE [BACK PANEL] [NEXT PAGE] [INSIDE LABEL] [INSIDE BOOKLET] [BELOW] FOR ADDITIONAL
PRECAUTIONARY STATEMENTS [STORAGE AND DISPOSAL] [AND] [DIRECTIONS FOR USE]**



EPA Reg. No. 74530-16

EPA Est No. _____

NET CONTENTS _____ [Gals.] [Gallons] [(L)] [(Liters)]

Manufactured for:
HELM Agro US, Inc.
401 E. Jackson St., Suite 1600
Tampa, FL 33602

ACCEPTED**2/6/2025**

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No.

74530-16

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN	
This product contains petroleum distillate. May pose an aspiration pneumonia hazard.	
FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC: 1-800-424-9300. For 24-Hour Medical Emergency Assistance call National Poison Control Center at 1-800-222-1222.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

CAUTION: Contains petroleum distillate. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Engineering controls are required for all mixers, loaders and applicators (see Engineering Controls for additional requirements).

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made Barrier Laminate, Nitrile Rubber ≥ 14 mils, Neo-prene Rubber ≥ 14 mils, or Viton ≥ 14 mils,
- Chemical-resistant apron, and.
- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Handlers performing tasks, such as spill clean-up or cleaning equipment, for which engineering controls are not feasible must wear:

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of Barrier Laminate, Nitrile Rubber ≥ 14 mils, Neo-prene Rubber ≥ 14 mils, or Viton ≥ 14 mils,
- Chemical-resistant apron, and
- Wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Engineering Controls

Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for dermal protection, and must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders,
- Wear protective eye wear, if the system operates under pressure,
- Either use a closed system that also meets the requirements in the WPS for inhalation protection or wear the type of respirator specified in the personal protective equipment sections of the labeling, and
- Be provided and have immediately available for use in an emergency, such as a spill or equipment breakdown: chemical-resistant footwear, and, if using a closed system cab that provides respiratory protection, a respirator of the type specified in the PPE section of this labeling.

Applicators must use motorized ground equipment that is equipped with an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, applicators must:

- Wear the personal protective equipment required in the PPE section of this labeling for applicators,
- Either wear the type of respirator specified in the PPE section of this labeling or use an enclosed cab with a properly functioning ventilation system that is used and maintained according to the manufacturer's written operating instructions and that is declared in writing by the manufacturer or by a government agency to provide at least as much respiratory protection as the type of respirator specified in the PPE section of this labeling.
- Be provided and have immediately available for use in an emergency for when they must exit the cab in the treated area: chemical-resistant gloves, and, if using an enclosed cab that provides respiratory protection, a respirator of the type specified in the PPE section of this labeling.
- Take off any PPE that was worn in the treated area before reentering the cab, and
- Store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

Additional Use Limitations for California Only: Mixers, loaders, applicators, and other handlers are prohibited from handling more than 93 gallons (560 pounds active ingredient) in any 21- day period. Property operators must include in their Pesticide Use Records the name of the person(s) that handled the product for each application.

User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Do not contaminate water when disposing of equipment wash water or rinsate.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Chemigation is prohibited. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Entry (including early entry that would otherwise be permitted under the WPS) by any person - other than a correctly trained and equipped handler who is performing a handling task permitted by the WPS - is PROHIBITED for 48 hours following applications. Do not allow entry by unprotected persons into the treated area until 48-hour period is expired and the warning signs are removed.

NOTIFICATION: Before the start of the application, notify workers of the application by warning them orally and by posting warning signs at entrances to the treated area. The signs must bear the skull and crossbones symbol and state: (1) “**DANGER/PELIGRO**”, (2) “**DO NOT ENTER/NO ENTRE**”, (3) the date and time of treatment, (4) “Ro-Neet Herbicide in use”, and (5) name, address, and telephone number of the applicator. Post the warning sign instead of the WPS sign for this application, but follow all WPS requirements pertaining to location, legibility, size, and timing of posting and removal.

Special Early Entry Exception: Workers may enter the treated area during the first 48 hours following application to plant crops, provided all of the following conditions are met:

(1) The special early entry workers must use motorized ground equipment that is equipped with an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5) for dermal protection. In addition, these workers must:

- Wear the personal protective equipment required in the PPE section of this labeling for applicators,
- *Either* wear the type of respirator specified in the PPE section of this labeling or use an enclosed cab with a properly functioning ventilation system that is used and maintained according to the manufacturer's written operating instructions and that is declared in writing by the manufacturer or by a government agency to provide at least as much respiratory protection as the type of respirator specified in the PPE section of this labeling.
- Be provided and have immediately available for use in an emergency when they must exit the cab in the treated area: chemical-resistant gloves, and, if using an enclosed cab that provides respiratory protection, a respirator of the type specified in the PPE section of this labeling,
- Take off any PPE that was worn in the treated area before reentering the cab, and
- Store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

(2) Employers of the special early entry workers must make sure that these workers:

- Have received training equivalent to WPS pesticide handler training before entering the treated area,
- Are provided with the specified type of respirator, and the respirator fits them correctly and is maintained as required in the WPS for handlers,
- Are provided all of the WPS protections for early entry workers, including PPE instructions, labeling information and instructions, decontamination sites, and duties related to providing, cleaning, and maintaining the PPE for early-entry workers.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep containers closed when not in use. Do not store near seeds, fertilizers, or foodstuffs. Protect from temperatures below 20° F. Product crystallizes at lower temperatures. Warm or store at higher temperatures and mix to re-dissolve crystals and assure uniformity before use. Do not store near sources of heat or open flame. **COMBUSTIBLE LIQUID.**

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable Container: Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

(Liquid Formulations) Nonrefillable container equal to or less than 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable container 5 gallons to bulk: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

RESISTANCE MANAGEMENT

For resistance management, Ro-Neet Herbicide is a Group 8 herbicide. Any weed population may contain or develop plants naturally resistant to Ro-Neet Herbicide and other Group 8 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Ro-Neet Herbicide or other Group 8 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving

- between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local HELM Agro US, Inc. representative.

PRODUCT INFORMATION

READ ALL LABEL DIRECTIONS BEFORE USING.

Ro-Neet Herbicide is a selective herbicide which is mixed (incorporated) into the soil, or applied below the surface of the soil for control of weeds listed on this label. Ro-Neet Herbicide controls weeds by interfering with normal seed germination and seedling development. It does not control established or germinated weeds present at application. All weed growth and crop stubble should be thoroughly worked into the soil before treatment.

PRODUCT USE PRECAUTIONS

Ro-Neet Herbicide must be used as directed at specified rates (do not overdose). Applied according to directions and under normal growing conditions, Ro-Neet Herbicide will not harm the treated crop. During germination and early growth, extended periods of unusually cold and wet or hot and dry weather, insect, nematode, or plant disease attack, carryover pesticide residues, the use of certain soil applied systemic insecticides, highly saline or alkaline soil conditions, improperly placed fertilizers or soil insecticides may create abnormal conditions that weaken crop seedlings. Ro-Neet Herbicide used under these abnormal conditions could result in crop injury.

Ro-Neet Herbicide may cause crop injury on very light sandy soil.

~~Tank mix this product with fungicides, insecticides, or herbicides only as recommended.~~

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

WEEDS CONTROLLED	
ANNUAL GRASSES	
Annual bluegrass	<i>Poa annua</i>
Annual ryegrass	<i>Lolium multiflorum</i>
Barley, volunteer	<i>Hordeum</i> spp.
Barnyardgrass (watergrass)	<i>Echinochloa</i> spp.
Crabgrass	<i>Digitaria</i> spp.
Foxtails	<i>Setaria</i> spp.
Wild oats	<i>Avena fatua</i>
ANNUAL BROADLEAF WEEDS	
Black nightshade	<i>Solanum nigrum</i>
Hairy nightshade	<i>Solanum villosum</i>
Henbit, deadnettle	<i>Lamium</i> spp.
Lambsquarters	<i>Chenopodium album</i>
Nettleleaf goosefoot*	<i>Chenopodium murale</i>
Purslane, common	<i>Portulaca oleracea</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Small stinging nettle, Burning nettle	<i>Urtica urens</i>
Velvetleaf**	<i>Abutilon theophrasti</i>

* California excluded.

** Partial control or suppression only.

The above broadleaf weeds will be controlled only if application is made when conditions are favorable for germination. Ro-Neet Herbicide WILL NOT CONTROL ESTABLISHED OR GERMINATED WEEDS PRESENT AT APPLICATION.

PERENNIAL WEEDS	
Yellow nutsedge	<i>Cyperus esculentus</i>
Purple nutsedge	<i>Cyperus rotundus</i>

Existing stands of nutsedge must be turned under and chopped thoroughly before treatment. Prior to planting, Ro-Neet Herbicide must be thoroughly mixed into the soil or applied subsurface to the soil with injectors for nutsedge control.

SOIL PREPARATION

Prepare the soil for seeding according to good agricultural practice. All weed growth should be thoroughly worked into the soil before treatment. DO NOT APPLY BEFORE PREIRRIGATION.

APPLICATION DIRECTIONS

Mechanical Soil Incorporation or Injection Applications: Apply Ro-Neet Herbicide to well-worked soil which is dry enough to permit thorough mixing with incorporation equipment or proper injection. Use standard low-pressure (20-50 psi) boom spray equipment. Application equipment should be carefully calibrated before use and checked frequently during application to be sure nozzles are free from clogging and delivering a uniform pattern. Apply UNIFORMLY the directed rate of Ro-Neet Herbicide in 10 to 50 gallons of water per acre. Avoid overlaps that will increase Ro-Neet Herbicide dosage above specified limits because plant injury will occur.

Sprinkler Irrigation: Begin sprinkler incorporation immediately following application, and complete within a 36-hour period using sufficient water to penetrate to a depth of 3-4 inches. The application and incorporation must be done within five days after the last tillage operation, since poor results will occur if weeds have germinated.

Ro-Neet Herbicide with Fluid Fertilizer: Ro-Neet Herbicide may be combined with fluid (solution, slurry, or suspension) fertilizers. However, physical compatibility with these fluid fertilizers must be determined before combining in the spray tank. See Appendix I giving directions for these combinations. Even though found to be compatible, constant agitation is necessary to keep the Ro-Neet Herbicide evenly mixed with the fluid fertilizer.

INCORPORATION DIRECTIONS

Ro-Neet Herbicide must be incorporated into the soil to prevent loss of the herbicide.

Timing: Ro-Neet Herbicide applications in water must be incorporated immediately (within minutes). Whenever possible, application and incorporation should be done in the same operation. Applications made in liquid fertilizer may have a maximum delay of 4 hours before incorporation. Applications made on dry bulk fertilizer must be incorporated within the same day of application. Immediate incorporation must be made with liquid and dry bulk fertilizer applications if soil surface is damp or if windy conditions (15 mph or greater) or rainfall are present or expected.

Depth: Ro-Neet Herbicide should be incorporated into the soil to a depth of 2-3 inches (**California only:** incorporate to a depth of 3 inches). Thorough soil mixing is necessary for good weed control. Use equipment which has been proven to incorporate thoroughly to the specified depth.

SOIL MIXING (INCORPORATION) BEFORE PLANTING

The following equipment commonly is used for soil mixing (incorporation) before planting:

For Broadcast (Overall) Application:

- Power Driven Cultivation Equipment (recommended on all soil types). Set to cut to a depth of 2 to 3 inches.
- Tandem Discs (recommended on all soil types). Set to cut to a depth of 4 to 6 inches, operated at 4 to 6 mph followed by a spiked-tooth harrow or some other leveling device which extends beyond the ends of the discs. For more thorough mixing (for nutsedge, heavier soils), disc in two different directions (cross disc).
- Field Cultivators (recommended on lighter soils in good tilth only). Do not use in California. Use 3 to 4 rows of sweeps, spaced at 7 inch or less intervals and staggered so that no soil is left unturned, followed by a spiked-tooth harrow or some other leveling device pulled behind the cultivator. Do not use chisels or points. Set the cultivator to cut 4 inches deep, operated at 5 mph or more. Run the equipment over the field twice, the second run at an angle to the first.
- Rotary Ground Driven or Spring-Tooth Cultivators (recommended on lighter soils in good tilth only). Do not use in California. Set to penetrate to a depth of 4 to 6 inches and operated at 5 to 8 mph in two different directions.

For Band (Row) Application:

Uniformly mix to a depth of 2 to 3 inches.

- Hooded power-driven rotary tillers.
- Hooded ground-driven rotary tillers.

Rolling: Under most conditions, the treated area should be firmly rolled between the incorporation unit and the planters.

PLANTING

For maximum weed control benefits, crops should be planted or seeded immediately after application except for fall applications. Do not use a drag behind the planter as it may concentrate Ro-Neet Herbicide over the seed row and cause crop injury.

CULTURAL PRACTICES FOLLOWING APPLICATION

Ro-Neet Herbicide is not persistent in the soil and susceptible weeds germinating later during the growing season may not be controlled. Shallow cultivation and approved postemergence herbicides may be necessary to control those susceptible weeds that escape control by Ro-Neet Herbicide as well as those weeds not susceptible to Ro-Neet Herbicide. Do not cultivate deeper than the depth of the herbicide incorporation.

When cultivating fields where Ro-Neet Herbicide has been banded in the row, use shields to prevent the movement of untreated soil into the treated row.

SUGAR BEETS, GARDEN BEETS, SPINACH USE RATES Broadcast Treatment			
CROP	APPLICATION TIMING	GALLONS Ro-Neet Herbicide PER ACRE	DIRECTIONS
Sugar Beets*	Preplant or At Planting or Immediately Post Planting	1/2 to 2/3	RESTRICTION: Use on mineral soils ONLY. Pre-Harvest Interval (PHI): Do not harvest sugar beets within 120 days after application. PRECAUTIONS: Use lower rate on sandy soils. Use higher dosage rate on heavier soil. Injury may result in highly saline or alkaline soils.
	Fall application	2/3	RESTRICTION: Apply only in the states of Idaho, Minnesota, Montana, North Dakota, Oregon, and Wyoming. Pre-Harvest Interval (PHI): Do not harvest sugar beets within 120 days after application. PRECAUTION: Apply and incorporate in late fall before the ground freezes.

Garden Beets	Preplant or At Planting or Immediately Post Planting	1/2 to 2/3	RESTRICTIONS: Use on mineral soils ONLY. Ro-Neet Herbicide in combination with fluid fertilizer, do not apply over 150 lbs. actual nitrogen per acre. Pre-Harvest Interval (PHI): Do not harvest garden beets within 45 days after application. PRECAUTIONS: Use lower rate on sandy soils. Use higher dosage rate on heavier soils.
Spinach	Preplant or At Planting or Immediately Post Planting	1/3 to 1/2 ¹ 1/2 to 2/3 ² 2/3 ³	RESTRICTION: Use on sandy mineral soils only. Pre-Harvest Interval (PHI): Do not harvest spinach within 21 days after application. Pre-Harvest Interval (PHI) in OREGON: Do not harvest spinach within 45 days after application.

* Make only one (1) application per year. If Ro-Neet Herbicide is applied in the fall, do not reapply Ro-Neet Herbicide the following spring.

¹ For use in Arkansas, Colorado, Connecticut, Delaware, Illinois, Maine, Maryland, Massachusetts, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oklahoma, Oregon, Pennsylvania, South Carolina, Texas, Vermont, Virginia, Washington, and western Tennessee.

² Arizona only.

³ California only.

RESTRICTIONS:

- Maximum of 2/3 gallon (4 pounds active ingredient) per acre per single application.
 - **WASHINGTON:** Maximum of 1/2 gallon (3 pounds active ingredient) per acre per single application.
- Do not exceed one application per year.
- Apply Ro-Neet Herbicide in a minimum of 10 gallons of spray solution per acre.
- **ARIZONA and OREGON:** Apply Ro-Neet Herbicide in a minimum of 20 gallons of spray solution per acre.
- **ARIZONA, OREGON and WASHINGTON:** Immediately (within minutes) after application, thoroughly mix Ro-Neet Herbicide into the soil to a depth of 2 to 3 inches using a finishing disk, harrow, rolling cultivator or similar implement capable of providing uniform incorporation.
- Do not use on Spinach Grown for Seed or canning, unless otherwise specified by Special Local Need Label.
- All spinach seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on farm disposal, disposal at a controlled dumpsite, incinerator, composter, or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
- No portion of the spinach seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Spinach seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Spinach seed may not be distributed for human consumption or animal feed.

Rotation Interval: Crops may be rotated immediately to other labeled crops (spinach, table beets or sugar beets); however, all crops without a cycloate registration need to be planted back at intervals of at least 45 days for leafy vegetables, 45 days for root and tuber vegetables and 120 days for all other crops, to ensure that quantifiable residues are not found in rotated crops.

Band Treatment

The rates given above are for broadcast treatment. For band treatment, reduce the amount of Ro-Neet Herbicide proportionately depending on the row spacing and band width to be treated. **DO NOT USE BAND APPLICATION ON ROCKY GROUND BECAUSE THOROUGH INCORPORATION IS NOT POSSIBLE.**

Soil Injection Application on Sugar Beets: Special equipment designed for soil injection must be used. The injector units must be rigidly mounted immediately ahead or immediately behind the planter unit.

Soil Injection: Injector shanks must be spaced 2 ½ to 3 inches apart and mounted in staggered positions to avoid trash building up on them. Injector shanks must be set to inject Ro-Neet Herbicide 1 1/2 to 2 inches below the soil surface. The width of the band in which weed control is desired will determine the number of injector shanks required per row (that is, four injector shanks for a 10 to 12-inch band, six injector shanks for a 15 to 18-inch band). The two injector shanks adjacent to the drill row must be 1 1/4 to 1 1/2 inches on either side of it.

RO-NEET HERBICIDE AND PRODUCTS CONTAINING THE ACTIVE INGREDIENT EPTC TANK MIX FOR PREPLANT USE IN SUGAR BEETS FOR MICHIGAN, MINNESOTA, OHIO, AND THE RED RIVER VALLEY AREA OF NORTH DAKOTA.

TANK MIX - DIRECTIONS FOR USE

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

The Ro-Neet Herbicide and EPTC combination is a selective tank mix which controls weeds by interfering with normal germination and seedling development. This tank mixture can be applied only once per year. The combination may be used in the fall or in the spring (except where noted, see table below).

The Ro-Neet Herbicide and EPTC combination may cause crop injury on very light sandy soil and when used under adverse environmental conditions that weaken crop seedlings.

A tank mixture of Ro-Neet Herbicide and products containing EPTC will give equal to or greater control of the following listed weeds than either product used separately. This combination does not control established weeds.

Green foxtail	<i>Setaria viridis</i>
Yellow foxtail	<i>Setaria glauca</i>
Wild oats	<i>Avena fatua</i>
Yellow nutsedge	<i>Cyperus esculentus</i>
Purple nutsedge	<i>Cyperus rotundus</i>
Lambsquarters, common*	<i>Chenopodium album</i>
Pigweed, redroot*	<i>Amaranthus retroflexus</i>

* Partial control only (suppression)

APPLICATION DIRECTIONS

During filling, pour the directed rates of both Ro-Neet Herbicide and EPTC into a properly calibrated, low-pressure boom sprayer having good agitation, and mix thoroughly. Apply the material uniformly in 10 to

50 gallons of water per acre.

- For Arizona and Oregon apply in a minimum of 20 gallons of spray solution per acre.

Check calibration frequently during application and observe the nozzles to ensure a uniform spray pattern. The soil should be well worked prior to application and dry enough to permit thorough mixing with incorporation equipment.

Removable chemical extraction probes (also known as "stingers") used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

SOIL INCORPORATION

The Ro-Neet Herbicide and EPTC tank mixture must be immediately incorporated (mixed) into the top 2 to 3 inches of soil after spraying to prevent loss of the herbicides.

SPRING* PREPLANT INCORPORATED APPLICATION RATES** (Pints/Acre***)			
SOIL TEXTURE	ORGANIC MATTER %	EPTC PRODUCT Containing 7 pounds active ingredient per gallon.	RO-NEET HERBICIDE
Coarse	3.0 to 4.5 > 4.5	1.0 1.0 to 1.25	3.0 3.0
Medium	3.0 to 4.5 > 4.5	1.0 to 1.25 1.0 to 1.75	3.0 to 4.0 3.0 to 4.0
Fine	3.0 to 4.5 > 4.5	1.0 to 1.5 1.0 to 2.0	3.0 to 4.0 3.0 to 4.0

* DO NOT USE THE TANKMIX IN THE SPRING IN MICHIGAN OR OHIO.

** Make only one application per year. Do not apply the Ro-Neet Herbicide / EPTC combination in the spring if either this combination or Ro-Neet Herbicide alone was applied in the fall.

*** Do not apply more than 5 pints/acre of Ro-Neet Herbicide and EPTC combined regardless of the ratio.

FALL PREPLANT INCORPORATED APPLICATION RATES* (Pints/Acre**)			
SOIL TEXTURE	ORGANIC MATTER %	EPTC PRODUCT Containing 7 pounds active ingredient per gallon.	RO-NEET HERBICIDE
Coarse	3.0 to 4.5 > 4.5	1.0 1.0 to 1.25	4.0 4.0 to 4.33
Medium	3.0 to 4.5 > 4.5	1.0 to 1.25 1.0 to 1.5	4.0 to 4.5 4.0 to 4.5
Fine	3.0 to 4.5 > 4.5	1.0 to 1.5 1.0 to 2.0	3.0 to 4.5 4.5

* Make only one application per year. Do not apply the Ro-Neet Herbicide / EPTC combination in the spring if either this combination or Ro-Neet Herbicide alone was applied in the fall.

** Do not apply more than 6 pints/acre of Ro-Neet Herbicide and EPTC combined regardless of the ratio.

RO-NEET HERBICIDE WITH FLUID FERTILIZERS

The following procedure is suggested for determining whether Ro-Neet Herbicide may be combined with a specific fluid fertilizer for spray tank application.

MATERIALS REQUIRED

1. Ro-Neet Herbicide
2. Fluid fertilizer to be used.
3. Adjuvant for fertilizer tankmix: Compex™, Sponto 168-D™, Unite™, or equivalent. The adjuvant which provides the best emulsification depends on the specific fertilizer under consideration.
4. Two one-quart, wide-mouth glass jars with lid or stopper.
5. Measuring spoons (a 25 ml pipette or graduated cylinder provides more accurate measurement).
6. Measuring cup, 8 oz. (237 ml).

PROCEDURE

1. Pour a pint (about 473 ml) of the fluid fertilizer into each of the quart jars.
2. Add adjuvant to one of the jars and mix (see rate table).
3. Add the Ro-Neet Herbicide to both jars (see rate table).
4. Close both jars with lid or stopper and mix the contents by turning the jars upside down ten times.
5. Inspect the surface and body of the mixtures.
 - (A) Immediately after completing the jar inversions.
 - (B) After allowing the jars to stand quietly for 30 minutes.
 - (C) And then again after turning the jars upside down 10 times.

If the uniform mix cannot be made, the mixture should not be used. If either mixture remains uniform for 30 minutes, the combination may be used. Should either mixture separate after 30 minutes, but readily remix uniformly with the jar inversions, the mixture can be used if adequate agitation is maintained in the tank. If the mixture with adjuvant is satisfactory, but the one without adjuvant is not, be sure to use the adjuvant in the spray tank. Add the adjuvant first at a rate of 3 pints per 100 gallons of fluid fertilizer; foaming can be minimized by using moderate agitation. If nondispersible oil, sludge, or clumps of solids form in the mixtures, the combinations should not be used.

RATE TABLE FOR Ro-Neet Herbicide AND ADJUVANT** WITH THE FLUID FERTILIZER		
Gallons of fluid fertilizer to be applied per acre	ml or tsp. of Ro-Neet Herbicide* to be added to 1 pint of fertilizer	
	Ro-Neet Herbicide	
	ml	tsp
10	8	1 2/3
15	5	1
20	4	3/4
25	3	2/3
30	2 1/2	1/2
40	2	2/5

* Based on field rate of 1 pound active ingredient per acre in the fertilizer volumes indicated. Increase volume proportionately to correspond with intended field rate in terms of pounds of active ingredient per acre (e.g., for field rate of 4 pounds actual Ro-Neet Herbicide in 40 gallons fertilizer per acre, add 8 ml or 2 tsp Ro-Neet Herbicide to each jar for compatibility testing purposes).

** Two (2) milliliters or one-half (1/2) teaspoon of adjuvant to be added to 1 pint of fluid fertilizer in order to equal the rate of 3 pints of adjuvant per 100 gallons of fluid fertilizer.

WARRANTY DISCLAIMER

The EPA approved directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.



HELM Agro US, Inc.

401 E. Jackson St., Suite 1600

Tampa, FL 33602

Tel. 813-621-8846

Ro-Neet® is a trademark of HELM Agro US, Inc.

Compex™ is a trademark of Kalo, Inc.

Micro-Cel™ is a trademark of Manville Sales Corporation.

Sponto 168-D™ is a trademark of Witco Corporation.

Unite® is a trademark of HACO, Inc.

(RV013025)

{Note to Reviewer

The following pages are for the following SUPPLEMENTAL LABELING (expiring on December 31, 2027):

- Use is from Washington 24(c) on Spinach / SLN No. WA-210008 }



CYCLOATE

GROUP

8

HERBICIDE

Ro-Neet® Herbicide

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. "Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental.

EPA Reg. No. 74530-16

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This supplemental label expires on September 5, 2027 and cannot be used or distributed after that date.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS IN THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL.

Read the label affixed to the container for Ro-Neet Herbicide before applying.

Use of Ro-Neet Herbicide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Ro-Neet Herbicide.

SPINACH IN WASHINGTON

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

SPINACH: Pre-Plant: Apply a surface broadcast application with ground application equipment prior to planting. Apply 1/3 to 1/2 gallons Ro-Neet Herbicide (2 to 3 lbs. ai) per acre on sandy mineral soils only in a minimum of 10 gallons of water. Immediately after application, thoroughly mix Ro-Neet Herbicide into the soil to a depth of 2 to 3 inches using a finishing disk, harrow, rolling cultivator or similar implement capable of providing uniform incorporation.

WEEDS CONTROLLED: For List of Weeds Controlled See EPA-registered Ro-Neet Herbicide Label.

RESTRICTIONS:

- Re-entry interval is 48 hours.
- DO NOT apply this product through any type of irrigation system.
- DO NOT harvest spinach for 45 days following application of this product.
- DO NOT make more than one (1) application per growing year. If Ro-Neet Herbicide is applied in the fall, do not reapply Ro-Neet Herbicide the following spring.

- DO NOT apply more than 3 lbs. active ingredient/acre per year.
- All spinach seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on farm disposal, disposal at a controlled dumpsite, incinerator, composter, or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
- No portion of the spinach seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Spinach seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Spinach seed may not be distributed for human consumption or animal feed.

WSDA CONTAINER DISPOSAL GUIDANCE: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program website at <https://agr.wa.gov/wastepesticide>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

IMPORTANT: Read the entire Directions for use and the Conditions of Sale and Warranty before using this product.

Ro-Neet is a registered trademark of HELM Agro US, Inc.

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

(RV013025)

Manufactured for:
HELM Agro US, Inc.
 401 E. Jackson St., Suite 1600
 Tampa, FL 33602