



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
1200 Pennsylvania Avenue, N.W.
Washington, D.C. 20460

EPA Reg. Number:
74530-31

Date of Issuance:
MAR 17 2008

Term of Issuance:
Conditional

Name of Pesticide Product:
KENDO EC Insecticide

NOTICE OF PESTICIDE:

- Registration
 Reregistration

(Under FIFRA as amended)

Name and Address of Registrant (include ZIP Code):

Helm Agro US, Inc.
8295 Tournament Drive, Suite 310
Memphis, TN 38125

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.


Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is **conditionally** registered in accordance with FIFRA sec. 3(c)(7)(A) subject to the following provisions:

1. You will submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. You will make the following label changes before you release the product for shipment:
 - a) Revise the EPA Registration Number to read "EPA Reg. No. 74530-31."
 - b) On page 21, under "Turfgrass" revise the entry for ticks to read "Ticks (including species which may transmit Lyme disease)"
 - c) Under "Conditions of Sale and Warranty" revise the sentence "The directions for use of this product are believed..." by changing the word "should" to "must."
 - d) The container disposal statements for pesticides have been revised. Refer to PR Notice 2007-4 and modify your label accordingly
 - e) Add "To the extent consistent with applicable law" in front of:
 - o "Helms Agro US, Inc. makes no other express or implied warranty..."
 - o "In no case shall Helms Agro US, Inc. or the seller..."

- 3. Please submit two (2) copies of your final printed labeling before releasing the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing revised labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact Olga Odiott of my team at (703) 308-9369.

Signature of Approving Official:  George T. LaRocca, Product Manager (13) Insecticide Branch/Registration Division (7505P)	Date: MAR 17 2008
--	--------------------------

Enclosure

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS
 For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Helm Agro US, Inc.
KENDO EC Insecticide
 For Agricultural and Turf and Ornamental Use

Active Ingredient

Lambda-cyhalothrin [1a(S*),3a(Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate	11.4%
Inert Ingredients	88.6%
Total	100.0%

Contains 1 lb. of active ingredient per gallon.
 Contains petroleum distillates.

EPA Reg. No. 74530-

EPA Est. No. 5905-GA-1

KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, give artificial respiration immediately, preferably by mouth-to-mouth. • Call a poison control center or doctor for treatment advice.
Note to Physician: Contains petroleum distillates. Vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.	

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed. Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as Barrier Laminate, or Viton ≥14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ACCEPTED
with COMMENTS
In EPA Letter Dated
MAR 17 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
74530-31



Helm Agro US, Inc.
 8295 Tournament Drive
 Suite 310
 Memphis, Tennessee 38125

User Safety Recommendations**USERS SHOULD:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE/clothing immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material such as Barrier Laminate, or Viton ≥ 14 mils

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep adults, children, and pets off treated areas until spray has dried following the application.

SPRAY DRIFT PRECAUTIONS

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

- Do not apply by ground within 25 ft., or by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 ft. when ultra-low volume (ULV) application is made.
- When treating very tall trees (> 15 ft. from the ground) with high pressure sprays or during any application with air assisted equipment (mist blowers), do not apply within 150 ft. of aquatic areas.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3-10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 ft. of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity or temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by production smoke and observing a smoke layer near the ground surface.

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage files must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial and ground application. For aerial applications, the 25 ft. vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

CHEMIGATION

Sprinkler Irrigation Application

Apply KENDO EC Insecticide using rates and timing described on this label. Consultation with your local State Extension Service or other local experts may be useful for recommendations on which adjuvants or diluent types to use, (see Tank Mix Applications section) as well as for rates and mixing instructions. Ascertain that the recommendations have been proven, through university and extension field trials, to be effective with this product applied by chemigation.

Be sure the irrigation system is providing uniform application of water to all areas, because good control requires thorough coverage of foliage. Maintain continuous agitation in the pesticide supply tank before and during the entire application period.

Inject the recommended rate of KENDO EC Insecticide into the irrigation system by means of a metering device that will provide a constant flow and distribute the product to the desired area in 0.1-0.2 inch of water. It is recommended that the minimum amount of water be used that will provide proper distribution and coverage. Inject the product into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Following application, flush the entire irrigation and injection system with clean water before stopping it.

If application is being made during a normal irrigation set of a stationary sprinkler, inject the recommended rate of KENDO EC Insecticide for the area covered into the system only during the end of the irrigation set for a sufficient time to provide adequate coverage and product distribution.

It is not recommended that KENDO EC Insecticide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

USE PRECAUTIONS: SPRINKLER IRRIGATION

APPLICATION

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-resource contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

GENERAL DIRECTIONS FOR USE

Thorough crop coverage is necessary for both initial and residual control. Apply by ground in at least 10 gal./A or by air in at least 2 gal./A using sufficient water to obtain full coverage of foliage unless this label specifies otherwise. In situations where foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), control can be improved by use of higher application volumes and/or higher use rates.

For cutworm control, KENDO EC Insecticide may be applied before, during, or after planting. When making soil incorporated applications, use higher rates for better control.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, it is possible that this product or other products with a similar mode of action, may not provide satisfactory control. If poor performance cannot be related to incorrect application or extreme weather conditions, a resistant strain of insect may be the reason. If control does not meet expectations and resistance is a possible cause, consult your local company representative or agricultural advisor immediately for the best alternative options for your area.

Tank Mix Applications

KENDO EC Insecticide may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. Adjuvants such as spreader stickers, wetting agents, and penetrants may also be added. Use a small volume mixing test with the other products to confirm compatibility. If other chemicals are added to the applicator tank, KENDO EC Insecticide should be added last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours. Observe all restrictions and precautions found on labels of products in the tank mix.

CROP USES AND SPRAY RECOMMENDATIONS

ALFALFA, ALFALFA GROWN FOR SEED

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Alfalfa Caterpillar Cutworm spp. Green Cloverworm Leafhopper spp. Looper spp. Three-cornered Alfalfa Hopper Velvetbean Caterpillar Webworm spp.	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use.
Alfalfa Seed Chalcid (Adult) Alfalfa Weevil Armyworm Bean Leaf Beetle (Adult) Blister Beetle spp. Blue Alfalfa Aphid Clover Leaf Weevil spp. Clover Root Borer (Adult) Clover Root Curculio spp. (Adult) Clover Stem Borer (Adult) Corn Earworm Cowpea Aphid Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle spp. (Adult) Egyptian Alfalfa Weevil Fall Armyworm ¹ Grape Colaspis (Adult) Grasshopper spp. Green June Beetle (Adult) Green Peach Aphid ³ Japanese Beetle (Adult) Meadow Spittlebug Mexican Bean Beetle Pea Aphid Pea Weevil (Adult) Plant Bug spp. Including Lygus spp. ³ Spotted Alfalfa Aphid Stink Bug spp. Thrips spp. Sweet Clover Weevil (Adult) Western Yellow-Striped Armyworm Whitefringed Beetle spp. (Adult) Yellow-Striped Armyworm	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	Do not apply more than 0.03 lb. a.i. (0.24 pt./A) per cutting. Do not apply more than 0.12 lb. a.i. (0.96 pt./A) per season. Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.
Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ²	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Apply only to fields planted to pure stands of alfalfa. Use scouting to determine applications. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. Apply in at least 2 gal./A by air or 10 gal./A by ground. In situations of dense foliage and/or high pest populations, use 5-10 gal./A by air or 20 gal./A by ground and higher use rates. Also use higher rates for improved residual control.</p> <p>Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters.</p>		

CANOLA

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Armyworm spp. Cabbage Seedpod Weevil Cutworm spp. Diamondback Moth Flea Beetle Grasshoppers Looper spp. Lygus Bug	0.015-0.03 lb. a.i. (1.92-3.84 fl. oz.)	See additional instructions below. Do not apply within 7 days of harvest. Do not apply more than 0.09 lb. a.i. (0.72 pt./A) per year.
Cabbage Aphid	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Use scouting to determine applications, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by air or ground with enough water to obtain full coverage of foliage. For air applications, apply a minimum of 2 gal. of water/A.</p>		

CEREAL GRAINS – CORN (AT PLANT): FIELD CORN, POPCORN, SEED CORN, SWEET CORN

Pests	Rate KENDO EC Insecticide per 1,000 ft. of Row	Remarks
Corn Rootworm Larvae (Western, Northern, Southern, Mexican) Cutworm spp. Lesser Cornstalk Borer Red Imported Fire Ant ¹ Seedcorn Beetle Seedcorn Maggot Wireworm spp. ¹ White Grub spp.	0.005 lb. a.i. (0.66 fl. oz.)	See additional instructions below. Do not harvest or graze livestock or cut treated crops for feed within 21 days of at plant application. Do not apply more than 0.09 lb. a.i. (0.72 pt./A) per crop at plant. For field corn, popcorn, and seed corn do not apply more than 0.12 lb. a.i./A per crop from at plant and foliar applications. For sweet corn do not apply more than 0.48 lb. a.i./A per crop from at plant and foliar applications. For Banded Applications – Make application at planting as a 5-7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel. For In-Furrow Applications – Make application into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel. Apply a minimum of 3 gal. finished spray/A. ¹ Suppression only.

lbs. a.i. and fl. oz./A of KENDO EC Insecticide applied at 0.66 fl. oz./1000 ft. of row for various row spacings.

Row Spacing	40"	38"	36"	34"	32"	30"
Linear ft./A	13,068	13,756	14,520	15,374	16,335	17,424
lbs. a.i./A	0.067	0.07	0.075	0.079	0.084	0.09
fl. oz./A	8.6	9.1	9.6	10.1	10.8	11.5

CEREAL GRAINS – CORN (FOLIAR): FIELD CORN, POPCORN, SEED CORN

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Corn Earworm ¹ Cutworm spp. Green Cloverworm Meadow Spittlebug Western Bean Cutworm ¹	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control before the larva bores into the plant stalk or ear. ² For control of first and second instars only.
Armyworm ² Bean Leaf Beetle Bird Cherry-Oat Aphid ³ Cereal Leaf Beetle Corn Leaf Aphid ³ Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) English Grain Aphid ³ European Corn Borer ¹ Fall Armyworm ² Flea Beetle spp. Grasshopper spp. Hop vine Borer ¹ Japanese Beetle (Adult) Lesser Cornstalk Borer Sap Beetle (Adult) Seedcorn Beetle Southwestern Corn Borer ¹ Stalk Borer ¹ Stink Bug spp. Tobacco Budworm ^{1,4} Webworm spp. Yellow-Striped Armyworm ²	0.02-0.03 lb./A (2.56-3.84 fl. oz.)	³ Suppression only. ⁴ See resistance statement under General Directions for Use. Do not apply within 21 days of harvest. Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. Do not apply more than 0.12 lb. a.i. (0.96 pt./A) per crop from at plant and foliar applications. Do not apply more than 0.06 lb. a.i. (0.48 pt.) after silk initiation. Do not apply more than 0.03 lb. a.i. (0.24 pt.) after corn has reached the milk stage (yellow kernels with milky fluid).
Beet Armyworm ^{2,4} Chinch Bug Green Bug ^{3,4} Mexican Rice Borer ¹ Rice Stalk Borer ¹ Southern Corn Leaf Beetle Sugarcane Borer ¹	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Use scouting or locally prescribed corn growth stages to determine application, usually at intervals of 7 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in at least 2 gal. of water/A.</p> <p>For chinch bug control, begin application when bugs migrate from small grains or grass weeds to small corn and direct the spray to the base of corn plants. Make additional applications at 3-5 day intervals if needed. KENDO EC Insecticide may only suppress heavy infestations and/or subsequent migrations.</p> <p>For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied corn rootworm control program use at least 3.84 fl. oz./A (0.03 lb. a.i./A).</p>		

CEREAL GRAINS – SWEET CORN (FOLIAR)

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Aphid spp. ^{2,3} Aster Leafhopper Beet Armyworm ^{1,2} Cereal Leaf Beetle Chinch Bug Common Cornstalk Borer Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) Corn Earworm Cutworm spp. European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Sap Beetle (Adult) Southern Armyworm ¹ Southwestern Corn Borer Spider Mite spp. ² Stink Bug spp. Tarnished Plant Bug Yellow-Striped Armyworm ¹ Western Bean Cutworm Webworm spp.	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under general Directions for Use. Do not apply within 1 day of harvest. Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. Do not apply more than 0.48 lb. a.i. (3.84 pt./A) per crop from at plant and foliar applications.
Chinch Bug Corn Silkfly (Adult) ² Green Bug ^{2,3}	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Use scouting or locally prescribed corn growth stages to determine application, usually at intervals of 4 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. For best results target control before insects enter the stalk or ear.</p> <p>Apply ground or air using enough water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in at least 2 gal. of water/A.</p> <p>For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied corn rootworm control program use a minimum of 0.025 lb. a.i. (3.2 fl. oz./A).</p>		

CEREAL GRAINS - RICE

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Bird Cherry-Oat Aphid Chinch Bug European Corn Borer ¹ Fall Armyworm Grasshopper spp. Green Bug Leafhopper spp. Mexican Rice Borer ¹ Rice Seed Midge Rice Stalk Borer ¹ Rice Stink Bug Rice Water Weevil (Adult) Sharpshooter spp. Sugarcane Borer ¹ True Armyworm Yellow-Striped Armyworm Yellow Sugarcane Aphid	0.025-0.04 lb. a.i. (3.20-5.12 fl. oz.)	See additional instructions below. ¹ For control before the larvae bore into the plant stalk. Do not release flood water within 7 days of an application. Do not apply more than 0.12 lb. a.i. (0.96 pt.)/A per season. Do not apply more than 0.08 lb. a.i. (0.64 pt.)/A within 28 days of harvest or more than 0.04 lb. a.i. (0.32 pt.)/A within 21 days of harvest. Do not apply within 21 days of harvest. Do not use treated rice fields for the aquaculture of edible fish and crustacea. Do not apply as an ultra-low volume (ULV) spray.

Use scouting to determine timing of application and the need for repeat applications, usually at 5-7 day intervals. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.

KENDO EC Insecticide can be safely used when propanil products are being used for weed control.

Apply by air or ground using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water (or a total carrier volume)/A but ensure that application is made in sufficient volume to provide adequate coverage. When applying at lower volumes by air, the addition of an emulsifiable crop oil (e.g. 1 pt./A) is recommended to help improve coverage, reduce evaporation, and improve efficacy.

For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not allow more than 10 days to elapse from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Treatment of adults may also be made at later stages of rice development to reduce overwintering populations.

For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars, usually when rice has emerged 1/2 inch above the waterline. When there is prolonged migration into the field, begin field scouting for adults and/or feeding scars 3-5 days after the first treatment and, if needed, make a second application within 7-10 days of the first application. Treatment of adults may also be made at later stages of rice development to reduce overwintering populations.

California: In addition to the directions above for control of rice water weevil in water seeded rice, KENDO EC Insecticide may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable both on levees and in the water. Larvae are vulnerable while feeding on the leaves before they enter the soil. Monitor field edges and levee areas for adults, then treat in one of the following ways: spray the inside perimeter of the field, or spray the entire field.

Because Green bug is known to have many biotypes, it is possible that KENDO EC Insecticide may only provide suppression. If the first application of KENDO EC Insecticide does not give satisfactory control, a resistant biotype may be present and use of an alternate chemistry may be necessary.

CEREAL GRAINS – SORGHUM (GRAIN)

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Sorghum Midge	0.015-0.02 lb. a.i. (1.92-2.56 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only.
Armyworm Beet Armyworm ^{1,3} Corn Earworm European Corn Borer ² Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Lesser Cornstalk Borer ² Southwestern Corn Borer ² Stink Bug spp. Webworm spp. Yellow-Striped Armyworm ¹	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	² For control before the larva bores into the plant stalk. ³ See resistance statement under General Directions for Use. Do not apply more than 0.08 lb. a.i. (0.64 pt.)/A per season. Do not apply more than 0.06 lb. a.i. (0.48 pt.)/A per season after crop emergence. Do not apply more than 0.02 lb. a.i. (0.16 pt.)/A per season once crop is in soft dough stage.
Chinch Bug	0.03 lb. a.i. (3.84 fl. oz.)	Do not apply within 30 days of harvest. Do not graze livestock in treated areas or harvest for fodder, silage, or hay.
<p>Use scouting to determine treatment, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of target location. When applying by air, apply in at least 2 gal. of water/A.</p> <p>For sorghum midge control, make first application when one quarter of the sorghum heads have emerged and are in tip bloom. If needed, repeat applications at 5-day intervals.</p> <p>For chinch bug control, start applications when bugs migrate from small grains or grass weeds to small sorghum, directing spray to the base of sorghum plants. If needed, repeat applications at 3-5 day intervals.</p> <p>KENDO EC Insecticide may only suppress heavy infestations and/or subsequent migrations.</p>		

CEREAL GRAINS – WHEAT, WHEAT HAY, AND TRITICALE

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Army Cutworm Cutworm spp.	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, KENDO EC Insecticide may provide suppression only. Higher rates and increased coverage will be necessary.
Armyworm Bird Cherry-Oat Aphid ¹ Cereal Leaf Beetle English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grasshopper spp. Hessian Fly ³ Orange Blossom Wheat Midge Russian Wheat Aphid ¹ Stink Bug spp. Yellow-Striped Armyworm	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	² See resistance statement under General Directions for Use. ³ Make application when adults emerge. Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pt.)/A per season.
Grass Sawfly	0.025-0.03 lb. a.i. (3.20-3.84 fl. oz.)	
Chinch Bug Corn Leaf Aphid ² Greenbug ^{1,2} Mite spp. ²	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Use scouting to determine treatment, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.</p> <p>For chinch bug control, repeat applications at 3-5 day intervals if needed. KENDO EC Insecticide may only suppress heavy infestations and/or migrations.</p> <p>Because Greenbug is known to have many biotypes, it is possible that KENDO EC Insecticide may only provide suppression. If this occurs, a second application using an alternative chemistry may be needed.</p>		

COLE CROPS – BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAVALO BROCCOLO, CAULIFLOWER, CHINESE BROCCOLI (GAI LON), CHINESE CABBAGE (NAPA), CHINESE MUSTARD CABBAGE (GAI CHOY), KOHLRABI

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Alfalfa Looper Cabbage Looper Cabbage Webworm Cutworm spp. Imported Cabbageworm Southern Cabbageworm	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use.
Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Lygus spp. ³ Meadow Spittlebug Aphid spp. ^{2,3} Plant Bug spp. – including Stink Bug spp. Spider Mite spp. ² Thrips spp. ² Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellow-Striped Armyworm	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	Do not apply within 1 day of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pt.)/A per season.
Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.		
Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gals. of water/A. ✓		

COTTON

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Soybean Thrips Tobacco Thrips	0.015-0.025 lb. a.i. (1.92-2.56 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only.
Cabbage Looper Cotton Fleahopper Cotton Leafperforator Cotton Leafworm Lygus Bug spp. ³ Pink Bollworm Saltmarsh Caterpillar	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	³ See resistance statement under General Directions for Use. Do not apply within 21 days of harvest. Do not graze livestock in treated areas. Do not apply more than 0.2 lb. a.i. (1.6 pt.)/A per season.
Bandedwing Whitefly ^{2,3} Beet Armyworm ^{1,3} Boll Weevil Brown Stink Bug Cotton Aphid ^{2,3} Cotton Bollworm European Corn Borer Fall Armyworm Green Stink Bug Southern Green Stink Bug Sweetpotato Whitefly ^{2,3} Tobacco Budworm ³ Twospotted Spider Mite ²	0.025-0.04 lb. a.i. (3.20-5.12 fl. oz.)	Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.
Use scouting to determine application, usually at intervals of 5-7 days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.		
Apply ground or air using enough water to obtain full coverage of foliage.		
Applications may also be made with equipment adapted and calibrated for ULV sprays. KENDO EC Insecticide may be mixed with once-refined vegetable oil and applied in a minimum of at least one qt. of finished spray/A.		
When bollworm/budworm infestation levels are light, 0.02 lb. a.i./A may be applied in together with intense field monitoring.		
Boll weevil: spray on a 3-5 day schedule.		
When applied according to the directions above for control of cotton bollworm and tobacco budworm, KENDO EC Insecticide also provides ovicidal control of unhatched <i>Heliothis</i> spp. eggs.		

FRUITING VEGETABLES – TOMATO, TOMATILLO, PEPPERS (BELL AND NONBELL), EGGPLANT, GROUND CHERRY, PEPINO

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cabbage Looper Cutworm spp. Hornworm spp.	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only.
Aphid spp. ^{2,3} Beet Armyworm ^{1,3} Blister Beetle spp. Colorado Potato Beetle ³ Cucumber Beetle spp. (Adult) European Corn Borer ⁴ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leafminer spp. ² Meadow Spittlebug Pepper Weevil (Adult) ² Plant Bug spp. Southern Armyworm ¹ Spider Mite spp. ² Stalk Borer ⁴ Stink Bug spp. Thrips ⁵ Tobacco Budworm ³ Tomato Fruitworm Tomato Pinworm Tomato Psyllid ^{2,3} Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellow-Striped Armyworm ¹	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	² Suppression only. ³ See resistance statement under General Directions for Use. ⁴ For control before the larva bores into the plant stalk or fruit. ⁵ Does not include Western Flower thrips. Do not apply within 5 days of harvest. Do not apply more than 0.36 lb. a.i. (2.88 pt./A) per season.
<p>Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.</p>		

LEGUME VEGETABLES (BEANS AND PEAS):

EDIBLE PODDED (ONLY): *Canavalia gladiata* – sword bean; *Canavalia ensiformis* – jackbean; *glycine max* – soybean (immature seed).

EDIBLE PODDED, SUCCULENT SHELLED OR DRIED SHELLED: *Phaseolus* spp. – includes field, kidney, lima, navy, pinto, runner, snap, tepary, and wax beans; *Vigna* spp. – includes adzuki, asparagus, moth, mung, rice, urd and yardlong beans, blackeyed pea, caljang, Chinese longbean, cowpea, Crowder pea, and Southern pea; *Pisum* spp. – includes dwarf, edible-pod, English, field, garden, green, snow, and sugar peas; *Cajanus cajan* – Pigeon pea.

SUCCULENT SHELLED OR DRIED SHELLED: *Vicia faba* – broadbean (fava bean).

DRIED SHELLED (ONLY): *Lupinus* spp. – includes grain, sweet, white and sweet white lupines; *Cicer arietinum* – chickpea (garbanzo bean), *Cyamopsis tetraganloba* – guar, *Lablab purpureus* – Lablab bean (hyacinth bean), *Lens esculata* – lentils.

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Green Cloverworm Imported Cabbageworm Mexican Bean Beetle Saltmarsh Caterpillar Velvetleaf Caterpillar	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control before the larva bores into the plant stalk or pods. ² For control of first and second instars only. ³ For suppression only.
Alfalfa Caterpillar Aphid spp. ⁴ Armyworm ² Bean Leaf Beetle Bean Leafskeletonizer Blister Beetle spp. Corn Earworm Corn Rootworm Beetle spp. (Adult) Cucumber Beetle spp. (Adult) Curculio and Weevil spp. ¹ (foliage and pod feeding adults and larvae) European Corn Borer ¹ Fall Armyworm ² Flea Beetle spp. (Adult) Flea Hopper spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leaf-tier spp. Looper spp. Meadow Spittlebug Painted Lady Butterfly (Larva) Plant Bug spp. including Lygus spp. ⁴ Stalk Borer ¹ Stink Bug spp. Three-Cornered Alfalfa Hopper Thrips spp. ⁴ Tobacco Budworm ⁴ Webworm spp. Western Yellow-Striped Armyworm ² Yellow-Striped Armyworm ²	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	⁴ See resistance statement under General Directions for Use. For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest. For dried shelled legume vegetables, do not apply within 21 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pt./A per season). For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay.
Beet Armyworm ^{2,3,4} Leafminer spp. ^{3,4} Lesser Cornstalk Borer ³ Spider Mite spp. ³ Whitefly spp. ^{3,4}	0.03 lb. a.i. (3.84 fl. oz.)	
Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.		

LEGUME VEGETABLES (PEAS AND BEANS): SOYBEANS

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Bean Leaf Beetle Cabbage Looper Chinese Aphid ⁴ Corn Earworm Corn Rootworm Beetle (Adult beetles including Mexican, Northern, Southern, Western) Cutworm spp. Green Cloverworm Mexican Bean Beetle Painted Lady (Thistle) Caterpillar Potato Leafhopper Saltmarsh Caterpillar Soybean Aphid ⁴ Three-Cornered Alfalfa Hopper Thrips spp. Velvetbean Caterpillar Woollybear Caterpillar	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use. ⁴ Use lower rates for early season applications and/or lighter populations. Do not apply within 45 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pt./A) per season.
Armyworm ¹ Blister Beetle spp. European Corn Borer ¹ Fall Armyworm ¹ Grasshopper spp. Japanese Beetle (Adult) Plant Bug spp. Silverspotted Skipper Stink Bug spp. Tobacco Budworm ³ Webworm spp. Yellow-Striped Armyworm ¹	0.025-0.03 lb. a.i. (3.20-3.84 fl. oz.)	
Beet Armyworm ^{1,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ²	0.03 lb. a.i. (3.84 fl. oz.)	
Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Do not graze or harvest treated soybean forage, straw or hay for livestock feed. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A. For control of adult corn rootworm beetles (<i>Diabrotica</i> spp.) as part of an aerial applied to corn rootworm control program use at least 2.56 fl. oz./A (0.02 lb. a.i./A).		

LETTUCE (HEAD AND LEAF)

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Alfalfa Cabbage Looper Cutworm spp. Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use.
Aphid spp. ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Plant Bug spp. including Lygus spp. ³ Southern Armyworm Spider Mite spp. ² Stink Bug spp. Tobacco Budworm ³ Vegetable Weevil (Adult) Whitefly spp. ^{2,3}	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	Do not apply within 1 day of harvest. Do not apply more than 0.3 lb. a.i. (2.4 pt./A) per season.
Use scouting to determine application; usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.		

ONION (BULB) AND GARLIC

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Leafminer spp. (Adult) Onion Maggot (Adult) Seedcorn Maggot (Adult)	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only.
Aphid spp. ² Armyworm spp. ¹ Flower Thrips ² Onion Thrips Plant Bug spp. Stink Bug spp. Tobacco Thrips Western Flower Thrips ^{2,3}	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	³ See resistance statement under General Directions for Use. Do not apply within 14 days of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pt.)/A per season.
<p>Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Use the higher label rates as thrips population increases and avoid rescue situations.</p> <p>Apply by ground or air using enough water and application methods to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.</p> <p>To control thrips by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturer's use directions) may improve the deposition of the spray and increase plant coverage.</p>		

PEANUT

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Green Cloverworm Potato Leafhopper Red-necked Peanut Worm Three-Cornered Alfalfa Leafhopper Velvetbean Caterpillar	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use.
Bean Leaf Beetle Corn Earworm Fall Armyworm ¹ Grasshopper spp. Southern Corn Rootworm (Adult) Stink Bug spp. Tobacco Thrips Vegetable Weevil Whitefringed Beetle (Adult)	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	Do not apply within 14 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pt.)/A per season.
Aphid spp. ² Beet Armyworm ^{1,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ²	0.03 lb. a.i. (3.84 fl. oz.)	
<p>Use scouting to determine application, usually at intervals of 7 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.</p> <p>Apply by ground or air using enough water to obtain full coverage of foliage. When applying by air, apply in at least 2 gal. of water/A.</p>		

POME FRUITS – APPLE, CRABAPPLE, LOQUAT, MAYHAW, ORIENTAL PEAR, PEAR, QUINCE

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Apple Aphid Apple Maggot (Adult) Cherry Fruit Fly spp. (Adult) Coding Moth Green Fruitworm Japanese Beetle Leafhopper spp. Leafroller spp. Lesser Appleworm Omnivorous Leafroller Orange Tortrix Oriental Fruit Moth Pear Psylla Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio Rosy Apple Aphid San Jose Scale (fruit infestations only) Spirea Aphid ¹ Stink Bug spp. Tent Caterpillar spp. Tentiform Leaf Miner spp. Tufted Apple Budworm Webworm spp.	0.02-0.04 lb. a.i. (2.56-5.12 fl. oz.)	See additional instructions below. Do not apply within 21 days of harvest. Do not apply more than 0.2 lb. a.i. (1.6 pt.)/A per year. Do not apply more than 0.16 lb. a.i. (1.28 pt.)/A per year post bloom.
<p>Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds and IPM recommendations.</p> <p>Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply in at least 5 gal. of water per acre, but use higher volumes as appropriate for thorough coverage.</p>		

STONE FRUITS – APRICOT, SWEET CHERRY, TART CHERRY, NECTARINE, PEACH, PLUM, CHICKASAW PLUM, DAMSON PLUM, JAPANESE PLUM, PLUMCOT, PRUNE

Pests	Rate KENDO EC Insecticide per Acre	Remarks
American Plum Borer Apple Maggot (Adult) Black Cherry Aphid Cherry Fruit Fly spp. (Adult) Codling Moth Green Fruitworm Japanese Beetle June Beetle Leafhopper spp. Leafroller spp. Oriental Fruit Moth Peachtree Borer spp. Peach Twig Borer Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio Rose Chafer Stink Bug spp. Tent Caterpillar spp. Thrips spp.	0.02-0.04 lb. a.i. (2.56-5.12 fl. oz.)	See additional instructions below. Do not apply within 14 days of harvest. Do not apply more than 0.2 lb. a.i. (1.6 pt.)/A per year. Do not apply more than 0.16 lb. a.i. (1.28 pt.)/A per year post bloom.
<p>Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds and IPM recommendations.</p> <p>Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply at least 5 gal. of water per acre, but use higher volumes as appropriate for thorough coverage.</p>		

SUGARCANE

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Mexican Rice Borer ¹ Pygmy Mole Cricket Rice Borer ¹ Sugarcane Aphid ³ Sugarcane Beetle (Adult) ² Sugarcane Borer ¹ West Indian Crane fly Yellow Sugarcane Aphid ³	0.025-0.04 lb. a.i. (3.20-5.12 fl. oz.)	See additional instructions below. ¹ For control before the larva bores into the plant stalk. ² Suppression only of beetles active above ground. ³ See resistance statement under General Directions for Use. Do not apply within 21 days of harvest. Do not apply more than 0.16 lb. a.i. (1.28 pt./A per season.

Use scouting to determine application, usually at intervals of 7 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic threshold.

Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply at least 2 gal. of water/A.

SUNFLOWER

Target Pests	Rate KENDO EC Insecticide per Acre	Remarks
Cutworm spp. Sunflower Beetle	0.015-0.025 lb. a.i. (1.92-3.20 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only.
Banded Sunflower Moth Fall Armyworm ¹ Grasshopper spp. Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Spotted Cabbage Looper Stem Weevil (Adult) Stink Bug spp. Sunflower Maggot (Adult) Sunflower Moth Woollybear Caterpillar	0.02-0.03 lb. a.i. (2.56-3.84 fl. oz.)	² Suppression only. ³ See resistance statement under General Directions for Use. Do not apply within 45 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pt./A per season. Do not apply more than 0.09 lb. a.i. (0.72 pt./A per season after bloom initiation. Do not apply as an Ultra Low Volume (ULV) spray.
Beet Armyworm ^{1,3} Spider Mite spp. ²	0.03 lb. a.i. (3.84 fl. oz.)	

Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.

Apply by ground or air using enough water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in at least 2 gal. of water/A.

TOBACCO (AIR DRIED) – BURLEY TOBACCO AND FLUE-CURED TOBACCO

Pests	Rate KENDO EC insecticide per Acre	Remarks
Aphid spp. ^{2,3} Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cucumber Beetle spp. (Adult) Cutworm spp. Grasshopper spp. Japanese Beetle (Adult) Katydid spp. Plant Bug spp. ³ Potato Tuberworm Salt Marsh Caterpillar Stinkbug spp. Thrips spp. ² Tobacco Budworm Tobacco Flea Beetle (Adult) Tobacco Hornworm Tree Cricket spp. Vegetable Weevil (Adult) Webworm spp.	0.015-0.03 lb. a.i. (1.92-3.84 fl. oz.)	See additional instructions below. ¹ For control of first and second instars only. ² Suppression only. ³ See resistance statement under General Directions for Use. Do not apply within 40 days of harvest. Do not apply more than 0.09 lb. a.i. (0.72 pt./A per year.

Use scouting to determine application, usually at intervals of 7 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds.

Apply by ground or air using enough water to obtain full coverage of the foliage. When applying by air, apply in at least 2 gal. of water/A.

TREE NUTS – ALMOND, BEECH NUT, BRAZIL NUT, BUTTERNUT, CASHEW, CHESTNUT, CHINQUAPIN, FILBERT (HAZELNUT), HICKORY NUT, MACADAMIA NUT (BUSH NUT), WALNUT-BLACK, WALNUT-ENGLISH (PERSIAN), PECAN

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Ants Chinch Bug Coddling Moth Filbertworm Leaffooted Bug Leafroller spp. Navel Orangeworm Peach Twig Borer Plant Bug spp. Stink Bug spp. Walnut Aphid Walnut Husk Fly spp. (Adult)	0.02-0.04 lb. a.i. (2.56-5.12 fl. oz.)	See additional instructions below. Do not apply within 14 days of harvest. Do not apply more than 0.16 lb. a.i. (1.28 pt.)/A per year. Do not apply more than 0.12 lb. a.i. (0.96 pt.)/A per year post bloom.
Hickory Shuckworm Pecan Aphid spp. Pecan Casebearer spp. Pecan Phylloxera spp. Pecan Spittlebug Pecan Weevil Stink Bug spp.	0.02-0.04 lb. a.i. (2.56-5.12 fl. oz.)	
Use scouting to determine application, usually at intervals of 5 or more days. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of the foliage or target area. When applying by air, apply in at least 5 gal. of water per acre, but use higher rates as appropriate for thorough coverage.		

CONIFER AND DECIDUOUS TREES – PLANTATIONS AND NURSERIES

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Bagworm Balsam Twig Aphid Balsam Woolly Aphid Birch Leafminer Black Pine Weevil Elm Leaf Beetle European Elm Leaf Beetle Gypsy Moth Japanese Beetle June Beetle spp. Leaf Beetle spp. Leafroller spp. May Beetle spp. Mealybug spp. ¹ Pales Weevil Pine Chafer Pine Colaspis Beetle Pine Conelet Bug Pine Leaf Chermid Pine Needle Scale Pine Sawfly spp. Pine Tip Moth spp. Pine Tortoise Scale Pine Weevil spp. Poplar Aphid spp. Sawfly spp. Spittlebug spp. Spruce Budworm Tent Caterpillar spp. Tussock Moth spp. Webworm spp.	0.02-0.04 lb. a.i. (2.56-5.12 fl. oz.)	See additional instructions below. Do not apply more than 0.24 lb. a.i. (1.92 pt.)/A per year.
Use scouting to determine timing for control of exposed foliage, flower, cone, seed and bark feeding insects. Base the timing and frequency of applications on the timing when insect populations reach local economic thresholds. Apply by ground or air using enough water to obtain full coverage of target site. When applying by air, apply in at least 2 gal. of water/A.		

CONIFER AND DECIDUOUS TREES – SEED ORCHARDS

Pests	Rate KENDO EC Insecticide per Acre	Remarks
Coneworm spp. Seed Bug spp. Thrip spp.	See Remarks	For high volume sprayers, dilute 5.12 oz. per 100 gal. of water and apply 5-10 gal. of finished spray per tree. For low volume sprayers, dilute 20 fl. oz. per 100 gal. of water and apply 100 gal. of finished spray/A. For aerial applications, apply 15 fl. oz./A in a minimum of 10 gal. of finished spray/A. Do not apply more than 0.5 lb. a.i. (4 pt.)/A per year.

NON-CROPLAND ADJACENT TO AGRICULTURAL AREAS (EXCLUDING PUBLIC LAND)

Pests	Instructions
See crop instructions in sections above for specific pest and rate information	Spray non-cropland next to agricultural areas to control insects which may migrate to and threaten crops. Follow the instructions and recommendations found on this label for the adjacent crop outlet and target pests. When foliage is dense/large, insect populations are high or larval stages are large, use the highest rates on this label. Repeat as necessary to maintain control. Do not apply more than 0.2 lb. a.i. (1.6 pt.)/A per year. Do not graze livestock in treated areas.

Rate Conversion Chart

Treated Acres/Gal.	66	50	40	33	25
Pts./A	0.12	0.16	0.20	0.24	0.32
Fl. oz./A	1.92	2.56	3.20	3.84	5.12
Lb. a.i./A	0.015	0.02	0.025	0.03	0.04

NON-CROP USES - TURF AND ORNAMENTALS

KENDO EC Insecticide may be used for applications to ornamentals grown in commercial greenhouses, shade houses, and nurseries, and turf grown on sod farms or for commercial seed production.

KENDO EC Insecticide may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown, such as residential landscape areas and non-residential landscapes around institutional, public, commercial, and industrial buildings, parks, recreational areas, golf courses, and athletic fields.

KENDO EC Insecticide may also be used for applications to golf course fairways, greens, greens aprons, and tee areas.

IMPORTANT: Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

Do not apply this product through any type of irrigation system for turf and ornamental uses. Do not apply this product to edible crops or crops grown for food/feed when applied to turf or ornamentals. Do not apply this product by aerial application for turf and ornamental uses.

SPRAY DRIFT PRECAUTIONS

Observe restrictions found elsewhere on this label. Do not make applications when wind speed is 15 miles per hour or greater. Low humidity and high temperatures increase the likelihood of spray drift to sensitive areas. Avoid spraying during conditions of low humidity and/or high temperature.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when the wind direction is toward the aquatic area. Do not make outdoor applications during temperature inversions. Inversions are characterized by stable air and increasing temperature with height above ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

APPLICATION

KENDO EC Insecticide mixes easily with water and may be used in all types of application equipment. Mix product with the required amount of water and apply as a dilute application to the point of runoff. Apply product using spray nozzles which produce a coarse droplet size. Formation of very small droplets may be minimized by appropriate nozzle selection and by avoiding excessive spray pressure. For application to plants like holly, pine, or ivy which have hard-to-wet foliage, add a spreader-sticker to enhance knockdown and increase residual activity. If application is made as a concentrate or mist-type application, use the same amount of product as would be used in a dilute application.

MIXING

KENDO EC Insecticide is to be diluted with water for spray application and may be used in all types of application equipment. First fill application tank with 1/2 - 3/4 volume of water. It is suggested that the pH of the water be between 5 and 7; use a buffering agent if necessary to adjust the pH. Next slowly add KENDO EC Insecticide to the applicator tank water with maximum agitation. Finally, fill tank to desired volume and continue to agitate while making applications. If application is interrupted, agitate or re-suspend spray solution before resuming sprays. Always add KENDO EC Insecticide last if other chemicals are to be added to the applicator tank. If mixed with EC formulations or oils, use within 24 hours. Make up only amount of application volume as required. See mixing charts below.

KENDO EC Insecticide Mixing Chart for Ornamental Insect Pest Control (KENDO EC Insecticide to add per spray tank)

Desired Rate of KENDO EC Insecticide per 100 gallons	25 gallons spray tank	50 gallons spray tank	100 gallons spray tank	200 gallons spray tank	300 gallons spray tank
1.3 oz.	0.33 oz.	0.65 oz.	1.3 oz.	2.6 oz.	4.0 oz.
2.6 oz.	0.65 oz.	1.3 oz.	2.6 oz.	5.2 oz.	7.9 oz.
4.4 oz.	1.1 oz.	2.2 oz.	4.4 oz.	8.8 oz.	13.3 oz.

KENDO EC Insecticide Mixing Chart for Turf Insect Pest Control (KENDO EC Insecticide to add per 100 gallon spray tank)

Rate of KENDO EC insecticide	2 gallons	4 gallons	6 gallons	8 gallons	10 gallons
4.4 oz./A	5.0 oz.	2.5 oz.	1.7 oz.	1.2 oz.	1.0 oz.
8.8 oz./A	10.0 oz.	5.0 oz.	3.3 oz.	2.5 oz.	2.0 oz.
17.6 oz./A	20.0 oz.	10.0 oz.	6.7 oz.	5.0 oz.	4.0 oz.

Conversion Rate: 1 Fluid ounce (fl. oz.) equals 29 milliliters (mL).

COMPATIBILITY

KENDO EC Insecticide has been found to be compatible with most commonly used fungicides, miticides, liquid fertilizers, and other insecticides. Use a jar test to check physical compatibility using the correct proportion of products if local experience is unavailable.

Note: While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution. It is advised to prespray a selection of ornamental plants and observe them for 7-10 days prior to treating large areas if local use experience is unavailable.

USE DIRECTIONS

ORNAMENTALS

Agricultural: Ornamentals in Greenhouses, Shadehouses, and Nurseries

Non-Agricultural: Ornamentals (including Trees, Shrubs, Flowers, Evergreens, Foliage Plants, and Ground Covers) in Residential Landscaped Areas and Landscaped Areas Around Institutional, Public, Commercial, and Industrial Buildings, Parks, Recreational Areas, Golf Courses, and Athletic Fields

Pest	Rate of KENDO EC Insecticide Per 100 gallons	Instructions
Ants (Including imported fire ants) Aphids Armyworms Azalea caterpillars Bagworms ¹ Black vine weevils (adult) Boxelder bugs Budworms California Oakworms Cankerworms Cockroaches Crickets Cutworms Eastern tent caterpillars Elm leaf beetles European sawflies Fall webworms Flea beetles Forest tent caterpillars Gypsy moth larvae Japanese beetles (adult) June beetles (adult) Lace bugs Leaf-feeding caterpillars Leafhoppers Leafminers, (adult) Leaf rollers Leaf skeletonizers Midges Mosquitoes Oleander moth larvae Pillbugs Pine sawflies Pine shoot beetles Pinetip moths Plant bugs Root weevils Sawflies Scale insects (crawlers) ² Spiders Spittlebugs Striped beetles Striped oakworms Thrips Tip moths Tussock moth larvae Wasps	1.3-4.4 fl. oz. (38-128 mL)	Begin application to ornamentals before high insect pest populations become established. Reapply as necessary to keep pest populations under control, using higher rates as pest pressure increases. Good spray coverage is necessary to provide the most effective level of control. For ornamentals with waxy, hard-to-wet foliage, add a spreader-sticker at recommended rates to enhance the control of insects. For spot treatments, use 0.44 fl. oz. KENDO EC Insecticide per 1-2.5 gallons of water. Apply at 7- day intervals if retreatment is necessary. Do not apply more than 0.36 lb. of the active ingredient (46 fl. oz. of concentrate) per acre per year. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. ¹ Bagworm: Apply KENDO EC Insecticide when bagworm larvae begin to hatch and spray directly on the larvae. Control will be best if the larvae are young. ² Scale: Cover the plant thoroughly with KENDO EC Insecticide spray, including trunks, stems, twigs, and foliage.
Broadmites Brown softscales California redscales (crawler) Clover mites Mealybugs Pine needlescales (crawler) Spider mites Whiteflies	2.6-4.4 fl. oz. (75-128 mL)	

TURFGRASS

Agricultural: Sod Farms, Grass Seed Farms

Non-Agricultural: Lawns around Residential, Institutional, Public, Commercial, and Industrial Buildings, Parks, Recreational Areas, Golf Courses, and Athletic Fields, Golf Course and Athletic Field Turf

Pest	Amount of KENDO EC Insecticide	Instructions
Ants (including imported fire ants) Armyworms Centipedes Crickets Cutworms Earwig Fleas (adult) Grasshoppers Japanese beetles (adult) Millipedes Mites Pillbugs Sod webworms Sow bugs Ticks (including species which transmit Lyme disease)	2.9-6 ml/1,000 sq. ft. (4.4-8.8 fl. oz./A)	Begin application to turf before the establishment of high insect pest populations and before significant turf damage has occurred. Reapply as necessary to keep pest populations under control, using higher rates as pest pressure increases. Apply at 7-day intervals if retreatment is necessary. Do not apply more than 0.36 lb. of active ingredient (46 fl. oz. of concentrate) per acre per year. For spot treatments, use 0.44 fl. oz. of KENDO EC Insecticide per 1-2.5 gals. of water.
Bluegrass billbugs (adult) Black turfgrass ateniensis (adult) Chiggers Fleas (adult) Grub (suppression) Hyperodes weevils (adult) Mole crickets (nymphs and young adults)	6 ml/1,000 sq. ft. (8.8 fl. oz./A)	Do not apply when turfgrass is waterlogged or when soils are saturated with water (i.e., will not accept irrigation). Keep children and pets off treated areas until spray has dried following the application.
Chinch bugs Mole crickets (mature adults) (Not for use on mature adult mole crickets and chinch bugs in New York State)	12 ml/1,000 sq. ft. (17.6 fl. oz./A)	See additional instructions below for specific pests.

Armyworms, cutworms, fleas, and other Surface Insects: For best results, apply KENDO EC Insecticide at recommended rates in 2-5 gals. of water per 1,000 sq. ft. If high rainfall amounts are forecast, a spreader-sticker may be useful; otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12-24 hours for optimum control of surface-feeding insect pests.

Chinch bugs, billbugs, and other Thatch Inhabiting Insects: For best results apply KENDO EC Insecticide at recommended rates in 2-10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant, or similar adjuvant is recommended at label rates. Irrigate lightly after application with up to 1/2 inch of water to move the KENDO EC Insecticide into the thatch layer. If irrigation is not available, then use high water application rates for optimum results.

Mole crickets, grubs, and other Subsurface Insects: For best results apply KENDO EC Insecticide at recommended rates in 4-10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant, or similar adjuvant is strongly recommended following label rates. Use the highest water application rates possible with your sprayer. Apply KENDO EC Insecticide to turf which is wet with dew, rain, or irrigation. Water-in immediately after application with 1/4 - 1/2 inch of water for optimum results.

Fire Ants: Treat individual mounds with a drench application by means of a watering can. Use 0.32 fl. oz. of KENDO EC Insecticide per 2.5 gals. of water. Thoroughly soak each mound as well as a 3 ft. diameter circle around each mound. Apply the mixture gently to avoid disturbing the mound; disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. For best results, apply in early morning or late evening hours. Make additional treatments if necessary, but not more than every 7 days.

Mosquitoes: Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. For best results, apply KENDO EC Insecticide at recommended rates in 2-5 gals. of water per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND WARRANTY

HELM AGRO US, INC. AND SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY.

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all variables and to eliminate all risks associated with its use. Injury or damage may result because of conditions which are beyond the control of HELM AGRO US, INC. or the Seller. HELM AGRO US, INC. warrants only that this product conforms to the chemical description on the label and is believed to be reasonably fit for the purposes referred to in the Directions for Use when used as directed under normal conditions. HELM AGRO US, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no case shall HELM AGRO US, INC. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Any variation or exception from this warranty must be in writing and signed by an authorized HELM AGRO US, INC. representative.



Helm Agro US, Inc.
8295 Tournament Drive
Suite 310
Memphis, Tennessee 38125

KENDO EC Insecticide is a trademark of Helm Agro US Inc.

Rev. 10/07