



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

74530-76

Date of Issuance:

12/27/16

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Helosate 75 SG PRO Herbicide

Name and Address of Registrant (include ZIP Code):

Helm Agro US, Inc.
 401 E. Jackson Street, Suite 1400
 Tampa, FL 33602

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Reuben Baris, Product Manager 25
 Herbicide Branch, Registration Division (7505P)

Date:

12/27/16

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 74530-76.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 8/18/2016
-

If you have any questions, please contact Emily Schmid by phone at 703-347-0189, or via email at schmid.emily@epa.gov.

Enclosure

GROUP 9 HERBICIDE

HELOSATE 75 SG PRO

HERBICIDE

For Use as a non-selective professional herbicide for use in non-crop areas and industrial sites.
Avoid herbicide contact with foliage, green stems, exposed non-woody roots or fruit of crops,
desirable plants and trees because severe injury or destruction may result.

INGREDIENTS

ACTIVE INGREDIENT:

*Glyphosate (n-(phosphonomethyl) glycine), in the form of its ammonium salt..... 75.7%

OTHER INGREDIENTS:..... 24.3%

TOTAL..... 100.0%

*Equivalent to 68.9% of glyphosate acid

EPA Reg. No. 74530-xx

EPA Est. No. xx-xx-xx

Net Contents: _____

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

ACCEPTED
12/27/2016
Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 74530-76

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed	<ul style="list-style-type: none">· Call a poison control center or doctor immediately for treatment advice.· Have person sip a glass of water if able to swallow.· Do not induce vomiting unless told to do so by a poison control center or doctor.· Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none">· Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.· Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.· Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the usage of gastric lavage.	

Manufactured by: **HELM AGRO US, Inc.**, 401 E. Jackson St., Suite 1400, Tampa, FL 33602

IMPORTANT PHONE NUMBER

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
KEEP OUT OF REACH OF CHILDREN
DANGER**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before re-use.

Domestic animals: Keep livestock and pets out of treated areas. Do not graze livestock on treated areas. This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long sleeved shirt and long pants
- shoes plus socks
- protective eyewear

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters and rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area at the time of application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Open dumping is prohibited. Store in original container only. Keep containers closed when not in use. Separate pesticides during storage to prevent cross-contamination of other pesticides, fertilizers, food, and feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed must be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

CONTAINER HANDLING:

Nonrefillable Containers: Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

Nonrefillable Plastic: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning; if burned, stay out of smoke.

Nonrefillable Paper or Plastic Bags: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRODUCT INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT

Product Description: This product is a systemic herbicide with no residual soil activity that should be used postemergent. It offers broad-spectrum non-selective control of many annual weeds, perennial weeds, woody brush and trees. This product is formulated as a water-soluble granule containing surfactant and no additional surfactant is required. However, the addition of a nonionic surfactant at 0.5% solution may improve weed control. May be applied through most standard sprayers after dissolution and thorough mixing with water according to label instructions.

Time to Symptoms: This product moves systemically through the plant from the point of foliage contact and into the root system. Visible effects on most annual weeds occur within 2 to 4 days. Visible effects on most perennial weeds may not

occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visible effects of control. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

No Soil Activity: Target weeds must be emerged at the time of application to be controlled by this product. Weeds emerging from seed after application will not be controlled. Plants growing from unattached underground rhizomes or root stocks of perennials that have not yet emerged at the time of application will not be affected by the herbicide and will continue to grow.

Cultural Considerations: Control may be reduced when applications are made to annual and perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment. Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Rainfastness: Rainfall or irrigation occurring within 2-4 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2-4 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

Spray Coverage: Uniform and complete spray coverage will provide best results. Do not spray weed foliage to the point of runoff. Always use the higher rate of this product per acre within the specified range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Annual Maximum Use Rate: The maximum allowed application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or as tank mixtures. For non-crop uses the combined total of all glyphosate treatments must not exceed 11.61 pounds of this product (8 pounds of glyphosate acid) per acre per year.

ATTENTION: TO PREVENT INJURY TO DESIRABLE PLANTS, EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, EXPOSED NON-WOODY ROOTS, OR DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT. AVOID DRIFT.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation. Herbicide solutions containing minute quantities of this product can cause severe damage or destruction to desirable plants or other areas on which treatment was not intended. Injury potential to desirable plants from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, do not combine of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **DO NOT APPLY AT EXCESSIVE SPEED OR PRESSURE**

NOTE: Use of this product in any manner not consistent with this label may result in injury to animals, crops, desirable plants, persons or other unintended consequences. To prevent spills and contamination, keep container closed.

MIXING INSTRUCTIONS

Mix only the amount of solution to be used during a 24 hour period in a clean sprayer. Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED WEED CONTROL MAY OCCUR IF WATER CONTAINING SOIL SEDIMENT IS USED AS A CARRIER. DO NOT MIX THIS PRODUCT WITH WATER FROM PONDS AND DITCHES THAT IS VISIBLY MUDDY OR MURKY.

Use caution to avoid siphoning back into the carrier source. An approved anti-back-siphoning device must be used where required by state or local regulations. Foaming of the spray solution may occur during mixing and application. To minimize or reduce foaming, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

Mix, store, and apply spray solutions of this product using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product does not provide residual weed control. HELOSATE 75 SG PRO Herbicide may be tank-mixed with other herbicides to provide residual weed control, a broader weed control spectrum or an alternate mode of action. Observe all precautions and limitations on the label, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines of all individual product labels when tank mixing. Use the most restrictive precautionary statements for each product in the tank mixture.

Always determine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Buyer and all users are responsible for any and all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly specified on this label. Performance may be reduced when mixing this product with herbicides or other materials not specified on this label.

Procedure for Preparing Spray Solution

Mixing with Water

Use the following procedure to mix this product in water. Fill the mixing or spray tank with the required amount of water. Add the specified amount of this product (see the DIRECTIONS FOR USE and WEEDS CONTROLLED sections of this label) near the end of the filling process and mix well.

Tank Mixture Instructions

Use the following procedure to mix this product when preparing tank mixtures with other labeled products.

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. Add HELOSATE 75 SG PRO Herbicide.
4. Add tankmix partners as follows:
 - a: Wettable powder - make a slurry with the water carrier, and add it SLOWLY through the screen into the tank.
 - b: Flowable formulation - premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank.
 - c: Emulsifiable concentrate formulation - premix one part emulsifiable concentrate with two parts water. Add

- diluted mixture slowly through the screen into the tank.
5. Continue agitation throughout process.
 6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
 7. Where nonionic surfactant is used, add this to the spray tank before completing the filling process.
 8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

When tank mixing this product with other products, maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

Always determine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Mixing for Hand-Held Sprayers

Prepare the required volume of spray solution by adding the amount of HELOSATE 75 SG PRO Herbicide as shown in the table below to a clean, empty sprayer. Add the appropriate amount of water and agitate to ensure dissolution of product. For best results when using backpack sprayers, mix the labeled amount of this product with the specified volume of water in a larger container. Fill sprayer with the mixed solution

Amount of HELOSATE 75 SG PRO Herbicide			
Mix Volume	Annuals	Perennials and Brush	Low Volume Directed
1 Gallon	1.2 oz.	1.5 oz.	4.0 - 8.0 oz.
3 Gallons	3.6 oz.	4.5 oz.	12.0 oz. - 1.5 lbs.
10 Gallons	12.0 oz.	15.0 oz.	2.5 lbs. - 5.0 lbs.

Colorants or Dyes

Commercially approved colorants or marking dyes may be added to this product. However, colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's directions.

Drift Control Additives

Drift control additives may be used with all equipment types. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Application Equipment and Techniques

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is

responsible for considering all these factors when making decisions.

RESTRICTIONS: DO NOT apply this product by air. DO NOT apply this product through any type of irrigation system. DO NOT allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to desirable plants or other areas on which treatment was not intended.

APPLY SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Ground Broadcast Equipment

For control of annual or perennial weeds listed on this label using broadcast equipment - Use the specified rates of this product in 10 to 80 gallons of water per acre as a broadcast spray unless otherwise specified on this label. As density of weeds increases, spray volume should be increased within the specified range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. With ground application equipment, use flat fan nozzles for best results. Check for even distribution of spray droplets.

Hand-Held and High-Volume Equipment

Use coarse sprays only.

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed in the Annual Weeds section of the WEEDS CONTROLLED section, apply 1.2 ounces of this product per 1 gallon of spray solution. See table in Mixing for Hand-Held Sprayers section for larger mixing volumes.

For control of perennials such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle, use 1.5 ounces of this product per 1 gallon of spray solution on harder-to-control perennials. See table in Mixing for Hand-Held Sprayers section for larger mixing volumes.

For directed low-volume spray applications, use 4.0 to 8.0 ounces of this product per 1 gallon of spray solution for control or partial control of brush weeds. See table in Mixing for Hand-Held Sprayers section for larger mixing volumes. Ensure spray coverage is uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of brush and tree seedlings when foliage is thick and dense, or where there are multiple sprouts

CDA (Controlled Droplet Application) Equipment

The rate of this product applied per acre by CDA equipment must not be less than the amount stated in this label when applied by conventional broadcast equipment.

For hand-held CDA units, apply a solution of 1.5 to 2.0 pounds of this product in one gallon of water at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mile per hour. For vehicle mounted CDA equipment, apply 3 to 15 gallons of water per acre.

Spray patterns produced by CDA equipment is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

SITE AND USE INSTRUCTIONS

Applications may be made to control any weeds listed in the annual, perennial and brush weeds and tree seedlings tables unless otherwise specified.

Non-crop Areas and Industrial Sites

This product may be used in non-crop areas including:

Airports		Industrial sites		Recreational areas
Apartment complexes		Landscape areas		Residential areas
Commercial sites		Lumberyards		Rights-of-way
Conservation Reserve Program (CRP)		Manufacturing sites		Roadsides
Ditch banks		Municipal sites		Schools
Driveways		Natural areas		Sports complexes
Dry ditches		Office complexes		Storage areas
Dry canals		Ornamental landscapes		Warehouse areas
Fencerows		Parks		Wildlife management areas
Golf courses		Parking areas		

RESTRICTIONS: DO NOT use this product on crops, timber, other plants being grown for sale, other commercial use, or for commercial seed production. DO NOT use this product for research purposes. Keep people and pets off treated areas until spray solution has dried.

Bare Ground, Trim-and-Edge and Weed Control in Non-Crop Areas

This product may be used for general weed control in non-crop areas. Apply with application equipment described in this label. May be used for spot treatment of unwanted vegetation, as trim-and-edge around objects in non-crop sites and to eliminate unwanted weeds growing in established ornamental plantings or shrub beds. May be used prior to planting an area to flowers, ornamentals, flowers, turfgrass (sod or seed), prior to laying asphalt or beginning construction projects. DO NOT use on plants grown for sale or other commercial use, or for commercial seed production. Applications of this product may be repeated for weed control as needed to maintain bare ground.

If tank mixed with a generic active ingredient (2,4-D, pendimethalin, etc.) described in this label, the user is responsible for ensuring that the specific application being made is included on the label of the specific product being used in the tank-mixture.

HELOSATE 75 SG PRO Herbicide may be tank-mixed with the following products. Refer to tank mix products' label for approved non-crop sites and application rates.

2,4-D
Banvel
Barricade 65WG
Certainty
Dimension 4 EC
Endurance
Pendimethalin
Pendulum 3.3EC
Pendulum WDG
Ronstar 50 WP
Surflan

When tank mixes are applied bare ground, HELOSATE 75 SG PRO Herbicide provides control of the emerged annual weeds and control or partial control of emerged perennial weeds.

Dormant Turfgrass

This product may be used to control or suppress numerous winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Application must be made prior to spring greenup only when turf is dormant.

RESTRICTION: DO NOT use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

RESTRICTIONS: Apply 5 to 16 ounces of this product per acre in 10 to 80 gallons of water per acre. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Application rates in excess of 9 ounces per acre may result in injury or delayed greenup in highly maintained areas, golf courses and lawns.

Turfgrass Renovation (Except for Commercial Sod Farms)

This product will control most existing vegetation prior to renovating turfgrass areas. To maximize control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. If repeat treatments are necessary, sufficient regrowth must be attained prior to application. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Desirable turfgrasses may be planted following the above procedures.

In existing turfgrass, hand-held equipment may be used for spot treatment of unwanted vegetation growing.

RESTRICTIONS: DO NOT use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. DO NOT use this product for renovation of bermudagrass or kikuyugrass sods. DO NOT disturb soil or underground plant parts before treatment. Delay tillage or renovation techniques such as vertical mowing, coring or slicing for 7 days after application to allow translocation into underground plant parts. DO NOT feed or graze treated turfgrass or feed treated thatch to livestock.

Parks, Recreational and Residential Areas

HELOSATE 75 SG PRO Herbicide may be used in parks, recreational and residential areas. Apply with any application equipment described in this label. Use this product to trim-and-edge around buildings, around trees, fences, other objects, paths and sidewalks in these areas. May be used for spot treatment of unwanted vegetation or to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. May be used prior to planting an area to flowers, ornamentals, turfgrass (sod or seed), prior to laying asphalt or beginning construction projects.

All of the instructions in the Non-Crop Areas and Industrial Sites section apply to park and recreational areas.

RESTRICTION: DO NOT use around plants being grown for sale or other commercial use.

WEEDS CONTROLLED

When weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area, ALWAYS use the higher rate of this product per acre within the range. If weeds are heavily covered with dust when treated, reduced results may occur. If weeds have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the label sections below for rates to control annual and perennial weeds. For difficult to control perennial weeds and where plants are growing under stressed conditions, or where infestations are dense, this product may be used at up to 11.61 pounds per acre for enhanced results.

Annual Weeds

Use 2.25 to 4.5 pounds per acre of this product as a broadcast spray to control annual weeds.

For spray-to-wet applications, apply 1.2 ounces of this product per 1 gallon of spray solution.

Anoda, spurred	Fleabane, annual*	Ragweed, giant
Barley*	Fleabane, hairy (<i>Conyza bonariensis</i>)*	Red rice
Barnyardgrass*	Fleabane, rough*	Russian thistle
Bassia, fivehook	Florida pusley	Rye*
Bittercress*	Foxtail*	Ryegrass*
Black nightshade*	Goatgrass, jointed*	Sandbur, field*
Bluegrass, annual*	Goosegrass	Shattercane*
Bluegrass, bulbous*	Grain sorghum (milo)*	Shepherd's-purse*
Brome, downy*	Groundsel, common*	Sicklepod
Brome, Japanese*	Hemp sesbania	Signalgrass, broadleaf*
Browntop panicum*	Henbit	Smartweed, ladysthumb*
Buttercup*	Horseweed/Marestail (<i>Conyza canadensis</i>)	Smartweed, Pennsylvania*
Carolina foxtail*	Itchgrass*	Sowthistle, annual
Carolina geranium	Johnsongrass, seedling	Spanishneedles
Castor bean	Junglerice	Speedwell, purslane*
Cheatgrass*	Knotweed	Sprangletop*
Cheeseweed	Kochia	Spurge, annual
(<i>Malva parviflora</i>)	Lambsquarters*	Spurge, prostrate*
Chervil*	Little barley*	Spurge, spotted*
Chickweed*	London rocket*	Spurry, umbrella*
Cocklebur*	Mayweed	Starthistle, yellow
Copperleaf, hophornbeam	Medusahead*	Stinkgrass*
Corn*	Morningglory	Sunflower*
Corn speedwell*	(<i>Ipomoea</i> spp.)	Teaweed/Prickly sida
Crabgrass*	Mustard, blue*	Texas panicum*
Dwarfdandelion*	Mustard, tansy*	Velvetleaf
Eastern mannagrass*	Mustard, tumble*	Virginia copperleaf
Eclipta*	Mustard, wild*	Virginia pepperweed*
Fall panicum*	Oats	Wheat*
Falsedandelion*	Pigweed*	Wild oats*
Falseflax, smallseed*	Plains/Tickseed coreopsis*	Witchgrass*
Fiddleneck	Prickly lettuce*	Woolly cupgrass*
Field pennycress*	Purslane, common	Yellow rocket
Filaree	Ragweed, common*	

* These species will be controlled or partially controlled when using field broadcast equipment (boom sprayers using flat-fan nozzles) in applications made using 10 to 80 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

Perennial Weeds

When perennial weeds are treated, best results are obtained when they are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). Best results are obtained for non-flowering plants when the plants reach a mature stage of growth. However, in many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the range.

Apply this product as a broadcast spray at 4.5 to 9.0 pounds per acre to control perennial weeds.

Apply 1.5 ounces of this product per 1 gallon of spray solution for spray-to-wet applications. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment.

Apply 4.0 to 8.0 ounces of this product per 1 gallon of spray solution when using hand-held equipment for low-volume directed spot treatments.

Allow 7 or more days after application before tillage.

Alfalfa*	Nutsedge; purple, yellow	Dock, curly
Johnsongrass	Bluegrass, Kentucky	Spurge, leafy*
Alligatorweed*	Orchardgrass	Dogbane, hemp
Kikuyugrass*	Blueweed, Texas	Thistle, artichoke
Anise (fennel)	Pampasgrass	Fescue
Knapweed	Bromegrass, smooth	Thistle, Canada
Bahiagrass	Paragrass	Fescue, tall
Lantana	Bursage, woolly-leaf	Timothy
Beachgrass, European	Pepperweed, perennial	German ivy
Lespedeza	Canarygrass, reed	Torpedograss*
(Ammophila arenaria)	Phragmites*	Guineagrass
Milkweed, common	Cattail	Trumpetcreeper*
Bentgrass*	Poison hemlock	Horsenettle
Muhly, wirestem	Clover; red, white*	Vaseygrass
Bermudagrass*	Quackgrass	Horseradish
Mullein, common	Cogongrass	Velvetgrass
Bermudagrass, water	Redvine*	Iceplant
Napiergrass	Dallisgrass	Wheatgrass, western
(knotgrass)	Reed, giant	Jerusalem artichoke
Nightshade, silverleaf	Dandelion	
Bindweed, field	Ryegrass, perennial	

*Partial Control

Brush Weeds and Tree Seedlings

When brush weeds and tree seedlings are treated, best results are obtained when they are in the seedling stage of growth. Retreatment is often required on larger plants. Under these conditions, use the higher application rate within the range.

Apply as a broadcast spray at 9.0 pounds per acre of this product to control brush weeds.

For spray-to-wet applications, apply 1.5 ounces of this product per 1 gallon of spray solution. Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment

When using hand-held equipment for low-volume directed spot treatments, apply 4.0 to 8.0 ounces of this product per 1 gallon of spray solution.

Allow 7 or more days after application before tillage.

Alder	Poison ivy*	Maple, red
Oak, southern red	black, pin	smooth, sugarbush,
Ash*	Poison oak*	Maple, sugar
Oak, white*	Dogwood*	winged*
Beech*	Poplar, yellow*	Oak, black*
Peppertree,	Elderberry	Sweetgum
Birch	Redbud, eastern	Oak, northern pin
Brazilian	Elm*	Vine maple*
Blackberry	Rose, multiflora	Oak, post
(Florida holly)*	Honeysuckle	Virginia creeper
Blackgum	Saltcedar*	Oak, red
Pine	Locust, black*	Waxmyrtle, southern*
Cherry; bitter,	Sumac; laurel, poison,	Oak, scrub

*Partial Control

CONDITIONS OF SALE AND WARRANTY

HELM AGRO US, INC. AND SELLER OFFER THIS PRODUCT AND THE BUYER AND USER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED CONDITIONS OF SALE AND WARRANTY.

The directions for use of this product are believed to be reliable and must be followed carefully. However, it is impossible to take into account all variables and to eliminate all risks associated with its use. Injury or damage may result because of conditions which are beyond the control of HELM AGRO US, INC. or the Seller. HELM AGRO US, INC. warrants only that this product conforms to the chemical description on the label and is believed to be reasonably fit for the purposes referred to in the Directions for Use when used as directed under normal conditions. To the extent consistent with applicable law, HELM AGRO US, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. To the extent consistent with applicable law, in no case shall HELM AGRO US, INC. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Any variation or exception from this warranty must be in writing and signed by an authorized HELM AGRO US, INC. representative.



HELM AGRO US, Inc.
401 E. Jackson St., Suite 1400

Tampa, FL 33602
info@helmagro.com
Tel. 813-621-8846