

74779-6

06/27/2007

5811299

1/21



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
74779-6

Date of Issuance:
JUN 27 2007

NOTICE OF PESTICIDE:
 Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance: conditional

Name of Pesticide Product:
Acephate Systemic Tree and Ornamental Insecticide

Name and Address of Registrant (include ZIP Code):
 Rainbow Treecare Scientific Advancements
 2239 Edgewood Ave. S.
 Minneapolis, MN 55426

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(2)(B) when the Agency requires all registrants of similar products to submit such data.
2. Make the following label corrections and submit one copy of the revised finished printed label incorporating these corrections before the product is released for shipment.
 - a. Add the following designation: EPA Reg. No. 74779-6.
 - b. Add "all" between "and" and "applicators" in the last sentence on the bottom of the second page so that it reads as follows: "In addition, all mixers and loaders and all applicators ___."
 - c. Correct "Niosh" to NIOSH in the PPE on the top of the third page.
 - d. Correct the text "Remove clothing immediately" in the second bullet of the User Safety Recommendations box to "Remove clothing/PPE immediately".
 - e. Add "Spray Drift Management" as the heading for the spray drift statements on the fourth page.

Signature of Approving Official:

Date:

JUN 27 2007

f. Replace the term "Recommendations" appearing in the use directions tables for the various use sites with more appropriate wording such as "Directions".

g. Add the appropriate minimum spray interval (3 days for rates up to and including 0.5 lb ai/A and 7 days for rates higher than 0.5 lb ai/A) wherever intervals are lacking throughout the Container Grown Nursery Stock and the Ornamental Trees and Shrubs use tables.

h. Add harvester ants to the list of excluded ants for the foliar spray application in the Non-Bearing Citrus table so the ant claim reads as follows: "ants (excluding Fire, Carpenter, Harvester and Pharoah)."


i. Correct the claim for ants in the Outdoor and Perimeter Spray table from "Ants (excluding imported fire ants and harvester ants)" to "Ants (excluding imported fire, carpenter, pharaoh and harvester ants)".

j. Correct the minimum spray interval specified in the Outdoor Floral Crops and Ground Covers table for aphids, thrips and lygus from 3 days to that specified below. This is needed because the maximum application rate specified for these insects on a per acre basis exceeds 0.5 lb ai/A.

The minimum spray interval is 3 days for application rates up to and including 0.5 lb ai/A and 7 days for rates exceeding 0.5 lb ai/A.

3. Submit your corrected application form as specified in item number 20 of our May 23, 2007 letter before you release the product for shipment.

A stamped copy of the label is enclosed for your records.


Marilyn A. Mautz
Biologist
Insecticide-Rodenticide Branch
Registration Division (7504P)

3/21

Acephate Systemic Tree & Ornamental Insecticide

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See Side/Back Panel for Additional Precautionary Statements, First Aid and Directions for Use

Acephate Systemic Tree & Ornamental Insecticide is absorbed through the plant root system and then moves throughout the plant, protecting it from the damage caused by sucking insects (aphids, lace bugs, leafhoppers, and thrips) and certain chewing insects (leafminers, leaf beetles, leaf tiers, and caterpillars) as listed.

Active Ingredient	
Acephate (O,S-Dimethyl acetylphosphoramidothioate).....	97.4%
<u>Other Ingredients...</u>	<u>2.6%</u>
Total	100%

EPA Reg. No. 74779- EPA Est. No.

Net Contents: X lbs (Y kg)

Rainbow Treecare Scientific Advancements
2239 Edgewood Ave. South
Minneapolis, MN 55426
1-877-ARBORIST
1-877-272-6747
www.rainbowscivance.com

ACCEPTED
with COMMENTS
in EPA Letter Dated:
JUN 27 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
As amended, for the pesticide
Registered under EPA Reg. No.
74779-6

FIRST AID: Acephate is an organophosphate, cholinesterase inhibitor	
IF SWALLOWED	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or doctor immediately for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call poison control center or doctor immediately for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call poison control center or doctor immediately for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOT LINE NUMBER	
For 24 hour medical emergency assistance (human or animal), or chemical emergency assistance (spill, leak or accident). Call CHEMTREC at 1-800-424-9300.	
NOTES TO PHYSICIAN	
This material contains a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure but decisions regarding treatment will usually need to be made before test results are available. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PROTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone.	

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are those made of any waterproof material. If you want more options, follow the instruction for category A on an EPA chemical-resistant category selection sheet.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

In addition, all mixers and loaders and applicators using low pressure hand wand application

equipment must wear:

- A Niosh-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this products concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

<p>USER SAFETY RECOMMENDATIONS:</p> <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
--

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean water high mark. Do not contaminate water when cleaning equipment or disposing of wastes. Cover or soil-incorporate spills.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

For use on turf, use limited to sod farms and golf courses, except when applying to mound or spot treatment for fire ant and harvester ant control.

Do not apply using low pressure hand wand except for control of fire ants as a mound treatment and when used on ornamental trees, shrubs, and floral plants grown for non-agricultural or non-commercial use.

Not for indoor residential use. For greenhouse use, use is limited to commercial greenhouses for use on ornamental, floral and foliage plants.

Do not apply by air.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

A variety of factors including weather conditions, (e.g., wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

- All application equipment must be properly maintained and calibrated using water as carrier. Do not apply this product as an ultralow (ULV) spray, or in any carrier other than water.
- Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Do not apply at wind speeds greater than 10 mph at the application site.
- Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage.
- For airblast applications, direct spray above foliage and turn off outward pointing nozzles at row ends and outer rows.
- Do not apply at heights greater than 4 feet.

GENERAL MIXING INSTRUCTIONS

Thoroughly clean all application components prior to mixing. Add approximately 1/2 of the required amount of water to the application tank and begin agitation. Add the required amount of Acephate Systemic Tree & Shrub Insecticide and mix thoroughly. Then add other additives and the remaining water. Maintain agitation during filling and application to ensure uniform mixture.

7/21

For application equipment that has minimal agitation, proper attention to mixing this product should be given. With such equipment, this product should be premixed with water to form a slurry prior to putting the product into the applicator. If premixing is not done, then adequate time should be allowed for the product to dissolve in the water prior to beginning applications.

SMALL VOLUME DILUTION CHART

Amount of product per 100 gallons of water	Amount of product per 25 gallons of water	Amount of product per 1 gallon of water
¼ lb (113.4 g)	1 oz (28.3 g)	1/3 tsp (1.1 g)
½ lb (226.8 g)	2 oz (56.7 g)	2/3 tsp (2.3 g)
¾ lb (340.2 g)	3 oz (85.0 g)	1 tsp (3.4 g)
1 lb (453.6 g)	4 oz (113.4 g)	1 1/3 tsp (4.5 g)

NON-BEARING CITRUS

DO NOT ALLOW LIVESTOCK TO GRAZE TREATED AREAS.

PLANTS	INSECTS	AMOUNT OF PRODUCT PER ACRE	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Non-Bearing Citrus (including citrus nursery seedlings)	Aphids Grasshoppers Katydid Mealybugs Orangedogs Plant Bugs (Lygus) Thrips Whiteflies	½ pound (8 oz) (226.8 g)	Apply for control of existing populations. The minimum spray interval for repeat applications is 3 days.	Apply this product in 100 to 200 gallons of water per acre. Spray individual juvenile or non-bearing trees for coverage with total application not to exceed specified rate in pounds per acre. Length of residual activity will depend upon spray coverage and the amount of moisture following application.
	Citrus Blackfly*	½ to ¾ pound (8-12 oz) (226.8 to 340.2 g)	The high rate should be used when a heavy infestation of Citrus Blackfly is present. Begin applications when eggs or adults first appear. The minimum spray interval for repeat applications is 7 days.	Do not allow livestock to graze treated areas. DO NOT HARVEST citrus for one year after treatment.
	Ants (excluding Fire, Carpenter, and Pharaoh*)	Foliar Spray ¾ lb (12 oz) (340.2 g)	Apply for control of existing	Applications made under prolonged hot and dry

	Ants (excluding Carpenter and Pharaoh)*	Mound Treatment – Drench Method: Mix ¾ ounce in 5 gallons of water. Apply 1 gallon of mix to each mound area by sprinkling mound until it is wet and treat a four (4) foot diameter circle around the mound.	populations	conditions may be ineffective due to the location of the ants deep within the nest.
--	---	--	-------------	---

*Not registered in CA for control of Citrus Blackfly and Harvester Ants.

NURSERY STOCK (NON-BEARING DECIDUOUS FRUIT TREES, NUT TREES AND VINES IN NURSERY FIELDS OR NON-BEARING ORCHARDS.)

PLANTS	INSECTS	AMOUNT OF PRODUCT PER ACRE	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Nuresery Stock (Non-bearing Deciduous Fruit Trees, Nut Trees, Vines) Almond Apricot Apple Cherry Kiwi Pear Pistachio Plum Prune Walnut	Aphids Plant Bugs (Lygus) Thrips	½ to 1 lb (8 – 16 oz) (226.8 to 453.6 g)	Apply as needed for control of existing populations. The minimum spray interval for repeat applications is 7 days.	Dilute in 50 to 200 gallons of water per acre and spray using ground equipment.
USE PRECAUTIONS: For application only to non-bearing trees and vines in nursery fields or in non-bearing orchards. Do not apply to interplants or single trees in a bearing orchard. Do not harvest any fruit from sprayed trees for one-year after application. Do not allow livestock to graze treated areas.				

9/21

SPECIALTY USES – NON CROP AREAS

PLANTS	INSECTS	AMOUNT OF PRODUCT PER ACRE	APPLICATION RECOMMENDATIONS
Non Crop Areas (Field borders, fencerows, roadsides, ditchbanks, and borrow pits)	Grasshoppers	¼ lb (4 oz) (113.4 g)	<p>For early to mid-season application to grasshoppers in field borders, fencerows, roadsides, ditchbanks and borrow pits</p> <p>Apply in 10 to 20 gallons of water per acre and spray using ground equipment. Use the higher volumes when spraying heavier foliage.</p> <p>Do not graze or feed vegetation cut from treated areas.</p>
	Fire Ants and Harvester Ants*	Use mound treatment-Drench Method: Mix ¾ ounce in 5 gallons of water. Apply 1 gallon of mix to each mound area by sprinkling mound until it is wet and treat a four (4) foot diameter circle around the mound.	<p>For best results, apply in the early morning or late afternoon when the ants are most active.</p> <p>Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest.</p> <p>Grass in treated area may be injured.</p> <p>Do not treat mound more than once per season.</p>

*Not registered in CA for control of Harvester Ants.

10/21

SPECIALTY USES – CONTAINER GROWN NURSERY STOCK (Arborvitae, Azalea, Camellia, Rhododendron, Roses, Viburnum, and Yew)

PLANTS	INSECTS	AMOUNT OF PRODUCT PER ACRE	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Container Grown Nursery Stock Arborvitae Azalea Camellia Rhododendron Roses Viburnum Yew	Black Vine Weevil, Strawberry Root Weevil	¾ lb (12 oz) (334.2 g)	Application should be made by mid-September for greenhouse stock and by mid-October for outdoor stock. Consult your local county extension agent for information on the identification and control of root weevils on ornamentals. The minimum spray interval for repeat applications is 7 days.	Use the specified amount per 100 gallons of water and thoroughly drench the root system.
	Ants (excluding fire, harvester, carpenter, and pharaoh)	¾ lb (12 oz) (334.2 g)	Apply as needed to control the pest.	

11/21

ORNAMENTAL TREES AND SHRUBS – FOLIAR SPRAY – Maximum single application rate is 1.0 lb active ingredient per acre.

PLANTS	INSECTS	AMOUNT OF PRODUCT PER 100 GALLONS	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Ornamental Trees and Shrubs (Except Flowering Crabapple)	Aphids Bagworms Birch Leafminer Lace Bugs Leafrollers Tent Caterpillars	¼ lb (4 oz) (113.4 g)	Begin treatments when insects first appear. The minimum spray intervals for repeat applications of 3 days up to and including 0.5 lb ai/acre and 7 days for rates exceeding 0.5 lb.ai/acre.	Apply as a full coverage spray using a hydraulic sprayer. Rainbow Treecare does not recommend applications be made to Huckleberry, Balm of Gilead, Cottonwood, Lombardy Poplar, and Viburnum suspensum. For nursery crops, spray only a few plants and observe for 2 weeks and evaluate for phytotoxicity before treating the entire crop.
	Douglas Fir Tussock Moth Larvae Gypsy Moth Larvae* Webworms	½ lb (8 oz) (226.8 g)		
	Scale (Crawlers)	½ lb (8oz) (226.8g)	Begin treatments when crawlers first appear. Repeat applications every 2 weeks as needed to achieve control.	The use of a sticker improves control of gypsy moth larvae. *Mist blower application: Adjust rates to 1 lb per 100 gallons of water for Gypsy Moth control. Adjust rate to 3/4lb per 100 gallons of water for tent caterpillar control.
	Ponderosa Pine Needle Miner	½ lb (8oz) (226.8g)	Time of application is important. Consult your County Extension Agent for proper timing.	Apply as a full coverage spray using a hydraulic sprayer.
	California Oakworm Cankerworms (Spring and Fall)	¼ lb to ½ lb (4 to 8 oz) (113.4 to 226.8g)	Begin treatments when insects first appear. Use the higher rate when larger larvae are present.	
	Nantucket Pine Tip Moth Larvae	¾ lb (12 oz) (340.2 g)	Time of application is important. Consult your County Extension Agent for proper timing. Repeat applications will be necessary for subsequent generations.	Apply as a full coverage spray using a hydraulic sprayer.

12/21

	Root Weevil Adults	¾ lb (12 oz) (340.2 g)	Begin treatments when feeding damage first appears. Repeat applications every four weeks until the first heavy frost may be necessary for complete foliage protection.	Apply as a full coverage spray using a hydraulic sprayer.
	Box Elder Bugs Budworms Leafhoppers Sawflies	¾ lb (12 oz) (340.2 g)	Begin treatments when insects first appear.	Apply as a full coverage spray using a hydraulic sprayer.
	Japanese Beetles Elm Leaf Beetles	1 lb (16 oz) (453.6 g)	Begin treatments when insects first appear. Repeat applications every 2 weeks as needed.	
Douglas Fir (Christmas Trees)	Douglas Fir Needle Midge	½ lb (8 oz) (226.8 g)	Application should be made no more than 2 weeks prior to bud burst. For additional pest management information, consult your County Extension Agent	Apply by ground as a full coverage spray in 100 gallons of water using a hydraulic sprayer.
Flowering Crabapples	Aphids Leafrollers Tent Caterpillars	¼ lb (4 oz) (113.4 g)	Begin treatments as the insects begin to appear	Apply as a full coverage spray using a hydraulic sprayer. Do not apply more often than 3 times in a growing season at a four (4) week interval Caution: Phytotoxicity has occurred on the following crabapple varieties: Hopa, Ichonoski, Malusfloribunda, Pink Perfection, Red Wine, and Snow Cloud.
Crape Myrtle	Aphids	2 – 3 level	As aphids begin to	Make a paint-on slurry by

13/21

		tablespoons (0.6 to 0.9 oz) (17 to 25.5 g) per 1 tablespoon of water	appear	mixing. Remove the loose bark from the trunk areas to be treated. Completely paint a band around each trunk to a width twice its diameter. Application should be made to trunks 6 to 8 inches above the ground and below the point where branching begins. For multi-trunk plants, treatments should be made as low as possible within the 6 to 12 inches of the ground.
--	--	--	--------	--

GOLF COURSES AND SOD FARM TURFGRASS

- Do not allow livestock to graze treated areas
- Do not feed treated grass to livestock.
- Do not apply by air
- Do not apply more than 4 1/8 lbs product/acre (4.0 lbs ai/acre) to golf courses in a single application.
- Do not apply more than 3.1 lbs product/acre (3.0 lbs ai/acre) to sod farm turfgrass in a single application.
- The minimum spray intervals for repeat applications is 3 days for application rates up to and including 0.5 lb ai/acre and 7 days for rates exceeding 0.5 lb ai/acre.
- A 3-day pre-harvest (PHI) for harvesting sod.

PLANTS	INSECTS	AMOUNT OF PRODUCT PER 1000 Sq. Ft.	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Turf (Golf Course and Sod Farm Turfgrass)	Fall Armyworm Yellow Striped Armyworm Southern Armyworm	0.4 to 0.9 oz (17 to 25.5 g)	Begin treatments when insects first appear. Repeat applications every 2 weeks as needed.	Dilute in 5 gallons of water to treat 1000 sq. ft.
	Cutworm	0.9 to 1.8 oz (25.5 to 51 g)		
	Chinch Bugs	0.9 to 1.8 oz (25.5 to 51 g)	Apply every 10 to 14 days as needed for knockdown of adult populations.	Dilute recommended amount of product in 1 to 15 gallons of water to treat 1000 sq. ft.

	Sod Webworm (Crambus spp.)	0.4 to 0.8 oz (17 to 34 g)	Begin applications when larvae first appear. Use the higher rate when quick knockdown is needed or with heavy populations. Reapply every 7 days as needed. Do not repeat at more than 1 week intervals.	
	Leafhopper	¾ oz (21.3 g)		
	Mole Crickets* Spittlebug*	0.8 to 1.4 oz (22.7 to 39.7 g)	Begin applications when insects first appear. More than one application may be needed throughout the growing season to obtain knockdown of existing populations. Use the higher rate when for heavy populations.	Dilute recommended amount of product in 1 to 15 gallons of water to treat 1000 sq. ft. Apply during late afternoon or early evening hours and following irrigation. Do not irrigate after application. The use of a lemon fragrance in the spray mix may enhance control by acting as a flushing agent and thus provide increased mole cricket contact. The following products have been shown to be effective flushing agents: Lemon Joy and Lemon Palmolive. The use rate for these lemon-scented products is 2 teaspoons per gallon of water for small total mix volumes or 6 fl. oz. per 50 gallons of water for large total mix volumes.
	Greenbug (Schizaphis graminun) Grasshoppers	0.4 oz (17 g)	Begin applications when insects first appear. Reapply every 7 days as needed.	Dilute product in 4 gallons of water to treat 1000 sq. ft. Do not mow turfgrass for at least 24 hours after application.
	Black Turfgrass Ataenius*	0.9 to 1.4 oz (25.5 to 39.7 g)	Begin applications when insects or their damage first appear.	Dilute product in 5 gallons of water to treat 1000 sq. ft. Irrigate lightly after application (not more than ½ inch). Use the higher rate for severe infestations.

15/21

Dichondra (Plant site for Golf Course and Sod Farm Turfgrass)	Cutworm Flea Beetle Southern Armyworm Yellow Striped Armyworm	0.8 to 1.4 oz (22.7 to 39.7 g)	Begin applications when insects first appear. Reapply every 2 weeks as needed.	Dilute product in 15 gallons of water to treat 1000 sq. ft.
---	--	-----------------------------------	--	---

*Not registered in CA for control of Mole Crickets, Spittlebug, or Black Turfgrass Ataenius.

MOUND TREATMENT OF FIRE ANTS AND HARVESTER ANTS IN TURFGRASS

DO NOT ALLOW LIVESTOCK TO GRAZE TREATED AREAS. DO NOT FEED TREATED GRASS TO LIVESTOCK.

PLANTS	INSECTS	AMOUNT OF PRODUCT	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Turfgrass Residential, Recreational, and Commercial Turf	Imported Fire Ants and Harvester Ants*	Mound Treatment - Drench Method: Use ¾ oz (21.3 g) of product in 5 gallons of water. Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound.	For best results, apply the material in the early morning or late afternoon when the ants are most active. Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest.	Grass in treated areas may be injured. Do not treat mound more than one per season.

*Not registered in CA for control of Harvester Ants

16/21

OUTDOOR AND PERIMETER SPRAY

Location	INSECTS	AMOUNT OF PRODUCT PER 1 (one) GALLON	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Outdoor and Perimeter Area	Wasp	1.2 oz (34 g)	Make treatments early in the morning or late in the evening, as wasps are less active at these times.	Apply as a spot treatment to the nest, nest entrance, and surrounding areas where the wasps may land.
	Ants (excluding Imported Fire Ants and Harvester Ants*) Crickets Cockroaches Earwigs Pillbugs	1.2 oz (34 g)	Begin applications when insects or their damage first appear.	Use specified amount of product in one (1) gallon of water. Apply to a band of soil 6 to 10 feet adjacent to the structure and to a height of 2 to 3 feet on the foundation where pests may be active or may find entrance. Also apply as a residual spray or with a paint brush to surfaces of buildings, window frames, shutters, entryways, screens, eaves, patios, garages, carports, around garage areas and other areas where these pests congregate.
Note: Not for use on residential turfgrass				

17/21

OUTDOOR FLORAL CROPS AND GROUND COVERS

PLANTS	INSECTS	AMOUNT OF PRODUCT PER 100 GALLONS	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Chrysanthemums Dahlias Daisies Easter Lily Gladioli Gypsophila Pachysandra Pansy Peony Roses Sedum Statice Strawflower Yarrow Zinnia	Aphids Thrips Lygus	½ lb (8oz) (226.8g) The maximum amount of product to be applied on a per acre basis is not to exceed 0.75 lb ai/acre for plants grown for cut flower production and 1.0 lb ai/acre for all other plants.	Begin treatments when insects first appear. Repeat applications as needed to achieve control. The minimum spray interval is 3 days	Apply the specified amount of product in 100 gallons of water using a hydraulic sprayer. Multiple applications may cause slight tip burn or marginal leaf necrosis on some varieties. Test on a few plants to determine varietal susceptibility.
Boston Ivy Roses	Japanese Beetles	1 lb (16 oz) (453.6 g) The maximum amount of product to be applied on a per acre basis is not to exceed 0.75 lb ai/acre for plants grown for cut flower production and 1.0 lb ai/acre for all other plants.	Begin treatments when insects first appear. Repeat applications every 2 weeks as needed.	

18/21

COMMERCIAL GREENHOUSE FLORAL AND FOLIAGE PLANT CROPS

Not for use in residential greenhouses

PLANTS	INSECTS	AMOUNT OF PRODUCT PER 100 GALLONS	APPLICATION TIMING	APPLICATION RECOMMENDATIONS
Roses	Leafrollers	<p>½ to ¾ lb (8 to 12 oz) (226.8 to 340.2 g)</p> <p>The maximum amount of product to be applied on a per acre basis is not to exceed 0.75 lb ai/acre for plants grown for cut flower production and 1.0 lb ai/acre for all other plants.</p>	Begin treatments when insects first appear. Use the higher rate when large larvae are present.	Apply the specified amount of product in 100 gallons of water using a hydraulic sprayer.
Foliage Plants Anthuriums Cacti Orchids Poinsettia	Aphids Mealybugs Scale (Crawlers) Thrips Whiteflies	<p>½ lb (8 oz) (226.8 g)</p> <p>The maximum amount of product to be applied on a per acre basis is not to exceed 0.75 lb ai/acre for plants grown for cut flower production and 1.0 lb ai/acre for all other plants.</p>	Begin treatments when insects first appear. A repeat application, at a 2 week interval, may be necessary for control of mealybugs, Scale crawlers, and whiteflies.	<p>Apply the specified amount of product in 100 gallons of water using a hydraulic sprayer. The addition of a wetting agent may be required on difficult to wet foliage.</p> <p>Caution: Phytotoxicity has occurred on the following foliage plants: Bletchum gibbum, Cissus Antarctica, Ficus triangularis, Fittonia verschaffelii, Maranta leuconeura kerchoveana, Pachystachya lutea, Plectanthus australis, Polypodium aureus, Polystichum, Pteris ensiformis, Tolmiea menziesii.</p> <p>As a precaution, spray only a few plants and wait for 2 weeks to observe any varietal phytotoxicity. Application on poinsettias after bract formation may result in Phytotoxicity on certain varieties.</p>

19/21

<p>Carnations Chrysanthemums Roses</p>	<p>Aphids Thrips</p>	<p>½ lb (8 oz) (226.8 g)</p> <p>The maximum amount of product to be applied on a per acre basis is not to exceed 0.75 lb ai/acre for plants grown for cut flower production and 1.0 lb ai/acre for all other plants.</p>	<p>Begin treatments when aphids first appear. As the thrips first appear or at the tight flower bud stage. Repeat applications may be necessary.</p>	<p>Apply the specified amount of product in 100 gallons of water using a hydraulic sprayer.</p> <p>Applications to Carnations and Chrysanthemums more than once every 28 days may cause phytotoxicity.</p> <p>Caution: Phytotoxicity has occurred on the following Chrysanthemum varieties: Albatross, Bonnie Jean, Dixie, Garland, Gem, Iceberg, Pride, Showoff, Statesman, Tally Ho, Westward Ho and Wild Honey.</p> <p>As a precaution, spray only a few plants and wait for 2 weeks to observe any varietal phytotoxicity. Applications to Chrysanthemums and Roses with open flowers may result in flower damage.</p>
--	--------------------------	--	--	--

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool, dry place. Protect from excessive heat. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. For help with any spill, leak, fire or exposure involving this material, call CHEMTREC at 1-800-424-9300

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

CONTAINER DISPOSAL

20/21

Completely empty canister into application equipment. Do not reuse container. Dispose of empty canister in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Note: This product is sold by weight. Settling may have occurred during shipment.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Tree injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or tree conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS or seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS and seller harmless for any claims relating to such factors.

RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions. This warranty does not extend to the use of the product contrary to label instructions, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, RAINBOW TREECARE MAKES NO WARRANTIES OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the fullest extent permitted by law, RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF RAINBOW TREECARE OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS and Seller offer this product, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of RAINBOW TREECARE SCIENTIFIC ADVANCEMENTS.

All TM property of RAINBOW Treecare Scientific Advancements.

© Copyright 2007 RAINBOW Treecare Scientific Advancements.

The RAINBOW logo is a trademark of RAINBOW Treecare Scientific Advancements.

21/21

OPTIONAL MARKETING CLAIMS

Committed to advancing the science of tree care.

Committed to bringing tools inside protocols that get predictable results for the arborist practitioner.

