

# Supplemental Labeling

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Restricted Use Pesticide

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

## Prolex™

EPA Reg. No. 74921-2

## Control of Insects Infesting Soybean

### ATTENTION


- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Prolex™ insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Prolex according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Prolex.

### Directions for Use

Refer to product label for General Use Precautions and Restrictions, Mixing and Application instructions.

**Note:** Numbers in parentheses refer to footnotes below table.

Target Pests	Rate	
	(lb ai/acre)	(fl oz/acre)
bean leaf beetle	0.0075 - 0.0125	0.77 - 1.28
cabbage looper		
corn earworm		
cutworm spp.		
green cloverworm		
Mexican bean beetle		
Mexican corn rootworm beetle (adult)		
northern corn rootworm beetle (adult)		
painted lady (thistle) caterpillar		
potato leafhopper		
saltmarsh caterpillar		
southern corn rootworm beetle (adult)		
soybean aphid (4)		
threecornered alfalfa hopper		
thrips spp. (5)		
velvetbean caterpillar		
western corn rootworm beetle (adult)		
woollybear caterpillar		



2/2

armyworm (1) blister beetle spp. European corn borer fall armyworm (1) grasshopper spp. Japanese beetle (adult) plant bug spp. silverspotted skipper stink bug spp. tobacco budworm (3) webworm spp. yellowstriped armyworm (1)	0.0125 - 0.015	1.28 - 1.54
beet armyworm (3) lesser cornstalk borer (2) soybean looper (2) (3) spider mite spp. (2)	0.015	1.54

<sup>1</sup> Use higher rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See resistance statement under General Use Precautions and Restrictions.

<sup>4</sup> Use a rate in the lower end of the rate range for early season applications and/or lighter populations.

<sup>5</sup> **Does not include western flower thrips.**

#### Remarks:

- Apply as required by scouting, usually at intervals of 5 days or more. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- **Do not** graze or harvest treated soybean forage, straw, or hay for livestock feed.
- Apply with ground or air equipment, using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial-applied corn rootworm control program, use a minimum of 1.02 fl oz per acre (0.01 lb active ingredient per acre).
- **Do not** apply more than 0.03 lb active ingredient (0.19 pint) per acre per season.
- **Preharvest Interval: Do not** apply within 30 days of harvest. **Note:** The 30-day PHI supercedes the previous PHI of 45 days.

™Trademark of Pytech

R005-005

EPA-accepted \_\_\_/\_\_\_/\_\_\_

Initial printing.