

74921-3

01105/2009

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

January 05, 2009

Mr. Eric Maurer,
Regulatory Director, North America
Authorized Agent for Pytech Chemicals GmbH
Cheminova, Inc.
1600 Wilson Blvd., Suite 700
Arlington, VA 22209

Subject: Supplemental labeling to add new uses, on pistachios and okra per approved IR-4 Tolerance
Petition #7E7287.
Proaxis™ (EPA Reg. No. 74921-3)
Your Submission dated October 25, 2007

Dear Mr. Maurer:

The above referenced amendment, submitted in connection with registrations under FIFRA Section 3(c)(7)(B), as amended, is acceptable.

Please note that we have approved the supplemental labels for the subject gamma-cyhalothrin product with the understanding that the labeling are consistent with the Agency's previously approved main and/or supplemental labels for similar registered lambda-cyhalothrin products. At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate the supplemental labeling into the product's main labeling.

Enclosed, please find stamped copies of the labels for your records. If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely,

A handwritten signature in black ink, appearing to read "Kimberly Nesci".

Kimberly Nesci, Product Manager 11
Insecticide Branch
Registration Division (7505P)

Enclosure:

Supplemental Labeling

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Restricted Use Pesticide
Due to toxicity to fish and aquatic organisms.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Proaxis™

EPA Reg. No. 74921-3

Control of Insects Infesting Pistachios

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Proaxis™ insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Proaxis according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Proaxis.

Directions for Use

Refer to product label for General Use Precautions and Restrictions, Mixing and Application instructions.

Note: Numbers in parentheses refer to footnotes below table.

Target Pests	Rate	
	(lb ai/acre)	(fl oz/acre)
ants	0.01 - 0.02	2.56 - 5.12
chinch bug		
codling moth		
filbertworm		
leaffooted bug		
leafroller spp.		
navel orangeworm		
peach twig borer		
plant bug spp.		
stink bug spp.		
walnut aphid		
walnut husk fly spp. (adult)		

ACCEPTED
JAN 05 2009
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 74921-3

Remarks:

- Apply as required by scouting, usually at intervals of 5 days or more. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment, using sufficient water to obtain full coverage of foliage or target area. When applying by air, apply in a minimum of 5 gallons of water per acre.
- **Do not** apply more than 0.08 lb active ingredient (1.28 pints of Proaxis) per acre per year.
- **Do not** apply more than 0.06 lb active ingredient (0.96 pints of Proaxis) per acre per year post bloom.
- **Preharvest interval: Do not** apply within 14 days of harvest.

™Trademark of Pytech
R004-010
EPA-accepted ___/___/___
Initial printing.

Supplemental Labeling

Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Restricted Use Pesticide
Due to toxicity to fish and aquatic organisms.
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Proaxis™ EPA Reg. No. 74921-3

Control of Insects Infesting Okra

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Proaxis™ insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Proaxis according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Proaxis.

Directions for Use

Refer to product label for General Use Precautions and Restrictions, Mixing and Application instructions.

Target Pests	Rate	
	(lb ai/acre)	(fl oz/acre)
cabbage looper cutworm spp. hornworm spp.	0.0075 - 0.0125	1.92 - 3.20
aphid spp. (2) (3) beet armyworm (1) (3) blister beetle spp. Colorado potato beetle (3) cucumber beetle spp. (adult) European corn borer (4) fall armyworm (1) flea beetle spp. grasshopper spp. Japanese beetle (adult) leafhopper spp. leafminer spp. (2) meadow spittlebug pepper weevil (adult) (2) plant bug spp. southern armyworm (1) spider mite spp. (2) stalk borer (4) stink bug spp. thrips (3) (5)	0.01 - 0.015	2.56 - 3.84

ACCEPTED
JAN 05 2009
Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 74921-3

4/4

tobacco budworm (3)		
tomato fruitworm		
tomato pinworm		
tomato psyllid (2) (3)		
vegetable weevil (adult)		
whitefly spp. (2) (3)		
yellowstriped armyworm (1)		

¹ For control of first and second instars only.

² Suppression only.

³ See resistance statement under General Use Precautions and Restrictions on container label.

⁴ For control before larvae bore into the plant stalk or fruit.

⁵ **Does not include western flower thrips.**

Remarks:

- Apply as required by scouting, usually at intervals of 5 days or more. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment, using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre.
- **Do not** apply more than 0.18 lb active ingredient (2.88 pints) per acre per season.
- **Preharvest Interval: Do not** apply within 5 days of harvest.

Note: This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife, and highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

™Trademark of Pytech
R004-011.
EPA-accepted ___/___/___
Initial printing.