

75266 -1

11/30/2010

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

NOV 30 2010

Jeff Jones, Agent for Deardorff Fitzsimmons Corp.
c/o Delta Analytical Corp.
12510 Prosperity Drive, Suite 160
Silver Spring, MD. 20904

Subject: Agency Initiated Action
Activate 5.25% Institutional Bleach
EPA Registration 75266-1

Dear Mr. Jones:

The Agency has reviewed your application for amendment submitted in accordance with continuing registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA), as amended, and determined the action acceptable.

In summary, you have submitted additional information to upgrade product performance data claims against *Pseudomonas aeruginosa*. See the enclosed review, DP 383414, for additional information.

A stamped copy of the accepted labeling is enclosed. Submit one copy of your final printed labeling before distributing or selling the product bearing the revised labeling. Should you have any questions or comments concerning this letter, please call Tom Luminello at (703) 308-8075.

Sincerely,

A handwritten signature in black ink that reads "Wanda Y. Henson".

Wanda Y. Henson
Acting Product Manager 32
Regulatory Management Branch II
Antimicrobials Division (7510P)

Activate™ 5.25% Institutional Bleach

For use only with the Activate™ Bleach Dilution Sprayer

The Activate™ Bleach Dilution Sprayer automatically dilutes and mixes this product to a ready-to-use 10% solution (9:1 ratio of water to bleach) equal to approximately 5,000 ppm available chlorine for hospital, laboratory and veterinary use on hard, non-porous surfaces.

Cleans & disinfects in one step

Kills HIV-1 & HBV in 30 Seconds
Kills Bacteria in 2 Minutes:

- *Staphylococcus aureus*
- *Methicillin resistant Staphylococcus aureus (MRSA)*
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Streptococcus pyogenes*
- *Vancomycin resistant Enterococcus faecalis (VRE)*
- *Acinetobacter baumannii*

Active Ingredient:

Sodium Hypochlorite 5.25%

Other Ingredients:..... 94.75%

TOTAL 100.00%

Contains 5% available chlorine by weight

KEEP OUT OF REACH OF CHILDREN DANGER

Read Complete Directions and Precautions on Package Rear & Side Panels

FIRST AID	
Call poison control center or doctor for treatment advice. Have product container or label with you when you call a poison control center or doctor or when going for treatment.	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin with plenty of water for 15-20 minutes.
If swallowed:	<ul style="list-style-type: none"> • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.	

EPA Reg. No. 75266-1
EPA Est. No. 75266-OR-001

NET CONTENTS:
Batch Code:

Manufactured by:
Deardorff Fitzsimmons Corporation
PO Box 539
Merlin, OR 97532
1-888-582-2700

ACCEPTED
NOV 30 2010
75266-1

with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) and Hepatitis B Virus (HBV).

Special Instructions for Cleaning and Decontaminating Against HIV-1 and HBV on Surfaces/Objects Soiled with Blood/Body Fluids: When handling items soiled with blood/body fluids, personal protective equipment such as disposable gloves, gowns, masks, and eye coverings must be used. Blood/body fluids must be thoroughly cleaned from surfaces/objects prior to disinfecting. Spray Activate™ on surface/object until all surfaces are thoroughly wet; let stand for 30 seconds. Wipe with a clean cloth or paper towel, and allow to air dry. Blood/body fluids and cleaning materials must be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

To Clean and Disinfect in a Veterinary Application: Use to clean and disinfect hard, non-porous surfaces such as feeding and watering equipment, cages, utensils, instruments, kennels, stables, catteries, etc. Remove all animals and feed from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings, and manure from walls, floors, and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering equipment. Pre-clean all surfaces with soap or detergent and rinse with water. Spray Activate™ on surface/object until all surfaces are saturated; let stand for 2 minutes. To kill TB, allow 30 seconds at room temperature. Rinse all surfaces with potable water. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treated surfaces have been thoroughly rinsed with water and allowed to dry. Thoroughly scrub all treating, feeding, and watering appliances with soap or detergent, and rinse with potable water before re-use.

STORAGE & DISPOSAL

Do not contaminate food or feed by storage, disposal or cleaning of equipment.

Pesticide Storage

This product degrades with age; test active level of sprayed product periodically with high-level chlorine test strips. Store in a cool (50° – 72°F), dry area away from direct sunlight and heat to avoid deterioration. In case of spill, flood area with large quantities of water.

Pesticide Disposal

Products or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer.

Container Handling: Non-refillable container. Do not re-use or refill this container. Offer for recycling if available or place in trash collection.

ACCEPTED
NOV 30 2010
75266-1


[Water Bottle Label]

WATER BOTTLE For use only with the Activate™ Bleach Dilution Sprayer

The Activate™ Bleach Dilution Sprayer automatically dilutes and mixes Activate™ 5.25% Institutional Bleach to a ready-to-use 10% solution (9:1 ratio of water to bleach) equal to approximately 5,000 ppm available chlorine for hospital, laboratory and veterinary use on hard, non-porous surfaces.

Activate™ Bleach Dilution Sprayer Instructions

1. Fill the sprayer's empty water bottle with cool tap water, and then insert the sprayer's tube down into the filled water bottle. Close the sprayer tab to lock the water bottle into place on the sprayer.
2. Remove the threaded plastic cap from the bleach cartridge. Do not puncture or tamper with the plug that is in the neck of the bleach cartridge.
3. Lock the bleach cartridge onto the other side of the sprayer.
4. Pointing the sprayer away from you into a sink or basin, pump the trigger 20 times to prime the system.
5. The sprayer is now ready to use. Hold the sprayer 6" to 8" from the surface, and spray thoroughly until surface is completely wet.
6. Refill the water bottle when empty.
7. Always re-prime the system when installing a new bleach cartridge.



Note that a safety feature ensures the sprayer will not function if either the bleach cartridge or the water bottle is empty.

The water in this bottle is not for human consumption.

Container Handling: Refillable container. Refill this container with water only. Do not re-use this container for any other purpose. Offer for recycling if available or place in trash collection.

Manufactured by:
Deardorff Fitzsimmons Corporation
PO Box 539
Merlin, OR 97532
1-888-582-2700

NOV 30 2010
75266-1