


79442-2

11/18/2004



1/3

[Front panel]

# EXOSEX - TPW®

A novel pheromone product for the suppression of tomato pinworm (*Keiferia lycopersicella*) on tomatoes

This package contains: 5 x high impact polystyrene autoconfusion station  
5 x tomato pinworm autoconfusion tray

## AUTOCONFUSION TRAY

[Net Weight: 2.5 g of pheromone impregnated wax powder]

### ACTIVE INGREDIENT

E-4-Tridecen-1-yl acetate

0.097%

Z-4-Tridecen-1-yl acetate

0.003%

### OTHER INGREDIENT

99.900%

Total

100.000%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 79442-E

EPA Establ. No. 79442-GBR-1  
Date of manufacture: mm/dd/yyyy

EXOSECT LTD.  
2 Venture Road  
Chilworth Science Park  
Southampton SO16 7NP  
United Kingdom

[Back panels]

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust, wear a dust mask, protective eyewear and gloves (appropriate for applying the product). Remove contaminated clothing and wash before reuse. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID**

**IF INHALED:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, <sup>then</sup> give artificial respiration, preferably mouth to mouth if possible.
- Call poison control center or doctor for treatment advice.

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment advice between the hours of 9.30 am and 7.30 pm eastern time.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

**STORAGE AND DISPOSAL STATEMENTS**

Do not contaminate water, food or feed by storage or disposal

**STORAGE:**

Store in unopened package at temperatures below 40°F. Do not open until ready to use.

**PESTICIDE DISPOSAL:**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

**CONTAINER DISPOSAL:**

Dispose of in a sanitary landfill or incineration, or as allowed by state and local authorities, by burning. If burned, stay out of the smoke.

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling**

Use within six months of manufacture  
Check front panel for date of manufacture.

Use experience with this product dictates a minimum treatment area of 7.5 acres

- 1. CROP: Tomatoes
- 2. PEST: Tomato pinworm [*Keiffera lycopersicella*]
- 3. APPLICATION: Deploy Exosex Autoconfusion Stations at the rate of 10 - 12 stations per acre. Place the stations at the top of the plant. Remove the plastic coverlid of the autoconfusion tray, bring the two wings of the station together, into a "click fit" to form a delta (triangle cross-section) The station is affixed to the plant, trellis or support stake with the provided ties which thread through the holes in the top ridge of the station.
- 4. TIMING: Deploy the stations prior to the onset of tomato pinworm emergence in the spring. The stations are active for an extended period of time, depending on the temperature. Make a second application prior to the emergence of the second moth flight but not later than October 1.
- 5. NOTES:
  - If the field borders other fields where tomato pinworm is not controlled, migration of these moths can reduce the level of control achieved. This may be counteracted by:
    - treatment of entire blocks of fields and not just sections of conventionally treated areas.
    - doubling the number of stations along such borders in a strip of 4 - 6 rows of tomato plants.
    - treatment of the infestation source with conventional insecticide.
    - monitor the occurrence of tomato pinworm in the treated area using weekly pheromone traps catches. If monitoring traps show an extremely high infestation in early spring, consider a pre-application treatment of insecticide. If monitoring traps indicate an extended period of infestation (longer than 8 weeks), replace the Exosex stations, to coincide with subsequent tomato pinworm flights or apply a conventional insecticide spray.

**WARRANTY**

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing, unfavorable temperatures, presence of other materials, weather, watering practices, nature of the soil, disease problem, condition of the crop, pre-existing conditions and other conditions influencing the use of this product are beyond the control of the seller. Buyer assumes all risks associated with the use, storage, or handling of this material not in strict accordance with directions given herewith. **NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.**