

79676-20

10/10/2008

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

OCT 10 2008

Mr. Michael Kellogg
Etigra
4110 136th St. N.W.
Gig Harbor, WA 98332

Dear Mr. Kellogg:

Subject: Double O E-Pro Herbicide
Label Amendment
EPA Registration No. 79676-20
Submission Dated: July 15, 2008

The labeling referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide act, as amended is acceptable.

Submit 1 (one) copy of your final printed labeling before you release the product for shipment. A stamped copy of the label is enclosed for your records.

Sincerely,

A handwritten signature in cursive script that reads "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

2/11

Double O E-Pro Herbicide Ornamental Herbicide

For Sale To, Use and Storage By Commercial Nursery, Cut Flower, Foliage And Landscape Personnel Only

Double O E-Pro Herbicide provides broad-spectrum preemergence control of weeds (including difficult weed species, such as spurge, chickweed, groundsel, and oxalis) in container, field grown and landscape ornamentals, cut flowers and ornamental foliage.

ACTIVE INGREDIENTS:

Oxyfluorfen*2.0%
Oryzalin**1.0%

OTHER INGREDIENTS:97.0%

TOTAL: 100.0%

*CAS No. 42874-03-3

**CAS No. 19044-88-3

**Keep out of Reach of Children
CAUTION**

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 79676-20

Manufactured for:
Etigra®
501 Cascade Pointe Lane
Suite 103
Cary, NC 27513
www.etigra.com

ACCEPTED
OCT 10 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

79676-20

EPA Est. No. 5905-IA-001

Double O E-Pro Herbicide contains oxyfluorfen and oryzalin, the active ingredients used in Rout®.

Net Weight:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Chemical-resistant apron for mixers and loaders

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)-(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

GENERAL INFORMATION

Double O E-Pro Herbicide is a pre-emergent herbicide that combines two herbicides into a clay granule to provide a broader spectrum of weed control and ornamental plant tolerance. Double O E-Pro Herbicide does not control established weeds and is not registered for use on vegetable or edible fruit bearing crops, turf, bedding plants or flowering crops except as described in the Landscape Ornamentals and Cut Flower and Ornamental Foliage sections.

Double O E-Pro Herbicide provides pre-emergent control of the following weeds:

Grassy Weeds

- | | |
|---|---|
| Annual bluegrass (<i>Poa annua</i>) | Foxtail, Green (<i>Setaria Viridis</i>) |
| Barnyardgrass (<i>Echinochloa crusgalii</i>) | Foxtail, Yellow (<i>Setaria lutescens</i>) |
| Crabgrass (<i>Digitaria Sanguinalis</i>) | Goosegrass (<i>Eleusine indica</i>) |
| Fall Panicum (<i>Panicum dichotomiflorum</i>) | Kikuyu Grass (<i>Pennisetum clandestinum</i>) |

Broadleaf Weeds

- | | |
|--|---|
| <i>Ageratum (Ageratum conyzoides)</i> | <i>Moss (Isopetyquim albesceus)</i> |
| <i>Artillery weed (Pilea micropylla)</i> | <i>Oxalis (Oxalis stricta)</i> |
| <i>Bittercress (Cardamine pennsylvanica)</i> | <i>Pearlwort (Sagina procumbens)</i> |
| <i>Buttonweed (Borreria laevis)</i> | <i>Pigweed (Amaranthus retroflexus)</i> |
| <i>Castor bean (Rincinus communis)</i> | <i>Primrose (Oenothera Cutleaf evening laciniata)</i> |
| <i>Chickweed, Common (Stellaria media)</i> | <i>Purslane (Portulaca oleracea)</i> |
| <i>Clover, White (Trifolium repens)</i> | <i>Sheperdspurse (Capsella bursa-pastoris)</i> |
| <i>Dandelion (Taraxacum officinale)</i> | <i>Sowthistle (Sonchus arvensis)</i> |
| <i>Fireweed (Epilobium angustifolium, Erechitites hieraciifolia)</i> | <i>Sowthistle, common (Sonchus oleraceus)</i> |
| <i>Groundsel (Senecio vulgaris)</i> | <i>Spurge, garden (Euphorbia hirta)</i> |
| <i>Lambsquarter (Chenopodium album)</i> | <i>Spurge, Hairy (Euphorbia vermiculata)</i> |
| <i>Marestail (Conyza canadensis)</i> | <i>Spurge, Prostrate (Euphorbia supina)</i> |
| <i>Medic (Medicago trunculata)</i> | <i>Spurge, Spotted (Euphorbia maculata)</i> |

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

USE GUIDELINES AND PRECAUTIONS

- For best results ensure an even distribution of particles at the proper application rate.
- Do not apply Double O E-Pro Herbicide to wet or dew covered foliage. However, should this happen, irrigate immediately to completely remove herbicide particles from leaf surfaces.
- Do NOT apply Double O E-Pro Herbicide in enclosed greenhouse or shadehouse type structures. Make applications at least two (2) weeks prior to enclosure in greenhouses or shadehouses.
- Do NOT apply Double O E-Pro Herbicide in greenhouse liner production beds; apply only to container, field grown and landscape ornamentals, cut flower and foliage crops listed.
- Do NOT apply Double O E-Pro Herbicide within two (2) weeks of shipping.
- Do NOT apply product directly to the bare roots of woody ornamentals or injury may result.
- In order to avoid foliage injury do NOT apply to whorl-leaved plants such as *Dracaena* spp., as their leaves channel the herbicide granules to the leaf base and makes removal of granules difficult (Note exception under 'Drancaena Field Stock').

- Newly forming leaves are particularly sensitive to this product and spotting or leaf drop may result. Do NOT apply the product two (2) weeks prior to or two (2) weeks following leaf bud break, or during periods of vigorous new growth.
- Within two (2) months following application, do NOT apply other herbicides to areas treated with Double O E-Pro Herbicide. Compatibility of Double O E-Pro Herbicide with other ornamental herbicides has not been verified.
- Do NOT disturb the soil surface following an application as the herbicide barrier will be broken resulting in less than optimum weed control.
- Do NOT incorporate this product into potting media or soil as dilution of the active ingredient and possible injury to plant roots may result.
- Do NOT apply in windy conditions that might cause uneven distribution of the granules or move the product onto sensitive species.
- Do NOT allow livestock to graze or feed on vegetation from treated fields or orchards.
- Do NOT apply by aerial application except where noted for agricultural uses in California.
- Do NOT apply by chemigation.

Application Restrictions - Container-grown ornamental: granular application rate of 2 lbs. a.i./A/application (6 lbs. a.i./season).

GENERAL APPLICATION INSTRUCTIONS

When applying this product, users should exercise reasonable judgment and caution. Because it is impossible to test this product with all possible combinations of plants and growing conditions, it is recommended that an initial test application be made to a subset of the plants to assure desired performance under your specific growing conditions. Users who wish to use Double O E-Pro Herbicide ornamental herbicide on ornamental species or nonbearing fruit trees not listed on this label may determine suitability for such uses by making trial application of this product at a recommended rate to a small number of plants. Prior to using this product on a large number of plants, the test treatment should be observed for 3 to 6 months for signs of herbicidal injury to determine if the treatment is safe for use on the target plant species.

Apply Double O E-Pro Herbicide at the rate recommended in the appropriate section of this label with a properly calibrated spreader in order to assure accurate, even particle distribution. Once this product has been applied, do not disturb the soil surface or the herbicide barrier will be broken. For best results, immediately following application of Double O E-Pro Herbicide, water in with approximately 1/2 inch of water in order to activate the herbicide and wash particles off the foliage.

	APPLICATION RATE PER				
	Square Foot	100 Square feet	Acre	Square Meter	100 Square Meters
Amount of Double O E-Pro Herbicide	1g	100 g	100 lbs.	10.8 g	1.1 kg

1 ounce = 28 grams (g)

PLANT TOLERANCE TO DOUBLE O E-PRO HERBICIDE

The following species are tolerant of Double O E-Pro Herbicide when used at the recommended application rates:

- Abelia
 - x *grandiflora* (Glossy Abelia)
- Achillea
 - Millefolium 'Cerise Queen'* (Yarrow)
- Acer
 - Palmatum 'Atropurpureum'* (Red Japanese Maple)
- Alyogyne
 - Huegelii* (Blue Hibiscus)
- Andromeda
 - Polifolia* (Bog Rosemary)
- Anthurium
 - Andraeanum* (Ozaki Red Anthurium)
- Arctostaphylos
 - Uva-ursi* (Bearberry/Kinnikinnick)
- Banksia
 - Menziesii 'Raspberry Frost'* (Raspberry Frost Banksia)
- Berberis
 - Thunbergii 'Crimson Pygmy'* (Crimson Pygmy Barberry)
- Bougainvillea
 - spp. *'Barbara Karst'* (Barbara Karst Bougainvillea)
- Buxus
 - Microphylla 'Green Velvet'* (Boxwood)
 - Microphylla japonica* (Japanese Boxwood)
 - Sempervirens 'Welleri'* (English Boxwood)
- Camellia
 - Japonica 'Lady Laura'*
- Carissa
 - Grandiflora 'Fancy'* (Natal Plum)
- Catharanthus
 - Roseus 'Vinca Blue Pearl'*
- Chamaecyparis
 - Obtusa 'Letinea Aurea'* (Gold Hinoki False C.) *pisifera*
 - 'Filifera Aurea Nana'* (Dwarf Golden Thread-Branch C.)
 - Pisifera 'Fillfera Aurea'* (Cypress)
- Cercis
 - Canadensis* (Redbud)
- Codiaeum
 - Norma 'Low Life'* (Low Life Croton)
- Coreopsis
 - Verticillata 'Moonbeam'*
- Cornus
 - Baileyi* (Bailey Dogwood)
 - Florida* (Flowering Dogwood)
- Cotoneaster
 - Dammeri 'Coral Beauty'* (Coral Beauty Cotoneaster)
 - Salicifolius 'Repens'* (Willowleaf Cotoneaster)
- Cupressus
 - Sempervirens 'Glauca'* (Italian Cypress)
- Deutzia
 - x *'Magician'*
- Dianthus
 - spp. *'Baths Pink'*
- Diosma
 - Ericoides* (Breath of Heaven)
- Dracaena
 - Massangeana* (Complant)
 - Deremensis 'Wameckii'* (Striped Dracaena)
- Elaeagnus
 - Pungens 'Fruitlandii'* (Fruitland Silverberry)
- Eriobotrya
 - Japonica* (Loquat)
- Escallonia
 - x *exoniensis 'Fradesii'* (Frades Escallonia)
- Euonymus
 - Alata 'Compacta'* (Dwarf Burning Bush)
 - Fortunei 'Emerald'n Gold'* (Emerald Gold E.)
 - Fortunei 'Gold Edge'* (Gold Edge E.)
 - Fortunei 'Vegeta'* (Euonymus Vegetus)
 - Klatschovica 'Manhattan'* (Manhattan E.)
- Euphorbia
 - Milii* (Crown of Thoms)
- Ficus
 - Benjamina 'Lady Diane'* (Weeping Fig)
 - Lyrata* (Fiddleleaf Fig)
 - Microcarpa nitida* (Little Leaf Ficus)
- Forsythia
 - x *intermedia 'Lynnwood'* (Lynnwood Gold F.)
- Fothergillia
 - Gargenii 'Mt Airy'*
- Fragaria
 - x *'Pink Panda'* (Sand Strawberry)
- Gardenia
 - Jasminoides 'August Beauty Gardenia'*
- Gelsemium
 - Sempervirens* (Carolina Jessamine)
- Gypsophila
 - spp. (Baby's Breath)
- Hemerocallis
 - spp. (Daylily)
- Hibiscus
 - Rosa-sinensis 'Brilliant'* (Brilliant Hibiscus)
 - Rosa-sinensis 'Crown of Bohemia'* (Crown of Bohemia H.)
- Hypericum
 - spp. (St. Johnswort)
- Iberis
 - Sempervirens* (Evergreen Candy Tuft)
- Ilex
 - Aquipemyi 'San Jose'* (San Jose Holly)
 - Cornuta 'Burfordii Nana'* (Dwarf Burford Holly)
 - Cornuta 'Needlepoint'* (Needle Point Holly)
 - Cornuta 'Rotunda'* (Dwarf Chinese Holly)
 - Crenata 'Convexa'* (Convex Japanese Holly)
 - Crenata 'Hetzii'* (Japanese Holly)
 - Opaca* (American Holly)
 - Vomitorea 'Schillings'* (Shillings Holly)
 - spp. *'Blue Girl'* (Dwarf Holly)
- Jasminum
 - Nitidum* (Angelwing Jasmine)
 - spp. (Confederate Jasmine)

- 7/11
- Juniperus*
Chinensis 'Gold Coast' (Gold Coast Juniper)
Chinensis 'Mint Julip' (Mint Julip Juniper)
Chinensis 'Pfitzerana Glauca' (Blue Pfitzer Juniper)
Chinensis 'Pfitzerana Old Gold' (Old Gold Juniper)
Chinensis 'Sargentii Viridis' (Green Sargent Juniper)
Chinensis 'Sea Green' (Sea Green Juniper)
Conferta 'Blue Pacific' (Blue Pacific Juniper)
Conferta (Shore Juniper)
Horizontalis 'Bar Harbor' (Bar Harbor Juniper)
Horizontalis 'Douglasii' (Waukegan Juniper)
Horizontalis 'Plumosa' (Andorra Juniper)
Sabina 'Broadmoor' (Broadmoor Juniper)
Sabina 'Tamariscifolia' (Tam Juniper)
Squamata 'Blue star' (Blue Star Juniper)
- Lantana*
Montevidensis (Weeping Lantana)
Montevidensis 'Radiation' (Radiation Lantana)
- Leucodendron*
x 'Safari Sunset' (South African Rose)
- Leucospermum*
Cordifolium (Pincushion Protea)
- Ligustrum*
Japonicum 'Texanum' (Waxleaf Privet, Texas Privet)
Lucidum (Glossy Privet)
Sinense 'Variegata' (Dwarf Variegated Chinese Privet)
spp. 'Jack Frost' (Jack Frost Ligustrum)
x 'Vaicaryi' (Vicary Golden Privet)
- Lonicera*
x *xylosteoides* 'Clavey's Dwarf' (Clavey's Dwarf Honeysuckle)
- Loropetalum*
Chinensis 'Burgundy'
- Mahonia*
Aquifolium (Oregon Grape Holly)
Chinensis 'Pfitzerana Glauca' (Blue Pfitzer Juniper)
Repens (Creeping Mahonia)
- Monarda*
Didyma 'Claire Grace'
- Microbiota*
Decussata (Russian Cypress)
- Nandina*
spp. 'Compacta'
Domestica 'Nana' (Dwarf Nandina)
- Nerium*
Oleander (Oleander)
- Osmanthus*
Frangrans (Sweet Osmanthus)
Heterophyllus 'Variegatus' (Variegated Osmanthus)
- Photinia*
x *fraseri* (Fraser, Photinia, Red Tip)
- Picea*
Abies 'Ohlendorffii' (Norway Spruce)
Glauca 'Conica' (Dwarf Alberta Spruce)
Pungens 'Glauca' (Colorado Blue Spruce)
- Pieris*
Japonica 'Mountain Fire' (Mountain Fire Andromeda)
- Pinus*
Elliotii (Slash Pine)
Mugo Mugo (Mugho Pine)
Radiata (Monterey Pine)
Strobus (Eastern White Pine)
Sylvestris (Scotch Pine)
Thunbergiana (Japanese Black Pine)
- Pittosporum*
Crassifolium (Evergreen Pittosporum)
Tenuifolium (Kohuhu Pittosporum)
Tobira (Green Pittosporum)
Tobira 'Wheeler's Dwarf'
- Podocarpus*
Gracilior (Weeping Podocarpus)
Occidentalis 'Pyramidalis' (Pyramidal Arborvitae)
- Potentilla*
Fruticosa 'Gold Drop' (Gold Drop Potentilla)
- Protea*
Nerifolia 'Pink Mink' (Pink Mink Protea)
- Prunus*
x *cistena* (Red-Leaf Plum)
- Pyracantha*
Fortuneana 'Graberii' (Graber Pyracantha)
'Mohave' (Mohave Pyracantha)
'Navajo' (Navajo Pyracantha)
- Quercus*
Shumardii (Shumard Red Oak)
- Raphiolepis*
Indica 'Ballerina' (India Hawthorn)
Indica 'Pinkie' (India Hawthorn)
Indica 'Pink Cloud' (Pink Cloud Raphiolepis)
- Rhododendron*
spp. 'Dora Amateis' (Dora Amateis R.)
spp. 'Coral Bell' (Azalea)
spp. 'Copperman' (Azalea)
spp. x *exbury* (Azalea)
spp. 'Pink Ruffles' (Azalea)
spp. 'Snow' (Azalea)
- Salvia*
Farinacea (Dwarf Blue Salvia)
Splendens 'Lavender Vista'
- Scabiosa*
Columbaria 'Butterfly Blue'
Columbaria 'Pink Mist'
- Schefflera*
Arboricola (Dwarf schefflera)
- Spirea*
x *bumalda* 'Anthony Waterer' (Anthony Waterer S.)
- Syringa*
Meyeri 'Pallbin' (Lilac)
- Taxus*
Media 'Hicksii'
Media 'Runyoni'
- Temstroemia*
Gymnanthera
- Thuja*
Occidentalis 'Nigra' (Dark Green Arborvitae)
Occidentalis 'Ellwangeriana Aurea' (Golden Arborvitae)
Occidentalis 'Hetz's Midget' (Hetz Midget Arborvitae)
Occidentalis 'Pyramidalis' (Pyramidal Arborvitae)
Occidentalis 'Smaragd' (Emerald Arborvitae)
Orientalis 'Aurea Nana' (Dwarf Golden Arborvitae)
Orientalis 'Blue Cone' (Blue Cone Arborvitae)
- Trachelospermum*
Asiaticum 'Solo' (Asiatic Jasmine)
Asiaticum 'White Mist' (Variegated Asiatic Jasmine)
Jasminoides (Confederate or Star Jasmine)
- Tsuga*
Canadensis (Canadian Hemlock)
- Vaccinium*
Ashei 'Homebell' (Homebell Blueberry)

- Verbena
 - Peruviana* (Sand Verbena)
- Virburnum
 - Carlesil* (Korean Spice Viburnum)
 - Davidii* (David Viburnum)
 - Macrocephalum* (Chinese Snowball)
 - Plicatum 'Tomentosum'* (Double Fire Viburnum)

- Vitis
 - x SV5279 '*Aurora*' (White Wine Grape)
- Weigela
 - Florida 'Java Red'* (Java Red Weigela)
 - spp. '*Boskoop Glory*' (Boskoop Glory W.)
- Xylosma
 - Congestum*

Double O E-Pro Herbicide is not recommended for use on the following species:

- Achillea
 - spp. '*Coronation Gold*' (Yarrow)
- Ageratum
 - spp. '*Hawaii Blue*'
- Calluna
 - spp. '*Forsey*' (Heather)
- Campanula
 - Persicifolia 'Blue'*
- Celosia
 - spp. '*Castle Yellow*'
- Chrysanthemum
 - spp. '*Legends*'
- Clematis
 - x '*Dutchess of Edinburgh*'
- Dianthus
 - Caryophyllus*
 - Deltoides 'Artic fire'*
- Digitalis
 - spp. '*Excelsior*'
- Echinacea
 - spp. '*Magnus*'
 - Pallida*
 - Purpurea 'Leuchstem'*
- Euonymus
 - Japonica 'Silver King'* (Silver King Euonymus)
- Euryops
 - Pectinatus* (Greyleaf Daisy)
- Ficus
 - Pumila* (Creeping Fig)
- Geranium
 - spp. '*Pinto Red*'
- Hemerocallis
 - spp. '*Happy Returns*' (Daylilly)
- Hydrangea
 - Arborescens 'Annabelle'*
 - Paniculata "Grandiflora"* (Peegee Hydrangea)
- Lantana
 - spp. '*Patriot Desert Sunset*'
- Leucanthemum
 - Maximum 'White Knight'* (Shasta Daisy)
- Lilium
 - Orientalis 'Mr. Ruud'*
- Llex
 - Crenata 'Helleri'* (Helleri Japanese Holly)
- Lupinus
 - spp. '*RH My Castle*' (Lupine)
- Magnolia
 - Stellata 'Pinkie'*

Papavor
Orientale 'Pizzicato Mix' (Poppy)

Penstemon
spp. 'Thorn'

Petunia spp.

Prunus
Glandulosa (Dwarf Flowering Almond)

Rhododendron
spp. 'English Roseum'

Rudbeckia
spp. 'Goldstrum' (Coneflower)

Salvia
spp. 'May Knight'

Scabiosa
spp. 'Pink Carnival'

Sedum
Spectabile 'Autumn Joy'
Brevifolium (Stonecrop)

Tagetes
Erecta (Marigold)

Trachelospermum
Asiaticum 'Oblanceolatum' (Hardy Star Jasmine)

Vinca
Minor (Dwarf Periwinkle)

Zinnia spp.

CONTAINER GROWN ORNAMENTALS

Apply Double O E-Pro Herbicide prior to weed seed germination or following complete weed removal in the spring or fall. Make repeat applications throughout the growing season at a *maximum* of three (3) month intervals (under most conditions three months is the maximum timeframe for residual effect for Double O E-Pro Herbicide). Make Fall applications two (2) weeks prior to placing plants in enclosed greenhouse structures.

Potted Liners - Prior to transplanting into containers, remove all established weeds from the liners. To settle the potting media around the root systems, after planting and prior to chemical application irrigate with approximately ½ inch of water and then allow foliage to dry. Broadcast Double O E-Pro Herbicide evenly over the container growing area and irrigate immediately after application with approximately ½ inch of water.

Bare-Root Liners - Remove all established weeds prior to application of Double O E-Pro Herbicide. Within three (3) weeks after potting, broadcast apply the Double O E-Pro Herbicide over the container growing area evenly and immediately after application irrigate with approximately ½ inch of water.

Recycled Water Precaution: When runoff water is recycled, care must be taken to locate pump intake line opening(s) close to the collection pond surface to avoid disturbing and recirculating clay fractions/organic matter accumulation near the pond bottom. This debris/settlement may contain amounts of previously applied pesticides tied-up on these clay/organic matter fractions and may lead to plant injury.

FIELD GROWN ORNAMENTALS AND LABELED CHRISTMAS TREES

Apply Double O E-Pro Herbicide either immediately after planting, in the spring prior to weed germination, or following complete weed removal using a rate of 50 lbs. per acre (one bag / acre) for a maximum of two (2) applications during the growing season or 100 lbs. per acre for a single application in a growing season. Do not reapply at less than two (2) month intervals for Christmas trees and at less than three (3) month intervals for field grown ornamentals.

NOTE: If irrigation is not available, Double O E-Pro Herbicide is NOT recommended for use on bare root stock.

Potted Liners - Prior to transplanting into the field, remove all established weeds from liners. To settle the soil around the root systems, after planting and prior to application irrigate with approximately 1/2 inch of water and then allow foliage to dry. Broadcast Double O E-Pro Herbicide evenly over the field growing area and irrigate immediately after application with approximately 1/2 inch of water and leave the soil undisturbed.

Bare-Root Liners - Remove all established weeds and then irrigate with approximately 1/2 inch of water after planting to settle the soil around the root systems. Broadcast Double O E-Pro Herbicide evenly over the field growing areas one (1) week after planting. Immediately after application, irrigate with approximately 1/2 inch of water and leave the soil undisturbed. Prior to re-application of Double O E-Pro Herbicide, loosen the soil by light cultivation.

LANDSCAPE ORNAMENTALS

IMPORTANT: Double O E-Pro Herbicide should be applied only to those species listed on the Double O E-Pro Herbicide label. Double O E-Pro Herbicide is NOT registered for use on vegetables or edible fruit bearing crops, turf, bedding plants, or flowering crops except Protea and Salvia species.

Apply Double O E-Pro Herbicide either immediately after planting, in the spring prior to weed germination, or following complete weed removal using a rate of 100 lbs. per acre. Immediately after application irrigate with approximately 1/2 inch of water and leave the soil undisturbed.

Do not reapply at less than two (2) month intervals and do not make more than two (2) applications per growing season.

CUT FLOWER AND FOLIAGE CROPS

Apply Double O E-Pro Herbicide prior to weed germination or following complete weed removal. To settle the soil around the root systems, after planting and prior to application irrigate with approximately 1/2 inch of water and then allow foliage to dry. Broadcast Double O E-Pro Herbicide evenly over the production area and irrigate immediately after application to activate the herbicide.

Wait three (3) weeks before applying to bare root plantings and do not reapply at less than three (3) month intervals.

NOTE: Be sure to follow all use precautions listed in this label unless specifically exempted in the specific crop instructions below.

FIELD GROWN PROTEA

Cover any shallow feeder roots exposed by cultivation or erosion prior to application. Apply Double O E-Pro Herbicide to field plantings of 1-2 year Protea nursery stock at a rate of 100 lbs. per acre. For soils with low organic matter levels (0.5-2%), apply Double O E-Pro Herbicide at a rate of 50 lbs. per acre. Reapplication may be made throughout the production cycle at three (3) month intervals.

SARAN HOUSE GROWN ANTHURIUMS

Make certain that all plant tissue (leaves / stems) are completely dry prior to application. Using a granular spreader that allows for direct placement of Double O E-Pro Herbicide to the base of plants and avoiding contact with leaf petiole attachments and flowering parts, apply Double O E-Pro Herbicide at a rate of 100 lbs per acre being sure to treat no more than 1/4 (25%) of any saran production unit in a single day. Reapplication may be made at no less than three (3) month intervals.

NOTE: To avoid personal injury due to inhalation of Double O E-Pro Herbicide vapors, be sure to provide optimum ventilation to all treated areas.

DRACAENA FIELD STOCK

When establishing new field stock plantings, bind the foliage of unrooted cuttings with rubber bands or string to prevent the Double O E-Pro Herbicide granules from lodging in the leaf whorls (see USE

GUIDELINES AND PRECAUTIONS). Using a granular spreader that allows for direct placement of Double O E-Pro Herbicide to the base of plants, apply Double O E-Pro Herbicide at a rate of 100 lbs. per acre. Reapplication may be made at no less than three (3) month intervals. Apply Double O E-Pro Herbicide after cuttings have been taken from field stock. Prior to re-application the remaining side shoots (with expanded leaves) must be removed to prevent the granules of Double O E-Pro Herbicide from lodging in the leaf whorls.

STORAGE AND DISPOSAL

Do not contaminate water, fertilizer, other pesticides, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a dry place. Protect from water and extreme heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by state law, neither Etigra nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Etigra.

Double O E-Pro Herbicide is not manufactured or distributed by The Scotts Company, seller of Rout®. Rout® is a registered trademark of The Scotts Company. Etigra® is a trademark of Etigra.

EPA [approval date]