

79676-51

03/14/2007

118



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Re

Number:

79676-51

Date of Issuance:

March 14, 2007

NOTICE OF PESTICIDE:

☒ Registration☐ Reregistration

(under FIFRA, as amended)

Term of Issuance: conditional

Name of Pesticide Product:

ETI 106 01 I-C

Name and Address of Registrant (include ZIP Code):

Etigra
c/o Michael Kellogg
Pyxis Regulatory Consulting, Inc.
4110 136th St, NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Per the indicated EPA Guidelines, submit the following within 18 months measured from the effective date of this registration.
 - a. one year storage stability (830.6317)
 - b. corrosion characteristics (830.6320)
3. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 79676-51."
 - b. In the Warranty section, add the phrase "To the extent consistent with applicable law, " to the beginning of the following two sentences:
 - 1) "Etigra makes no warranties ... nor any other express or implied warranty" in paragraph 3.
 - 2) "The exclusive remedy of the user or buyer ... for all claims, losses," in paragraph 4.
4. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label stamped "accepted with comments" is enclosed for your records.

Signature of Approving Official:

Thomas C. Harris
Biologist
Insecticide Rodenticide Branch
Registration Division (7505P)

Date:

MAR 14 2007

2/18

RESTRICTED USE PESTICIDE
TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR
PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES
COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

ETI 106 01 I - C
AGRICULTURAL MITICIDE/INSECTICIDE

ACTIVE INGREDIENT:
 Abamectin (CAS No.'s 65195-56-4 and 65195-55-3): 2.0%*
OTHER INGREDIENTS: 98.0%
TOTAL: 100.0%
 *1 gallon contains 0.15 lb. abamectin

KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
<p>Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness), and muscle tremors. Toxicity following accidental ingestion of ETI 106 01 I - C can be minimized by inducing vomiting within one-half hour of exposure. If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms, and measurements.</p> <p>In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin exposure.</p>	

See inside label booklet for additional PRECAUTIONARY STATEMENTS

EPA Reg. No. 79676- *LA*
 Manufactured for:
 Etigra™
 501 Cascade Pointe Lane, Suite 103
 Cary, NC 27513
 www.etigra.com

EPA Est. No.

ETI 106 01 I - C contains abamectin, the
 active ingredient used in Agri-Mek®.

Net Contents:

ACCEPTED
 with COMMENTS
 in EPA Letter Dated:
March 14, 2007
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 As amended, for the pesticide
 Registered under EPA Reg. No.
79676-51

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear (goggles or faceshield). Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

This product contains a chemical (N-methyl pyrrolidone) known to the state of California to cause birth defects or other reproductive harm.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. For more options, follow the instructions for Category B on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber \geq 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwater or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrip), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber \geq 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Do not apply this product through any type of irrigation system. Thorough coverage is essential for good mite and insect control on crops.

Do not apply ETI 106 01 I - C by aerial application in New York State.

ETI 106 01 I - C is an emulsifiable concentrate which will control certain pests on the crops listed on this label when applied according to the Directions for Use.

ETI 106 01 I - C has been tested for phytotoxicity and has a wide margin of safety on a variety of crops. ETI 106 01 I - C has also been shown to be compatible with many commonly used pesticides, crop oils and nutritional sprays. However, since it is not possible to test a large number of possible mixtures, the user should pre-test to assure the physical compatibility and lack of phytotoxicity effect of any proposed mixtures with ETI 106 01 I - C.

Spray Drift Precautions for ETI 106 01 I - C Use on Cotton, Celery, Celeriac, Cucurbits, Grapes, Head Lettuce, Pepper, Potatoes, Strawberries, and Tomatoes.

Do not apply by ground within 25 ft., or by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds; estuaries and commercial fish farm ponds.

All aerial and ground application equipment must be properly calibrated and maintained using appropriate carriers. For aerial application, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure. Sprays should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward aquatic areas. Do not cultivate within 25 ft. of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground service

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that ETI 106 01 I - C be used in a sound resistance management program which includes rotation with other products with different modes of action.

Almonds and Walnuts: For control of two-spotted spider mites, strawberry spider mites, Pacific spider mites, and European red mites (Ground Application Only).

ETI 106 01 I - C may be applied using conventional dilute or concentrated sprayers calibrated to deliver sufficient water for thorough coverage. Gallonage of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control. Apply in combination with horticultural spray oil (not a dormant oil) as directed. Applications without horticultural spray oil* may result in reduced efficacy and/or residual control. Do not graze treated orchards. To reduce the risk to aquatic environments, maintain a 25-yard buffer zone from all water bodies containing aquatic life (streams, ponds, lakes, rivers, springs, irrigation canals containing water year-round, swamps, bogs and marshes).

Use ETI 106 01 I - C in combination with horticultural spray oil (not a dormant oil) approved for use on almonds and walnuts. Apply ETI 106 01 I - C when spider mites first appear. Residual spider mite control is greater from spray deposits on newer leaves compared to older leaves. A second application may be made if necessary to maintain control.

Almonds and Walnuts

Application Rates

Pests	For Dilute Sprays ¹	For Concentrate Sprays ²
	fl. oz. ETI 106 01 I - C/100 gals.	fl. oz. ETI 106 01 I - C/A
Two spotted spider mite Pacific spider mite Strawberry spider mite European red mite	2.5-5.0	10-20
Horticultural Spray Oil*	0.25% or 1 gal./A	Minimum of 1 gal./A

* Not a dormant oil

¹ The rate of ETI 106 01 I - C per 100 gals. is based on a volume of 400 gals./A dilute spray.

² To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.

Use Restrictions- Almonds and Walnuts

- Do not exceed 20 fl. oz. of ETI 106 01 I - C per acre per application or 40 fl. oz./A in a growing season.
- Do not make more than 2 applications per growing season.

- Do not apply in less than 40 gals. of water per acre.
- If a second application is necessary, do not retreat within 21 days.
- Do not apply within 21 days of harvest.
- Do not apply by aircraft.
- Do not graze treated orchards.

Apple: For control of white apple leafhopper, tentiform leafminers, European red mites, two-spotted spider mites, and McDaniel spider mites (Ground Application Only).

ETI 106 01 I - C may be applied using conventional dilute or concentrate ground sprayers calibrated to deliver sufficient water for thorough coverage. Gallonage of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control. For best results, apply in combination with horticultural spray oil* (not a dormant oil) as directed. Application without horticultural spray oil* may result in reduced efficacy and/or residual control. Do not graze treated orchards.

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas.

Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

Do not apply ETI 106 01 I - C when weather conditions favor drift to aquatic areas.

Do not apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph.

Do not apply during a temperature inversion.

For best results, use ETI 106 01 I - C in combination with a horticultural spray oil* approved for use on apples. Apply ETI 106 01 I - C when spider mite or insect thresholds are reached. A second application may be made if necessary to maintain control.

White Apple Leafhopper (Not for use west of the Rocky Mountains):

Application of ETI 106 01 I - C is limited only to first generation white apple leafhopper. Apply soon after petalfall in combination with horticultural spray oil*.

Spider Mites: For best results, apply before a threshold of 5 spider mites per leaf is reached. Residual spider mite control with ETI 106 01 I - C in combination with Horticultural spray oil* is greater from spray deposits on newer leaves compared to older. Application of ETI 106 01 I - C for spider mite control should be limited to a period extending from petalfall through six weeks following petalfall.

Tentiform Leafminers: For best results, apply against egg (to control new hatch) and early sap feeder stages of first and second generation tentiform leafminers when locally established thresholds have been reached. Do not apply during bloom.

Apples**Application Rates**

Pests	Application Rates	
	For Dilute Sprays¹ fl. oz. ETI 106 01 I - C/100 gals.	For Concentrate Sprays² fl. oz. ETI 106 01 I - C/A
Two-spotted spider mite McDaniel spider mite European red mite Tentiform leafminers White apple leafhopper	2.5-5.0	10-20
Horticultural Spray Oil* (For best results, use oil with ETI 106 01 I - C)	0.25% or 1 gal./A	Minimum of 1 gal./A

* Not a dormant oil

¹ The rate of ETI 106 01 I - C per 100 gals. is based on a volume of 400 gals./A dilute spray.

² To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.

Use Restriction- Apples

- Do not exceed 20 fl. oz. of ETI 106 01 I - C per acre per application or 40 fl. oz./A in a growing season.
- Do not make more than 2 applications per growing season.
- Do not apply in less than 40 gals. of water per acre.
- If a second application is necessary, do not retreat within 21 days.
- Do not apply within 28 days of harvest.
- Do not apply by aircraft.
- ETI 106 01 I - C in combination with horticultural spray oil* may cause fruit injury to certain varieties of apples, e.g., russetting on light-skinned varieties such as Golden Delicious, when used alone or when other products are applied sequentially. The Directions for Use and Precautions on horticultural spray oil labels, as well as precautions on official spray guides should be followed carefully when using ETI 106 01 I - C in combination with horticultural spray oil (not a dormant oil).
- Do not graze treated orchards.

Celeriac (*Apium graveolens*): For control of two-spotted spider mites

ETI 106 01 I - C may be applied using conventional or concentrate sprayers calibrated to deliver sufficient water for thorough coverage of the foliage. For best results, use in combination with a non-ionic surfactant. Apply when mites first appear and repeat as necessary to maintain control.

Celeriac**Application Rate**

Pests	fl. oz. ETI 106 01 I - C/A
Two-spotted spider mites	16.0

Use Restrictions-Celeriac

- Do not apply at less than 7-day intervals.
- Do not exceed 48 fl. oz./A after transplanting.
- Do not apply within 7 days of harvest
- Do not apply in less than 20 gals. of water per acre.
- For resistance management purposes, use no more than 2 sequential applications.

Celery: For control of *Liriomyza* leafminers and spider mites (two-spotted and carmine)

Mix with water as indicated below and apply thoroughly by ground or air as a foliar spray to assure good upper and lower leaf coverage. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations. For spider mites, apply when mites first appear and repeat as necessary to maintain control. For leafminers, apply when adult flies are first observed and repeat application as necessary to maintain control.

Celery	Application Rate
Pests	fl. oz. ETI 106 01 I - C/A
<i>Liriomyza</i> leafminers	8.0-16.0
Spider mites (two spotted and carmine)	

Use Restrictions- Celery

- Do not apply at less than 7-day intervals
- Do not exceed 48 fl. oz./A after transplanting.
- Do not apply within 7 days of harvest
- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals. of water by air.
- For resistance management purposes, use no more than 2 applications on celery being grown for transplanting.
- Do not make more than 2 sequential applications after transplanting.
- Do not apply by aircraft in New York State.

Note: ETI 106 01 I - C may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Citrus: For control of citrus rust mites, broad mites, two-spotted spider mites, citrus bud mites, citrus leafminers, and citrus thrips.

Apply using conventional dilute or concentrate ground sprayers calibrated to deliver sufficient water for thorough coverage (or outside coverage for citrus leafminers and citrus thrips). Gallonage of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good mite and insect control.

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas.

Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

Do not apply ETI 106 01 I - C when weather conditions favor drift to aquatic areas.

Do not apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph.

Do not apply during a temperature inversion.

Applications should be made with a minimum of 0.20% horticultural spray oil (not a dormant oil) in the spray mixture or not less than 1.0 gal. of oil per acre (except where specified differently below).

Mites: Apply when mites first appear during spring, summer, and/or fall.

Citrus Bud Mites: For the best results, time the spray at "budswell."

Citrus Leafminers: Apply to protect new growth during spring, summer, or fall.

Citrus Thrips: Spray applications targeted for citrus thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun- preferably early to mid-hatch).

Citrus

Application Rates

Pests	For Dilute Sprays ¹	For Concentrate Sprays ²
	fl. oz. ETI 106 01 I - C/100 gals.	fl. oz. ETI 106 01 I - C/A
Citrus rust mite	0.5-2.0	5-20
Broad mite	1.0-2.0	10-20
Two-spotted spider mite	1.0-2.0	10-20
Citrus bud mite	1.0-2.0	10-20 ³
Citrus leafminer	—	10-20 ⁴
Citrus thrips	—	10-20 ⁵
Horticultural Spray Oil*	Minimum 0.2%	Minimum 1 gal./A

¹ The rate of ETI 106 01 I - C per 100 gals. is based on a volume of 1,000 gals./A dilute spray.

² For concentrate sprays, adjust the dosage to apply an amount per acre equal to that used in full cover dilute spray.

³ For best results, use a minimum of 500 gals./A of spray mix with a minimum of 0.5% horticultural spray oil*.

⁴ For best results, use 100-150 gals./A of spray mix applied to achieve outside coverage.

⁵ For best results, use 100-250 gals./A of spray mix applied to achieve outside coverage.

* Not a dormant oil

Use Restrictions- Citrus

- Do not apply within 7 days of harvest
- Do not apply within 30 days of last treatment
- Do not apply more than 3 applications or 40 fl. oz./A in any growing season.
- Always apply in combination with horticultural spray oil (not a dormant oil) as directed.
- For resistance management purposes, do not use in citrus nurseries.
- Do not apply by aircraft.

Cotton: For control of two-spotted spider mites, carmine spider mites, Pacific spider mites, and strawberry spider mites (West of the Mississippi river only).

ETI 106 01 I - C is an emulsifiable concentrate which, when mixed with water according to the Directions for Use will control spider mites (two-spotted, carmine, Pacific and strawberry) on cotton.

For control of spider mites on cotton, mix with water and apply at the rates given in the following table:

Cotton		Application Rates	
Spider Mites		fl. oz. ETI 106 01 I - C/A	
		Mid-season/Lay-by	Early Season
Two-spotted Carmine Pacific Strawberry		8-16	4-6

Use 8-16 fl. oz. of ETI 106 01 I - C per acre depending on size of the plant and density of the foliage. Apply with ground sprayers or aerially by fixed-wing aircraft or helicopter in a minimum of 5 gals. of water per acre to provide thorough coverage. Begin application when mites are first noticed and repeat if necessary.

For use on early season cotton, apply when cotton is under 10 inches in height.

Apply at a rate of 4-6 fl. oz. of ETI 106 01 I - C per acre by ground equipment only. Do not use less than 4 fl. oz./A.

Use Restrictions

- Do not apply more than 32 fl. oz. or more than 2 applications of ETI 106 01 I - C per acre in any growing season.
- Do not apply within 21 days of last treatment.
- Do not apply this product east of the Mississippi River.
- Do not apply within 20 days of harvest.
- Do not graze or feed cotton foliage.
- Do not apply this product through any type of irrigation system.

Cucurbits (Chayote, Chinese Waxgourd, Citron Melon, Cucumbers, Gherkin, Edible Gourds, Momordica spp., Melons, Pumpkin, Summer Squash, Winter Squash, Watermelon): For control of spider mites and leafminers.

Mix with water as indicated below and apply thoroughly by ground or air as a foliar spray to ensure good upper and lower leaf coverage. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations.

Apply when adult leafminer flies or spider mites are first observed and repeat applications as necessary to maintain control within constraints of a sound resistance management program.

Cucurbits		Application Rates
Pests		fl. oz. ETI 106 01 I - C/A
Spider mites		8.0-16.0

Pests	fl. oz. ETI 106 01 I - C/A
Leafminers	

Use Restrictions- Cucurbits

- Do not exceed 48 fl. oz./A in a growing season.
- Do not apply within 7 days of harvest.
- Do not apply at less than 7-day intervals.
- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals. of water by air.
- Do not make more than 2 sequential applications.
- Do not apply by aircraft in New York State.

Grapes: For control of western grapeleaf skeletonizer, two-spotted spider mites, Pacific spider mites, and Willamette spider mites; and for quick knockdown control of western grape leaf hoppers and variegated leaf hoppers.

ETI 106 01 I - C may be applied using conventional ground sprayers calibrated to deliver sufficient water for thorough coverage. Thorough coverage is essential for good spider mite and insect control. Always apply in combination with a nonionic surfactant to improve the wetting of the foliage and smooth out spray deposits. Do not spray alternate rows. ETI 106 01 I - C must be applied to both sides of each row for maximum coverage. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations. Do not graze treated vineyards.

Apply ETI 106 01 I - C plus a nonionic surfactant when western grapeleaf skeletonizer larvae are first observed. For optimal control, apply shortly after egg hatch. For spider mite control, apply when mites first appear, but before motiles exceed 5 per leaf.

Grapes

Pests	Application Rate fl. oz. ETI 106 01 I - C/A
Two-spotted spider mite Pacific spider mite Willamette spider mite Western grapeleaf hoppers Variegated leaf hoppers	8.0-16.0 Plus a nonionic surfactant

Note: ETI 106 01 I - C must be applied in combination with a nonionic surfactant to improve wetting of foliage and to smooth out spray deposits.

Use Restrictions- Grapes

- Do not exceed 16 fl. oz. of ETI 106 01 I - C per acre per application or 32 fl. oz./A in a growing season.
- Do not make more than 2 applications per growing season.
- If a second application is necessary, do not retreat within 21 days of first application.
- Do not apply in less than 50 gals. of water per acre.
- Do not apply within 28 days of harvest.
- Do not apply by aircraft in New York State.
- Do not graze livestock in treated vineyards.

Although ETI 106 01 I - C in combination with a nonionic surfactant has been tested for safety to grapes, it is impossible to test on all grape varieties under the variety of biotic and abiotic conditions that may

cause phytotoxicity. Therefore, Directions for Use and Precautions on nonionic labels and in official spray guides should be followed carefully when using ETI 106 01 I - C in combination with a nonionic surfactant.

Head Lettuce: For control of *Liriomyza* leafminers

Mix with water as indicated below and apply thoroughly by ground or by air as a foliar spray to assure good leaf coverage. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations. Apply when adult flies are first observed and repeat applications as necessary to maintain control.

Head Lettuce	Application Rate
Pests	fl. oz. ETI 106 01 I - C/A
<i>Liriomyza</i> leafminers	8.0-16.0

Use Restrictions- Head Lettuce

- Do not apply at less than 7-day intervals.
- Do not exceed 48 fl. oz./A per growing season.
- Do not apply within 7 days of harvest.
- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals. of water by air.
- Do not make more than 2 sequential applications.
- Do not apply by aircraft in New York State.

Note: ETI 106 01 I - C may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Hops: For control of two-spotted spider mites (Not registered for use in California).

Apply ETI 106 01 I - C when two-spotted spider mites reach treatment thresholds. Mix with water as indicated below and apply by ground equipment as a foliar spray to assure good upper and lower coverage. Gallonage of spray may vary with the size of plants and density of foliage. In any case, thorough coverage is essential for good control of two-spotted mites. When using concentrate sprays, the amount of water is reduced, but the amount of ETI 106 01 I - C per acre should remain the same as for dilute sprays. If a second application is necessary for control, do not retreat within 21 days of the first application and only after an alternative miticide with a different mode of action has been used.

Hops	Application Rate
Pests	fl. oz. ETI 106 01 I - C/A
Two-spotted spider mites	16.0

Use Restriction – Hops

- Do not use less than 16 fl. oz. per application or exceed 32 fl. oz./A per growing season.
- Do not apply within 28 days of harvest.
- Do not apply in less than 40 gals. of water per acre.
- If a second application is necessary to maintain control, do not retreat within 21 days of the first application.
- Do not apply by aircraft.
- Do not permit livestock to graze treated hop yards.

Note: ETI 106 01 I - C may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Spray Drift Precautions- Hops

All ground application equipment must be properly maintained and calibrated using appropriate carriers.

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not apply by ground equipment within 25ft. of the above mentioned aquatic areas. Avoid applications when wind direction is toward these areas.

Do not make ground applications during temperature inversions or if heavy rainfall is imminent. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Make ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity or wind gusts approach 15 mph. Inclusion of a compatible drift reducing agent is encouraged.

Low humidity and high temperatures increase the evaporative rate of droplets and therefore increase spray drift potential. When making applications under these conditions, set up equipment to produce larger droplets to compensate for evaporation.

Pears (including oriental pear trees): For control of two-spotted spider mites, McDaniel spider mites, European red mites, pear rust mites, yellow mites and pear psylla (Ground Application Only).

ETI 106 01 I - C may be applied using conventional dilute or concentrate ground sprayers calibrated to deliver sufficient water for thorough coverage. Gallonage of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control. Always apply in combination with horticultural spray oil (not dormant oil) as directed. Do not graze treated orchards.

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds:

Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas. Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row and passing tree gaps in rows.

Do not apply ETI 106 01 I - C when weather conditions favor drift to aquatic areas. Do not apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph. Do not apply during a temperature inversion.

Use ETI 106 01 I - C with a minimum of 0.25% horticultural spray oil (Not a dormant oil) in the dilute spray mixture and not less than 1.0 gal. of horticultural spray oil (Not a dormant oil) per acre in the final finished spray.

Pears

Application Rates

Pests	For Dilute Sprays ¹ fl. oz. ETI 106 01 I - C/100 gals.	For Concentrate Sprays ² fl. oz. ETI 106 01 I - C/A
Two-spotted spider mite McDaniel spider mite European red mite Pear rust mite Yellow mite	2.5-5.0	10-20
Pear psylla	2.5-5.0	10-20
Horticultural Spray oil*	0.25%	Minimum of 1 gal./A

* Not a dormant oil

¹ The rate of ETI 106 01 I - C per 100 gals. is based on a volume of 400 gals./A dilute spray.

² To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.

Use Restrictions-Pears

- Do not exceed 20 fl. oz. of ETI 106 01 I - C per acre per application or 40 fl. oz./A in a growing season.
- Do not make more than 2 applications per growing season.
- Do not apply less than 40 gals. of water per acre.
- If a second application is necessary, do not retreat within 21 days of the first application.
- Do not apply within 28 days of harvest.
- Do not apply by aircraft.
- Do not graze treated orchards.

Peppers (Includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper): For control of *Liriomyza* leafminers, *Thrips palmi*, spider mites, and broad mites.

Mix with water as indicated below and apply thoroughly by ground or air as a foliar spray to assure good upper and lower leaf coverage. For broad mites and spider mites, apply when mites first appear. For leafminers, apply when adult flies are first observed. For *Thrips palmi*, apply when thrips are first observed. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations. In all cases, repeat applications as necessary to maintain control.

Peppers

Application Rate

Pests	fl. oz ETI 106 01 I - C/A
<i>Liriomyza</i> leafminers Spider mites Broad mites <i>Thrips palmi</i>	8.0-16.0

Use Restrictions- Peppers

- Do not exceed 48 fl. oz./A per growing season.
- Do not apply at less than 7-day intervals.
- Do not apply within 7 days of harvest.
- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals. of water by air.
- Do not make more than 2 sequential applications.
- For resistance management purposes, do not use on peppers being grown for transplanting.
- Do not apply by aircraft in New York State.

Note: ETI 106 01 I - C may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Potatoes: For control of Colorado potato beetle, *Liriomyza* leafminers, and spider mites.

Mix with water as indicated below and apply thoroughly by ground or air as a foliar spray to ensure thorough coverage on the upper and lower leaf surfaces. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations. The addition of a nonionic surfactant or organosilicone-based surfactant, at the manufacturer's recommended rate is suggested for optimum spider mite and insect control. Insect and spider mite control may be reduced if ETI 106 01 I - C is used in combination with a sticker or binder type of product such as Bravo Weather Stik.

For Colorado potato beetle, do not make more than 2 applications per crop. Make the first application after approximately 50% of the egg masses have hatched and larvae are present. If 2 applications are made; they should be restricted to a single Colorado potato beetle generation per crop and should not be made at less than 7-day intervals.

For *Liriomyza* leafminers, make the first application when adult flies are first observed. Repeat applications as necessary to maintain control, but not at intervals less than 7 days.

For spider mites, make the first application when mites first appear. Repeat application as necessary to maintain control, but not at intervals less than 7 days.

Potatoes**Application Rate**

Pests	fl. oz. ETI 106 01 I - C/A
Colorado potato beetles	8.0-16.0
Spider mites	
<i>Liriomyza</i> leafminers	

Use Restriction- Potatoes

- Do not exceed 32 fl. oz. of ETI 106 01 I - C per acre per crop for spider mite or Colorado potato beetle control.
- Do not exceed 48 fl. oz. of ETI 106 01 I - C per acre per crop for leafminer control.
- Do not apply within 14 days of harvest.
- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals. of water per acre by air.
- Do not graze or feed treated foliage to livestock.
- Do not make more than 2 sequential applications.
- Do not apply by aircraft in New York State.

- Do not apply at less than 7 day intervals.

Strawberries: For control two-spotted spider mites.

Use 16 fl. oz. of ETI 106 01 I - C per acre. Make 2 applications 7-10 days apart when mites first appear. Repeat this sequence of application if necessary to maintain control.

Strawberries		Application Rate
Pests		fl. oz. ETI 106 01 I - C/A
Two-spotted spider mites		16.0

Adjust spray volume and nozzle placement to assure maximum coverage of tops and undersides of leaves.

Use Restrictions- Strawberries

- Do not exceed 16 fl. oz./A per application or 64 fl. oz./A in a growing season.
- Do not apply within 3 days of harvest.
- For resistance management purposes, do not use in strawberry nurseries.
- Do not apply in less than 100 gals. of water per acre.
- Do not repeat treatment within 21 days of second application.
- Do not apply by aircraft.

Note: ETI 106 01 I - C may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Tomatoes: For control of Colorado potato beetle, *Liriomyza* leafminers, spider mites, russet mites, and tomato pinworm.

Mix with water as indicated below and apply thoroughly by ground or air as a foliar spray to assure good upper and lower leaf coverage. For Colorado potato beetle, apply when newly emerged larvae are present. For russet and spider mites, apply when mite damage first appears. For leafminers, apply when adult flies are first observed. For tomato pinworm, apply when larvae are first observed. In all cases, repeat applications as necessary to maintain control. Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations.

Tomatoes		Application Rates
Pests		fl. oz ETI 106 01 I - C/A
Colorado potato beetles <i>Liriomyza</i> leafminers Spider mites Tomato russet mite		8.0-16.0
Tomato pinworm		16.0

Use Restrictions – Tomatoes

- Do not exceed 48 fl. oz./A per growing season
- Do not apply at less than 7-day intervals
- Do not apply within 7 days of harvest

- Do not apply in less than 20 gals. of water per acre by ground or less than 5 gals of water by air.
- Do not make more than 2 sequential applications.
- For resistance management purposes, do not use on tomatoes being grown for transplanting.
- Do not apply by aircraft in New York State.

Note: ETI 106 01 I – C may be used without a wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPILLS: For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-424-9300, day or night.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. **ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Etigra nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Etigra.

ETI 106 01 I – C is not manufactured or distributed by Syngenta, seller of Agri-Mek®.
Agri-Mek® is a trademark of a Syngenta Group Company.
Etigra™ is a trademark of Etigra.