



U.S. ENVIRONMENTAL PROTECTION
AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg.

Number:

79676-53

Date of Issuance:

APR 13 2007

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance: **Conditional**

Name of Pesticide Product:

ETI 103 01 G

Name and Address of Registrant (include ZIP Code):

Etigra
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit within one year of the date of this registration notice, storage stability (830.6317) and corrosion characteristics (830.6320) studies.
3. Make the following label changes:
 - a. Change the Engineering Controls Statements to read "Pilots must use an enclosed cockpit that meets the requirements listed in the WPA for agricultural pesticides [40 CFR 170.240(d)(6)]."
 - b. The second sentence for User Safety Requirements must read "If no such instructions for washables exist, use detergent and hot water."
 - c. Add the following statement to the User Safety Recommendations statements, "Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

APR 13 2007

- d. Under Environmental Hazards add the following statements:
“This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product’s contribution to surface water contamination.”
“This chemical has properties and characteristics associated with chemicals detected in ground water. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.”
- e. In the Agricultural Use Requirements box, in the Early Reentry Personal Protective Equipment for products, the second bullet must read “Chemical-resistant gloves made of any waterproof material”.
- f. The following label recommendations must be reflected in the directions for use:
“General Use Rate Information (under normal conditions) for Cotton:
0.125 lbs ai/A
Maximum application rate of 0.3 lbs ai/A/year
Maximum of 2 applications, but not to exceed 0.3 lbs/A/year.
- General Use Rate Information (under variable conditions) for cotton:
Maximum application rate of 0.2 lbs ai/A
Maximum application rate of 0.3 lbs/A/year
Maximum of 2 applications, but not to exceed 0.3 lbs ai/A/year.
- For cutout and mature cotton under normal weather patterns, use (insert product name) at 0.05 to 0.10 lb. active ingredient per acre. The 0.05 lb. rate is most effective when used in a tank mix with other cotton defoliant products.
- Increased rates of (insert product name) above 0.10 lb. of active ingredient per acre may be needed to defoliate and control regrowth during periods of rank growth/high fertilizer conditions, extreme weather conditions, such as extended periods of rain and/or low temperatures (60 -65 F), and on full-season cotton varieties.”

page 3

EPA Reg. No. 79676-53

To Achieve an Application Rate of:	Use This Amount of Product	At the Indicated Rate, One Gallon of Product Will Treat:
0.07 lbs. ai/Acre	2.3 fluid oz/A	56 Acres
0.17 lbs. ai/Acre	5.4 fluid oz/A	23.5 Acres

- g. Revise the EPA Registration Number to read, "EPA Reg. No. 79676-53".
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

4 8 11
ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 13 2007
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

ETI 103 01 G

Cotton Defoliant - For Agricultural Use Only

ACTIVE INGREDIENT:		
Thidiazuron (N-phenyl-N'-1,2,3-thiadiazol-5-ylurea)*	42.4%	79676-53
OTHER INGREDIENTS:	57.6%	
TOTAL:	100.0%	

*Contains not less than 4.0 lbs. of thidiazuron per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 79676-26

EPA Est. No.

Manufactured for:
 Etigra™
 501 Cascade Pointe Lane
 Suite 103
 Cary, NC 27513
 www.etigra.com

ETI 103 01 G contains thidiazuron,
 the active ingredient used in Dropp®.

Net Contents:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or butyl rubber gloves

- Chemical-resistant footwear plus socks
- Protective eyewear

GENERAL INFORMATION

ETI 103 01 G provides growers a tool to help in the harvest of cotton. ETI 103 01 G readily disperses in water for use in various types of application equipment. Applications by ground or air of ETI 103 01 G just prior to harvest causes the leaves to fall off the plants (defoliation) and inhibits additional plant growth. This effect helps to reduce any staining of lint that may occur during harvest and ginning.

ETI 103 01 G's effects may not be seen for several days after treatment. Cooler weather or other adverse conditions may require an adjustment in the dose level (i.e., higher rates) or require longer periods of time for defoliation to occur.

USE PRECAUTIONS

- Do not apply ETI 103 01 G through any type of irrigation system.
- To avoid inconsistent or ineffective results, do not apply ETI 103 01 if rainfall is expected within 24 hours of application.
- Do not feed gin trash, treated foliage or immature crops to livestock.

Rotation Crop Restrictions: Do not replant the following crops unless the minimum specified time has elapsed after the last application of ETI 103 01 G.

ROTATION CROP	PLANTBACK INTERVAL
Small grains, sorghum, corn, or root crops (except carrots, onions or sugar beets)	Two (2) weeks
Legumes (including alfalfa) or leafy vegetables (except lettuce and spinach)	Two (2) months
Sugar beets	Four (4) months
Carrots, onions and spinach	Nine (9) months
Lettuce (replant ONLY if the soil has been deep-plowed down to 12-15 inches.)	Nine (9) months

ETI 103 01 G will defoliate and cause injury to other plant species. Avoid spray drift to crops other than the target crop of cotton.

Increased non-target phytotoxicity may be observed when tank-mixing ETI 103 01 G with organophosphates or when particularly sensitive crops such as lettuce, citrus, and cantaloupes are planted adjacent to fields treated with ETI 103 01 G.

STEPS TO REDUCE SPRAY DRIFT POTENTIAL - NON-TARGET CROPS

Drift to non-target crops may occur during application of ETI 103 01 G especially under windy weather conditions. Take the following steps to reduce the potential for off-site spray drift:

- Do not make aerial or ground applications of ETI 103 01 G when wind speeds are greater than ten (10) miles per hour at the application site.
- Use the recommended low nozzle pressures of 20-30 psi.
- Do not apply this product during a temperature inversion or if a temperature inversion appears likely to occur during the application.
- Do not make ground applications of ETI 103 01 G within 100 feet of growing lettuce.
- Do not make aerial applications of ETI 103 01 G within 0.5 miles of growing lettuce.

STEPS TO REDUCE SPRAY DRIFT POTENTIAL - NON-TARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

The following steps can help reduce the potential for drift:

- Do not make aerial or ground applications of ETI 103 01 G when wind speeds are greater than ten (10) miles per hour at the application site. If wind speeds are less than ten (10) miles per hour, consult your State Extension Cotton Specialist for local application recommendations.
- Use the recommended low nozzle pressures of 20-30 psi.
- Do not apply this product during a temperature inversion or if a temperature inversion appears likely to occur during the application.
- Use the maximum nozzle orifice size that provides appropriate coverage and deposition of this product.
- **Citrus Areas:** Do not make ground applications of ETI 103 01 G within 0.5 miles of citrus in flush. Do not make aerial applications of ETI 103 01 G within 0.5 miles downwind of citrus in flush.

APPLICATION TIMING

Make applications of ETI 103 01 G defoliant only to mature cotton plants and when bolls are mature. Bolls are considered mature when the following are observed: 1) the boll does not dent when squeezed between thumb and finger; and/or 2) the boll is difficult to cut using a sharp knife; and/or 3) the seeds have fully developed cotyledons (determined by the fact that the seeds lack liquid within the seed coat if cut in cross section or if the outer edges of the seed coat are light brown).

ETI 103 01 G applications may be made at least five (5) days before the expected day of harvest.

Note: The success of ETI 103 01 G as a cotton defoliant depends on the following factors

- Cotton plants are actively growing.
- Humid weather conditions exist.
- Cotton plant leaves are high in moisture content

APPLICATION CONDITIONS

If the temperatures at night fall below 60°F, ETI 103 01 G applied as a stand-alone product (i.e., without a tank mix partner), may give inconsistent or ineffective defoliation and/or regrowth suppression. Improved performance of ETI 103 01 G is achieved if application is timed so that the overnight temperatures exceed 60°F for 2-3 days prior to and 2-3 days after application.

Heat- and/or drought-stressed cotton (low leaf moisture content, thick cuticle) may not be completely defoliated and/or suppressed in growth when ETI 103 01 G is applied to these plants.

SEQUENTIAL ETI 103 01 G APPLICATIONS

To avoid reduced or unsatisfactory defoliation, apply a sequential application of ETI 103 01 G **only** if the first application was ETI 103 01 G as a stand-alone or as a tank mix with other defoliants. Also refer to the section on **APPLICATION CONDITIONS** of this label.

USE OF CROP OIL/ADJUVANTS

Improved defoliation and regrowth suppression has been shown when crop oil concentrates (COC's) or penetrating oils approved for use on agricultural crops are mixed with ETI 103 01 G and applied when nighttime temperatures are low (60°-65°F) or when drought conditions exist. Consult the adjuvant label for use directions and rates.

When applying ETI 103 01 G in the desert Southwest (i.e., Arizona and California), use of adjuvants is recommended.

NOTE: The use of adjuvants is not recommended during high temperature periods and may result in the freezing and/or desiccation of leaves.

MIXING INSTRUCTIONS

1. Add one-half the amount of water required to the spray applicator tank.
2. Begin agitation and continue agitation during the addition and application steps below.
3. Add ETI 103 01 G.
4. Fill the spray tank with the remainder of the water.

Prepare only the amount of spray solution needed for a single day of application. Do not allow the spray tank mixture to stand overnight.

TANK MIX ADJUVANTS

To improve defoliation and spray tank clean-out, addition of a compatibility agent or surfactant to tank mixes of ETI 103 01 G with organophosphate defoliants is recommended. Conduct a compatibility test of all components before adding to the spray tank.

Only products which are exempt from tolerance (refer to 40 CFR 180.1001) are permitted for this use.

Consult the adjuvant label for mixing and application directions and use rates.

DIRECTIONS FOR APPLICATION

Apply the recommended amount of ETI 103 01 G in sufficient volume of water to ensure uniform and thorough coverage of the leaves. Ground equipment: apply 10-25 gallons per acre. Aerial applications: apply 3-10 gallons per acre.

Ensure adequate agitation is maintained during application.

APPLICATION USE RATES

Apply ETI 103 01 G at rates between 0.1 lb AI/A (3.2 fl. oz./A) and 0.2 lb. AI/A (6.4 fl. oz./A): at these rates, one gallon of product treats between 40 and 20 acres, respectively. Use the higher rate if application is made when temperatures are between 60°-65°F.

For defoliation of rank cotton, a second application may be required to ensure adequate defoliation. Do not make more than two applications and do not exceed a total of 9.6 fl. oz. (0.3 lb. AI/acre).

DO NOT APPLY MORE THAN 9.6 FL. OZ. (0.3 LB. AI/ACRE) PER SEASON.

ARIZONA ONLY- BOTTOM DEFOLIATION

Use ETI 103 01 G to improve air circulation within the cotton canopy and reduce boll rot and hard lock caused by moisture under the canopy.

Time the date of application so that application equipment can enter the fields without causing damage to the cotton crop.

Apply 0.05 lb. AI/acre (1.6 fl. oz. ETI 103 01 G/A); direct the sprays to the lower 1/3 part of the plants.

ARIZONA AND CALIFORNIA ONLY: PRECONDITIONING

ETI 103 01 G can be applied to pre-condition the cotton seven to ten days prior to application of other defoliant compounds and results in improved efficacy of the defoliants. Consult the label of the other defoliant prior to use, and observe all product label directions, recommendations and prohibitions.

Apply ETI 103 01 G at rates between 0.05 lb AI/A (1.6 fl. oz./A) and 0.1 lb. AI/A (3.2 fl. oz./A).

TANK MIXES

ETI 103 01 G PLUS ETHEPHON-CONTAINING COMPOUNDS – BOLL OPENING

Use of ETI 103 01 G and Ethephon 6 or its equivalents will improve defoliation and speed the opening of mature bolls. For best results, apply to mature cotton plants (refer to the **APPLICATION TIMING** section of this label). Ensure a sufficient number of bolls have matured prior to application in order to achieve good cotton yields. Certified cottonseed producers should consult seed companies to determine the boll and seed maturity requirements for their varieties. Reduced defoliation or suppression of regrowth may occur if applications are made when nighttime temperatures are expected to be below 60°F.

Apply ETI 103 01 G at rates between 0.05 lb. A/A (1.6 fl. oz./A) and 0.2 lb. A/A (6.4 fl. oz./A) plus 1-2 lbs. A/acre (32-64 fl. oz./A) of Ethephon 6.

The higher rates of ETI 103 01 G are used when excessive regrowth is expected or if temperatures are between 60-65°F.

ETI 103 01 G PLUS ETHEPHON - ENHANCED DEFOLIATION

Use the lower rates of Ethephon 6 or its equivalents plus ETI 103 01 G to enhance defoliation without affecting boll opening. After this tank mix application, use Ethephon 6 alone to promote boll opening but do not apply more than 2.0 lb. A/acre of Ethephon 6 (approximately 42 fl. oz. of product) for all ethephon applications to cotton. Read and follow the Ethephon 6 label directions, recommendations and prohibitions.

Apply ETI 103 01 G at rates between 0.05 lb. A/A (1.6 fl. oz./A) and 0.2 lb. A/A (6.4 fl. oz./A) plus 0.25 lbs. A/acre (32-64 fl. oz./A) of Ethephon 6. The higher rates of ETI 103 01 G are used when excessive regrowth is expected or if temperatures are between 60-65°F.

ETI 103 01 G PLUS TRIBUFOS PRODUCTS (DEF[®]/FOLEX[®]) – NOT FOR USE IN CAMERON, HIDALGO, STARR AND WILLACY COUNTIES OF THE STATE OF TEXAS

If plants are treated when growing under less favorable conditions, such as when overnight temperatures are expected to be below 60°F one day prior to and three days after application, ETI 103 01 G and DEF[®] 6/ FOLEX[®] 6 EC tank mixes are recommended for improved regrowth suppression and defoliation. For optimum results, apply the tank mix to cotton plants that have at least 60% open bolls.

Apply ETI 103 01 G at rates between 0.05 lb. A/A (1.6 fl. oz./A) and 0.2 lb. A/A (6.4 fl. oz./A) plus DEF[®] 6 at 0.5- 2.0 pints per acre.

Apply ETI 103 01 G at rates between 0.05 lb. A/A (1.6 fl. oz./A) and 0.2 lb. A/A (6.4 fl. oz./A) plus FOLEX[®] 6 EC at 0.5-2.0 pints per acre.

The higher rates of ETI 103 01 G are used when excessive regrowth is expected.

The lower rates of DEF[®] 6/ FOLEX[®] 6 EC are used when nighttime temperatures are expected to be 65°F or above. Use the higher rates if conditions are not ideal.

If a second application of this tank mix is necessary, do not exceed a total of 0.3 lb. A/acre (9.6 fl. oz./A) of ETI 103 01 G.

Read and follow the DEF[®] 6 and FOLEX[®] 6 EC label directions, recommendations and prohibitions.

NOTE: Use caution when tank mixes of phosphate defoliant plus ETI 103 01 G are applied at the highest use rates and during extremely high temperatures since leaf desiccation may occur.

Preparation of Tank Mixes of ETI 103 01 G plus DEF[®] 6/ FOLEX[®] 6 EC

Use the directions below to prepare tank mixes of ETI 103 01 G plus DEF[®] 6/ FOLEX[®] 6 EC:

1. Add one-half the amount of water required to the spray applicator tank.
2. Begin agitation and continue through addition and application steps.
3. Add the required amount of ETI 103 01 G and agitate to ensure complete mixing with water.
4. Add the required amount of DEF[®] 6 or FOLEX[®] 6 EC to the spray tank.

SPRAY TANK CLEANOUT INSTRUCTIONS

IMPORTANT: Residues may form in the spray equipment of tank mixes of ETI 103 01 G plus organophosphates. The residues can be removed using a commercial tank cleaner as long as the spray equipment is cleaned while the residue is still fresh and moist. **DO NOT ALLOW SPRAY EQUIPMENT TO DRY BEFORE CLEANING.** Consult your State Extension Cotton Specialist for recommended tank cleaners and cleaning procedures.

Immediately after application, flush the entire system (nozzles, booms, application tanks, sumps, pumps, and transfer lines, etc.) using a tank cleaner and water.

Should the spray solution be allowed to dry in the application equipment, the residues will be more difficult to remove. Use the commercial tank cleaner diluted with water to completely fill the equipment and let it stand for 7 days. Then, completely flush and rinse all equipment with water.

If spray application equipment is not thoroughly cleaned, residues of ETI 103 01 G may remain and be released during subsequent applications and lead to crop damage.

Etigra assumes no liability for crop damage due to spray equipment that has not been properly cleaned before re-use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store product at temperature less than 100°F. Re-mix product prior to use if product is exposed to freezing conditions.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. **ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, neither Etigra nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Etigra.

ETI 103 01 G is not manufactured or distributed by Bayer CropScience, seller of Dropp®.
DEF® and Dropp® are trademarks of Bayer.
FOLEX® is a trademark of AMVAC Chemical Corporation.
Etigra™ is a trademark of Etigra.