

79676-54

10/16/2008

Y12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

OCT 16 2008

Michael Kellogg
Agent for Etigra
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

SUBJECT: Application for Pesticide Notification – Removal of Statement that is Not
Applicable for Product
Imazuron E-Pro Herbicide
EPA Reg. No. 79676-54
Application Dated October 16, 2007

Dear Mr. Kellogg:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

3/12

PYXIS REGULATORY CONSULTING, INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
Email: www.PyxisRC.com

October 16, 2007

COURIER DELIVERY

Jim Tompkins (PM25)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202-4501

RE: Etigra – Imazuron E-Pro Herbicide (EPA Reg. No. 79676-54)
Notification of a Minor Label Revision per PRN 98-10

Dear Mr. Tompkins,

On behalf of Etigra please find the enclosed notification of a minor label revision for Etigra's Imazuron E-Pro Herbicide (EPA Reg. No. 79676-54). The statement "Application with a spoon, a pump-feed backpack spreader or a gravity-feed backpack spreader is PROHIBITED" has been removed from the USE PRECAUTIONS section of the label as these methods of application imply the product is being applied as a granule. Imazuron E-Pro Herbicide is a dispersible granule that is **mixed with water** and a spray adjuvant and applied as a **spray solution**. Therefore, application with a spoon, backpack spreader or gravity-feed backpack spreader is not applicable. No other changes to the labeling have been made.

In support of this notification, we submit the following documents:

1. Completed Application for Registration (EPA Form 8570-1)
2. One (1) copy of the Imazuron E-Pro Herbicide label with changes tracked
3. One (1) copy of the Imazuron E-Pro Herbicide label with changes incorporated
4. Letter of Authorization

Please feel free to call me if you have any questions or need any additional information.

Sincerely,



Michael Kellogg

Enclosures



4/12

NOTIFICATION

OCT 16 2008

Imazuron E-Pro Herbicide

A broad-spectrum dispersible granule herbicide for use on specified non-crop sites and under paved areas.

ACTIVE INGREDIENTS:

Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)	7.78%
Diuron (3-[3,4-dichlorophenyl]-1,1-dimethylurea).....	62.22%
OTHER INGREDIENTS:	30.00%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

EPA Reg. No. 79676-54

EPA Est. No.

Manufactured for:

Etigra™
501 Cascade Pointe Lane, Suite 103
Cary, NC 27513
www.etigra.com

Imazuron E-Pro Herbicide contains imazapyr and diuron, the active ingredients used in Sahara®.

Net Weight:

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, or viton ≥ 14 mils. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- Long sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material ≥ 14 mils
- NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix "TC-84A". Groundboom applicators do not need to wear a respirator.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of Imazuron E-Pro Herbicide should be mixed, stored and applied only in stainless steel, fiberglass, plastic, and plastic-lined steel containers. DO NOT mix, store or apply Imazuron E-Pro Herbicide or spray solutions of Imazuron E-Pro Herbicide in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE PRECAUTIONS

- Do NOT enter or allow others to enter treated areas until sprays have dried.
- Aerial application is prohibited.
- DO NOT USE IN CALIFORNIA.
- Do NOT apply more than a total of 12 lbs. a.i. of diuron (19 pounds per acre of Imazuron E-Pro Herbicide) in a 12-month period.

- DO NOT mix, store or apply Imazuron E-Pro Herbicide or spray solutions of Imazuron E-Pro Herbicide in unlined steel (except stainless steel) containers or spray tanks.
- DO NOT use on food or feed crops.
- DO NOT treat irrigation ditches, or water used for crop irrigation or for domestic purposes.
- Do NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- DO NOT contaminate water when disposing of equipment washwaters.
- DO NOT drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- DO NOT use on turfgrass at residential sites (including homes, apartment complexes, condominium grounds, daycare facilities, schools, playgrounds, parks, recreational areas, and sports fields).
- DO NOT side trim desirable vegetation with this product.
- DO NOT allow Imazuron E-Pro Herbicide to come in contact with other fertilizers, insecticides, fungicides and seeds.
- Take all measures possible to prevent drift of spray to desirable plants.
- Be sure to clean application equipment after using this product by thoroughly flushing with water.

GENERAL INFORMATION

Imazuron E-Pro Herbicide is a dispersible granule that is mixed with water and a spray adjuvant and applied as a spray solution to the following noncropland areas where bare ground is desired: industrial non-crop areas including utility plant sites, petroleum tank farms, pumping installations, storage areas, railroads, utility, and pipeline rights-of-way; highway rights-of-way; non-irrigation ditchbanks; fence rows; farmyards; and non-crop areas around farm buildings. Imazuron E-Pro Herbicide may also be used to control weeds under paved surfaces.

Imazuron E-Pro Herbicide controls most annual and perennial grasses and broadleaf weeds in addition to many brush and vine species. Imazuron E-Pro Herbicide also provides residual control of weeds that germinate in treated areas.

For annual weed control, either preemergence or postemergence applications may be used; however, a late preemergence to early postemergence application provides the best results in most situations.

For perennial weed control, Imazuron E-Pro Herbicide is only effective when applied postemergence and will not control perennial weeds that have not emerged at the time of application. For best results, applications should be made when the weeds are growing vigorously and the spray solution should include a spray adjuvant. For specific recommendations, see the "Adjuvants" section of this label.

The duration of residual weed control depends upon the types of weeds present, the application rate, and weather conditions. Longer residual control occurs in areas with sensitive weed species, higher Imazuron E-Pro Herbicide use rates, lower precipitation and cooler soil temperatures. Higher than average rainfall or warmer than normal temperatures can significantly affect the residual control Imazuron E-Pro Herbicide provides and shorten the overall length of control.

Precautions for Avoiding Injury to Non-Target Plants

Untreated trees may be affected by root uptake of Imazuron E-Pro Herbicide through movement into the topsoil and injury or loss of desirable trees or other plants may result if Imazuron E-Pro Herbicide is applied on or near desirable trees or other plants, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots. Treatment of powdery dry soil or light sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to desirable plants when soil particles are moved by water and/or wind. Exposure to Imazuron E-Pro Herbicide may injure or kill most crops and injury to crops may result if treated soil is washed, blown or moved onto land used to produce crops.

Use best practices to avoid drift to all other crops and non-target areas. Do not apply when conditions favor drift from target areas. The interaction of many equipment-and weather-related factors determine

the potential for spray drift. Avoiding spray drift at the application site is the responsibility of the applicator. The applicator must follow the most restrictive precautions to avoid drift, including those found in this labeling as well as applicable state and local regulations and ordinances. A drift control agent may reduce drift, however, it may also decrease weed control.

Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and Coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater.

Apply only when wind speed is less than or equal to 10 miles per hour.

Do not apply into temperature inversions.

When applying to non-crop areas, use lowest nozzle height consistent with safety and efficacy. Direct spray into target vegetation.

APPLICATION INSTRUCTIONS

For rights-of-way and non-crop areas:

- The maximum rate per application is 19 lbs./acre Imazuron E-Pro Herbicide (equivalent to 12 lbs. diuron active ingredient per acre) in areas of high rainfall or dense vegetation.
- For all other areas, the maximum rate per application is 13 lbs./acre Imazuron E-Pro Herbicide (equivalent to 8 lbs. diuron active ingredient per acre).
- Make a maximum of two applications per year.
- The minimum retreatment interval is 90 days.

Mix Imazuron E-Pro Herbicide in water and apply the recommended gallons per acre of spray volume using properly calibrated equipment to deliver a uniformly distributed spray pattern. Imazuron E-Pro Herbicide should be applied at 7-19 pounds of product per acre, although rates as low as 5 pounds per acre may be used *only* if tank mixed with another herbicide (see the TANK MIXES section below). For retreatment purposes within the same growing season, Imazuron E-Pro Herbicide may be applied at a rate of less than 7 pounds per acre.

Rainfall may significantly affect length of residual weed control achieved with Imazuron E-Pro Herbicide and in cases of increasing rainfall amounts, higher rates may need to be applied to achieve the desired residual control. Refer to the following table for Imazuron E-Pro Herbicide rates for different annual rainfall amounts. Actual use rates will depend upon the length of residual control desired as well as weed pressure and environmental conditions.

Average Annual Rainfall	Imazuron E-Pro Herbicide
Less than 15 inches	7-10 pounds [†]
Between 15 and 35 inches	8-13 pounds
Greater than 35 inches	13-19 pounds

[†] Initial applications of Imazuron E-Pro Herbicide may be made at rates as low as 5-6 pounds per acre, but must be tank mixed with another herbicide (see the TANK MIXES section below).

When both mixing and spraying, be sure to maintain sufficient agitation to keep the Imazuron E-Pro Herbicide suspended in spray mixture.

Postemergence Applications: When making postemergence applications, always use a spray adjuvant (see "Adjuvants" section of this label). For best results on tough to control perennial weeds, applications should be made in combination with one quart per acre of methylated seed oil and at a total volume of 100 gallons per acre or less. For faster burndown or brown-out of target weeds, Imazuron E-Pro Herbicide may be tank-mixed with products such as Roundup[®], or Finale[®] (refer to the TANK MIXES section for specific recommendations).

Tank Mixes

Imazuron E-Pro Herbicide may be tank-mixed with Roundup[®], Karmex[®] (Diuron), Oust[®], Garlon[®], Finale[®], MSMA, Banvel[®], Vanquish[®], Pendulum[®], Plateau[®], or Arsenal[®]. Do not use a tank mix product if the tank mix product label prohibits such mixing. Consult the manufacturer's labels for specific rates and weeds controlled, and always follow the more restrictive label instructions and restrictions on all labels used when making a tank-mix application.

NOTE: Tank-mixes with 2,4-D or products that contain 2,4-D have resulted in reduced perennial weed control performance.

For Control of Undesirable Weeds under Paved Surfaces

Imazuron E-Pro Herbicide should be applied only to areas that have been prepared according to good construction practices. All rhizomes, stolons, tubers, or other vegetative plant parts present in the site should be removed by scalping with a grader blade to a depth sufficient to ensure their complete removal.

Imazuron E-Pro Herbicide may be applied under asphalt, pond liners and other paved areas *only* in industrial sites or where the pavement has a barrier along the perimeter that will prevent encroachment of roots of desirable plants. Sites should be paved as soon as possible after application.

NOTE: This product is not recommended for use where landscape plantings could be anticipated, or under paved areas such as driveways or parking lots on residential properties, nor is it recommended for use in recreational areas such as under bike or jogging paths, golf cart paths, or tennis courts. Injury or death of desirable plants may result if this product is applied where roots are present or where they may extend into the treated area.

Application Directions for Paved Surfaces:

When final grade is established, apply Imazuron E-Pro Herbicide in sufficient water (at least 100 gallons per acre) to ensure thorough and uniform wetting of the soil surface, including the shoulder areas. Do not move soil following application. In the spray tank, mix clean water with 19 pounds of product per acre during the filling operation and be sure to agitate before spraying.

Incorporation of Imazuron E-Pro Herbicide is required to activate the herbicide if the soil is not moist prior to treatment. Use a rototiller or disc to incorporate the Imazuron E-Pro Herbicide into the soil to a depth of 4 to 6 inches. One inch of rainfall or irrigation will also provide uniform incorporation. If using water to incorporate, do not allow treated soil to wash or move into untreated areas.

Adjuvants

Nonionic Surfactants: A nonionic surfactant at a rate 0.25% v/v or higher (see manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons) may be used with this product. For best results, select a nonionic surfactant with a HLB (hydrophilic to lipophilic balance) ratio between 12 - 17 and that has at least 70% surfactant in the formulated product (alcohols, fatty acids, oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet the above requirements).

Methylated Seed Oils or Vegetable Oil Concentrates: Methylated seed oils are the adjuvant of choice and research indicates that these oils may aid in Imazuron E-Pro Herbicide deposition and uptake by plants under moisture or temperature stress and will increase control of perennial weeds. Use a methylated seed oil or vegetable-based seed oil concentrate at the rate of 1.5 - 2 pints per acre in place of a surfactant. When using spray volumes greater than 30 gallons per acre, methylated seed oil or vegetable based seed oil concentrates should be mixed at a rate of 1% of the total spray volume, or a nonionic surfactant as described above may be used instead.

Silicone-Based Surfactants: Silicone-based surfactants may allow greater spreading on the leaf surface as compared to conventional nonionic surfactants by reducing the surface tension of the spray droplets. However, some silicone-based surfactants may limit herbicide uptake by drying too rapidly. Refer to the manufacturer's label for specific rate recommendations.

Fertilizer/Surfactant Blends: Use 2 - 3 pints of nitrogen-based liquid fertilizers (such as 28%N, 32%N, 10-34-0, or ammonium sulfate) per acre in combination with the recommended rate of nonionic surfactant, methylated seed oil or vegetable/seed oil concentrate. Using fertilizers in a tank mix without a nonionic surfactant, methylated seed oil or vegetable/seed oil concentrate is not recommended.

Weeds Controlled by Imazuron E-Pro Herbicide

When used at the rates listed in the APPLICATION INSTRUCTIONS section, Imazuron E-Pro Herbicide provides preemergence or postemergence control with residual control (control of newly germinating seedlings) of the following target vegetation species. In general, annual weeds may be controlled by preemergence or postemergence applications of Imazuron E-Pro Herbicide while postemergence applications of Imazuron E-Pro Herbicide are recommended for established biennials and perennials. Imazuron E-Pro Herbicide should be used only in accordance with the recommendations on this label.

Note Regarding Resistant Biotypes: Naturally occurring biotypes of some of the weeds listed on this label (pigweed, kochia and Russian thistle for example) may not be effectively controlled by this and/or other herbicides with the ALS/AHAS enzyme inhibiting mode of action (such as OUST). To ensure control if naturally occurring ALS/AHAS resistant biotypes are present in an area, Imazuron E-Pro Herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action.

WEEDS CONTROLLED¹

GRASSES			
Name (Species)	Growth Habit ²	Name (Species)	Growth Habit ²
Annual bluegrass (<i>Poa annua</i>)	A	Kyllinga (<i>Cyperus brevifolius</i>)	A
Annual ryegrass (<i>Lolium multiflorum</i>)	A	Lovegrass (<i>Eragrostis</i> spp.)	A/P
Annual sweet vernalgrass (<i>Anthoxanthum odoratum</i>)	A	Maidencane (<i>Arundinaria amabilis</i>)	P
Bahiagrass ⁷ (<i>Paspalum notatum</i>)	P	Orchardgrass (<i>Dactylis glomerata</i>)	P
Barnyardgrass (<i>Echinochloa crusgalli</i>)	A	Paragrass (<i>Brachiaria mutica</i>)	P
Beardgrass (<i>Andropogon</i> spp.)	P	Peppergrass (<i>Lepidium virginicum</i>)	A
Bermudagrass ^{7,8,9} (<i>Cynodon dactylon</i>)	P	Phragmites (<i>Phragmites australis</i>)	P
Big bluestem ⁷ (<i>Andropogon gerardii</i>)	P	Prairie cordgrass (<i>Spartina pectinata</i>)	P
Broadleaf signalgrass (<i>Brachiaria platyphylla</i>)	A	Prairie threeawn (<i>Aristida oligantha</i>)	P
Canada bluegrass (<i>Poa Compressa</i>)	P	Quackgrass (<i>Agropyron repens</i>)	P
Cattail (<i>Typha</i> spp.)	P	Rattail fescue (<i>Vulpia myuros</i>)	A
Cheat (<i>Bromus secalinus</i>)	A	Reed canarygrass (<i>Phalaris arundinacea</i>)	P
Cogongrass (<i>Imperata cylindrical</i>)	P	Ricegrass (<i>Oryzopsis hymenoides</i>)	A
Crabgrass (<i>Digitaria</i> spp.)	A	Saltgrass ^{7,8,9} (<i>Distichlis stricta</i>)	P
Dallisgrass ⁷ (<i>Paspalum dilatatum</i>)	P	Sand dropseed ⁷ (<i>Sporobolus cryptandrus</i>)	P
Downy brome (<i>Bromus tectorum</i>)	A	Sandbur (<i>Cenchrus</i> spp.)	A
Fall panicum (<i>Panicum dichotomiflorum</i>)	A	Smooth brome (<i>Bromus inermis</i>)	P
Feathertop (<i>Pennisetum villosum</i>)	P	Sprangletop ^{6,7} (<i>Leptochloa</i> spp.)	A
Fescue (<i>Festuca</i> spp.)	A/P	Timothy (<i>Phleum pretense</i>)	P
Foxtail (<i>Setaria</i> spp.)	A	Torpedograss (<i>Panicum repens</i>)	P
Goosegrass (<i>Eleusine indica</i>)	A	Vaseygrass (<i>Paspalum urvillei</i>)	P
Guineagrass (<i>Panicum maximum</i>)	P	Velvetgrass (<i>Holcus lanatus</i>)	A
Italian ryegrass (<i>Lolium multiflorum</i>)	A	Wild barley (<i>Hordeum</i> spp.)	A
Johnsongrass (<i>Sorghum halepense</i>)	P	Wild oats (<i>Avena fatua</i>)	A
Kentucky bluegrass (<i>Poa pratensis</i>)	P	Wirestem muhly (<i>Muhlenbergia frondosa</i>)	P
		Witchgrass (<i>Panicum capillare</i>)	A

BROADLEAF WEEDS			
Name (Species)	Growth Habit ²	Name (Species)	Growth Habit ²
Arrowwood (<i>Pluchea sericea</i>)	A	Nettleleaf goosefoot (<i>Chenopodium murale</i>)	A
Ageratum (<i>Asteraceae houstonianum</i>)	P	Oxeye daisy (<i>Chrysanthemum leucanthemum</i>)	P
Broom snakeweed ³ (<i>Gutierrezia sarothrae</i>)	P	Pennycress (<i>Thlaspi</i> spp.)	A
Bull thistle (<i>Cirsium vulgare</i>)	B	Pepperweed (<i>Lepidium</i> spp.)	A
Burdock (<i>Arctium</i> spp.)	B	Pigweed ⁶ (<i>Amaranthus</i> spp.)	A
Canada thistle ⁷ (<i>Cirsium arvense</i>)	P	Pineapple weed (<i>Matricaria matricarioides</i>)	P

BROADLEAF WEEDS			
Name (Species)	Growth Habit ²	Name (Species)	Growth Habit ²
Carolina geranium (<i>Geranium carolinianum</i>)	A	Plantain (<i>Plantago</i> spp.)	P
Carpetweed (<i>Mollugo verticillata</i>)	A	Pokeweed (<i>Phytolacca Americana</i>)	P
Clover (<i>Trifolium</i> spp.)	A/P	Prickly sida (<i>Sida spinosa</i>)	A
Cocklebur (<i>Xanthium strumarium</i>)	A	Primrose (<i>Oenothera kunthiana</i>)	P
Common chickweed (<i>Stellaria media</i>)	A	Puncturevine (<i>Tribulus terrestris</i>)	A
Common ragweed (<i>Ambrosia artemisiifolia</i>)	A	Purple loosestrife ³ (<i>Lythrum salicaria</i>)	P
Corn spurry (<i>Spergula arvensis</i>)	P	Purslane (<i>Portulaca</i> spp.)	A
Dandelion (<i>Taraxacum officinale</i>)	P	Ragweed (<i>Ambrosia</i> spp.)	A
Dayflower (<i>Commelina</i> spp.)	A/P	Rush skeletonweed ³ (<i>Chondrilla juncea</i>)	B
Desert Camelthorn (<i>Alhagi pseudalhagi</i>)	P	Russian knapweed (<i>Centaurea repens</i>)	P
Diffuse knapweed (<i>Centaurea diffusa</i>)	A	Russian thistle ³ (<i>Salsola kali</i>)	A
Dock (<i>Rumex</i> spp.)	P	Saltbush (<i>Atriplex</i> spp.)	A
Dogfennel (<i>Eupatorium capillifolium</i>)	A	Sesbania (<i>Sesbania</i> spp.)	A
Filaree (<i>Erodium</i> spp.)	A	Sicklepod (<i>Cassia obtusifolia</i>)	A
Fleabane (<i>Erigeron</i> spp.)	A	Silverleaf nightshade (<i>Solanum elaeagnifolium</i>)	P
Giant ragweed ¹ (<i>Ambrosia trifida</i>)	A	Shepherd's-purse (<i>Capsella bursa-pastoris</i>)	A
Goldenrod (<i>Solidago</i> spp.)	P	Smartweed (<i>Polygonum</i> spp.)	A/P
Grey rabbitbrush (<i>Chrysothamnus nauseosus</i>)	P	Sorrell (<i>Rumex</i> spp.)	P
Gromwell (<i>Lithospermum</i> spp.)	A	Sowthistle (<i>Sonchus</i> spp.)	A
Groundcherry (<i>Physalis</i> spp.)	A/P	Speedwell (<i>Veronica</i> spp.)	A
Hawksbeard (<i>Crepis</i> spp.)	A	Stinging nettle ³ (<i>Urtica dioica</i>)	P
Hoary vervain (<i>Verbena stricta</i>)	P	Sunflower (<i>Helianthus</i> spp.)	A
Horsenettle (<i>Solanum Canadensis</i>)	P	Sweet clover (<i>Melilotus</i> spp.)	A/B
Horseweed (<i>Conyza Canadensis</i>)	A	Tansymustard (<i>Descurainia pinnata</i>)	A
Indian mustard (<i>Brassica juncea</i>)	A	Texas thistle (<i>Cirsium texanum</i>)	P
Japanese bamboo (<i>Polygonum cuspidatum</i>)	P	Velvetleaf (<i>Abutilon theophrasti</i>)	A
Knawel (<i>Scleranthus annuus</i>)	A	Western ragweed (<i>Ambrosia psilostachya</i>)	P
Kochia ³ (<i>Kochia scoparia</i>)	A	Wild buckwheat (<i>Polygonum convolvulus</i>)	A
Lambsquarters (<i>Chenopodium album</i>)	A	Wild carrot (<i>Daucus carota</i>)	B
Lespedeza (<i>Lespedeza</i> spp.)	P	Wild lettuce (<i>Lactuca</i> spp.)	A/B
Little mallow (<i>Malva parviflora</i>)	B	Wild parsnip (<i>Pastinaca sativa</i>)	B
Marigold (<i>Tagetes</i> spp.)	P	Wild radish (<i>Raphanus raphanistrum</i>)	B
Milkweed (<i>Asclepias</i> spp.)	P	Wild turnip (<i>Brassica campestris</i>)	B
Miners lettuce (<i>Montia perfoliata</i>)	A	Woolly leaf bursage (<i>Franseria tomentosa</i>)	P
Morningglory (<i>Ipomoea</i> spp.)	A/P	Yellow starthistle (<i>Centaurea solstitialis</i>)	A
Mullein (<i>Verbascum</i> spp.)	B	Yellow woodsorrel (<i>Oxalis stricta</i>)	P

VINES AND BRAMBLES			
Name (Species)	Growth Habit ²	Name (Species)	Growth Habit ²
Blackberry ⁴ (<i>Rubus</i> spp.)	P	Morningglory (<i>Ipomoea</i> spp.)	A/P
Dewberry ⁴ (<i>Rubus</i> spp.)	P	Poison Ivy (<i>Rhus radicans</i>)	P
Field bindweed (<i>Convolvulus arvensis</i>)	P	Redvine (<i>Brunnichia cirrhosa</i>)	P
Greenbriar (<i>Smilax</i> spp.)	P	Trumpetcreeper ⁷ (<i>Campsis radicans</i>)	P
Hedge bindweed (<i>Calystegia sepium</i>)	A	Virginia creeper ⁷ (<i>Parthenocissus quinquefolia</i>)	P
Honeysuckle (<i>Lonicera</i> spp.)	P	Wild buckwheat (<i>Polygonum convolvulus</i>)	P
Kudzu ⁵ (<i>Pueraria lobata</i>)	P	Wild grape (<i>Vitis</i> spp.)	P
		Wild rose (<i>Rosa</i> spp.)	P

BRUSH
Imazuron E-Pro Herbicide controls over 30 species of brush.

¹ Where heavy or well-established infestations occur, use the higher recommended rates.

- 2 Growth Habit: A= Annual, B= Biennial, P= Perennial
- 3 Early postemergence applications are required for best results.
- 4 The degree of control is species dependent; some *Rubus* species may not be completely controlled.
- 5 Use a minimum of 75 GPA; repeat applications may be required to control established stands.
- 6 Control is species dependent; for preemergence control a tank-mix with Pendulum herbicide and/or a postemergence application of a labeled herbicide may be required.
- 7 A minimum of 13 pounds of Imazuron E-Pro Herbicide per acre is required.
- 8 Tank-mix with OUST for best results.
- 9 Repeat applications may be required to control established stands.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in an approved sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use. This warranty does not extend to the use of this product contrary to label instructions, or, to the extent allowable by applicable law, under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. **ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Etigra nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT ALLOWABLE BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Etigra.

Imazuron E-Pro Herbicide is not manufactured or distributed by BASF, seller of Sahara®.

12/12

Arsenal[®], Banvel[®], Pendulum[®], Plateau[®] and Sahara[®] are trademarks of BASF Corporation.
Finale[®] is a trademark of Bayer.
Garlon[®] is a trademark of Dow AgroSciences Company.
Karmex[®] and Oust[®] are trademarks of E.I Du Pont de Nemours and Company.
Etiga[™] is a trademark of Etiga.
Roundup[®] is a trademark of Monsanto Company.
Vanquish[®] is a trademark of a Syngenta Group Company.

EPA 20071016