

80286-3

3-11-2008

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

820925	322
rw	1300

MAR 11 2008

ISCA Technologies, Inc.
c/o Robert Rose, Ph.D.
2060 Chicago ave., Suite C2
Riverside, CA 92507

Subject: Product Name: Solat
EPA Reg. No: 80286-3
Application for label Notification dated November November 26, 2007 to
change the "Warranty Statement".

Dear Dr. Rose

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PRN 98-10 and it has been determined that the action(s) requested falls within the scope of PRN 98-10. Our records have been duly noted, and the label submitted with this application has been stamped "Notification, received and reviewed" and will be placed accordingly in our records.

Questions concerning this action should be directed to Mr. Raderrio Wilkins at (703) 308-1259 or email at wilkins.raderrio@epa.gov.

Sincerely,

Linda Hollis

Linda Hollis, Chief
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention Division

2/15



United States
Environmental Protection Agency
Washington, DC 20460

Registration
Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number Company No. 80286-CA-003 / Product No. 80286-3		2. EPA Product Manager Andrew Bryceland	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restrictive
4. Company/Product (Name) ISCA Technologies, Inc. / SPLAT Cydia		PM# BPPD, 7504P	
5. Name and Address of Applicant (Include ZIP Code) ISCA Technologies, Inc., 2060 Chicago Ave., Suite C2 Riverside, CA 92507, USA <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. 80286-1, 80286-2, 80286-4 Product Name SPLAT OFM 30M-1, PBW 30M-1 & GM	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of change in warranty statement only per PR Notice 98-10, II. Labeling Notification, J. Change in Warranty Statement. See attachment.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 0.2, 1, 2, 2.5, 3.5, 5, 5.5, 6.5, 11, 55 gallons		5. Location of Label Directions <input checked="" type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Robert I. Rose	Title Authorized agent for ISCA Tech. Inc.	Telephone No. (Include Area Code) 301-693-0659
------------------------	---	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Regulatory Agent for ISCA Technologies	3. Date Application Received (Stamped)
4. Typed Name Robert I. Rose	5. Date November 27, 2007	

2

11

3/15

ROBERT I. ROSE, PH.D.
Biotechnology Regulatory Consultant

8322 Sharon Drive
Frederick, Maryland 21704

Phone & Fax 301-874-5593
Mobile 301-693-0659
rirose1@juno.com

November 26, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs, (7504P)
U.S. Environmental Protection Agency
Room S- 4910, 4th floor, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Attn: Mr. Andrew Bryceland
Regulatory Action Leader, Biochemical Pesticide Branch
Biopesticides and Pollution Prevention Division (7504P)

NOTIFICATION
Date Reviewed By: [Signature]
Reviewed By: [Signature]

**Subject: Fast-Track Label Notification according to PR Notice 98-10
Change of warranty statement only
EPA Reg. No. 80286-3**

Dear Mr. Bryceland:

Enclosed are the following:

1. EPA Form 8570-1, Application for pesticide label Notification according to PR Notice 98-10, II, Labeling Notifications, J. Changes in Warranty Statement. Only the warranty statement has been changed. This new warranty statement has already been accepted by the Agency on the SPLAT OFM 30M-1, EPA Reg. No. 80286-1, label stamped accepted by the Agency on August 14, 2007.
2. Copy of SPLAT Cydia EPA stamped accepted on May 24, 2007.
3. Copy of the SPLAT OFM 30M-1 label stamped accepted by the Agency on August 14, 2007.
4. Copy of SPLAT Cydia draft label with change of warranty statement, bold font, as the only change. I understand this change of warranty statement was requested by the California Department of Pesticide Registration for registration in the State of California.

This application for label change by notification is submitted in triplicate. Please contact me by e-mail or phone if there are any questions or problems concerning this notification for pesticide amendment.

Sincerely yours,



Robert I. Rose
Regulatory Consultant/Agent
cc: ISCA Technologies, Inc.

4/15

Attachment to EPA Form 8570-1.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations as 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

RER
11/27/07

5/15

SPLAT **Cydia**

NOTIFICATION
Date Reviewed: 3/11/08
Reviewed By: [Signature]

For Mating Disruption of the Codling Moth, *Cydia pomonella* and Hickory Shuckworm, *Cydia caryana*

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained release of insect pheromones. SPLAT Cydia provides control by disrupting mating behavior.

ACTIVE INGREDIENT:

Codling Moth Technical Pheromone	
(E,E)-8,10-Dodecadien-1-ol:.....	10.0%
OTHER INGREDIENTS:.....	90.0%
Total:.....	100.0%

KEEP OUT OF REACH OF CHILDREN **CAUTION**

See side/back panel for Additional Precautionary Statements

EPA Reg. No. 80286-3
EPA Est. No. 80286-CA-003

Net Contents: ___ lbs ___ kg

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

If on skin or clothing:

- Take off contaminated clothing.
- Wash skin immediately with soap and water and rinse with plenty of water for 5-10 minutes.
- Call poison control center or doctor for treatment advice.

If swallowed:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless advised to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Hotline Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR MEDICAL EMERGENCY INFORMATION CALL 1-800-222-1222.**

6/15

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals:

CAUTION. Causes moderate eye irritation. Harmful if swallowed, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before use.

Personal Protective Equipment (PPE): Applicators and handlers must wear long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves, and protective eyewear. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands thoroughly with soap and water after handling, before eating, drinking, chewing gum, using tobacco, or using the toilet.
Users should remove clothing/PPE immediately if pesticide gets inside. Then shower cleaning skin thoroughly with soap and water and put on clean clothing.
Users should remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. Remove and wash contaminated clothing before reuse.
As soon as possible, wash body and scalp thoroughly with soap and water and change into clean clothing.

ENVIRONMENTAL HAZARDS:

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE:

It is a violation Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State and Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI), which for this product is 4 hours. For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemically resistant gloves, shoes plus socks, and protective eyewear.

SPLAT Cydia is an insect pheromone mating disruptant for control of many moth species of the genus *Cydia*, principally the codling moth, *Cydia pomonella*, on pome fruit, walnut, quince, apple, pear, crabapple and cherry and the hickory shuckworm, *Cydia caryana*, on pecan. The disruption of adult moth mating will reduce larval damage to fruit and nuts through reduction of larval pest densities. The application rate of SPLAT Cydia is 400 grams up to 1,000 grams (1 kg) of undiluted product per acre. This is equivalent to 40 to 100 grams of the pheromone active ingredient (AI) per acre. Never exceed

7/15

2 applications of 100 grams first and then 40 grams second of the pheromone AI per acre per year. Do not apply more than 140 grams of the active ingredient per acre per year.

Size and Spacing of SPLAT Cydia dollops: SPLAT is unlike other pheromone dispensing products because the drop or dollop size can be made small or large depending on pest population density, the amount of pheromone applied, and the duration of control desired. A high pest infestation is more quickly and effectively controlled by many small dollops that emit more pheromone in a shorter time than fewer large dollops. The larger dollops emit the same amount of pheromone as small dollops, but are longer lasting making them more useful for long-term suppression of moderate to low density *Cydia* populations. High pest populations require two applications; the first with a high number of smaller point sources or dollops and a second application a month later with fewer larger dollops for more persistent control. For a maximum of two applications per year, apply 100 grams AI per acre first followed by 40 grams AI per acre in the second application.

Method of Application: Do not open containers until ready to use. Do not use SPLAT from damaged or punctured containers. Apply SPLAT Cydia early in the season or immediately after the first moth is detected in pheromone-baited insect traps, because the pheromone has to be applied before most mating is expected. The early or first application is effective for about 100 days. Reapply SPLAT Cydia, based on monitoring result and field scouting.

Apply SPLAT Cydia evenly over the acreage to be treated. If insect scouts are employed to monitor fruit pests, they may advise on the size, timing, and number of deposits or dollops to use per acre after they have read this label. Apply dollops of the product on branches, tree trunks, bark, or foliage as high as can be reached by an adult and out of reach of children. The dollop size can vary depending on the number of pheromone sources per acre. To place 100 sources on one acre, evenly deposit dollops of 10 ± 2 grams. To place 1,000 sources on one acre, deposit dollops of 1 ± 0.2 grams.

Use a small spatula, small putty knife, or household-measuring spoon to apply SPLAT Cydia directly out of its container without diluting it. The same size dollop or same amount of product should be applied on each tree. If measuring spoons are used for application, one tablespoon contains 14 grams of product; one teaspoon 4.6 grams; $\frac{1}{2}$ teaspoon 2.3 grams; and $\frac{1}{4}$ teaspoon 1.15 grams. If 1000 grams are to be used for the application, divide 1,000 grams by the approximate number of trees per acre to find out how many grams of product to use per tree. Next, decide if you want to treat every tree, every second tree, or one tree out of three or four. If you treat every tree, use that amount per tree; if you treat every second tree, use twice that amount per tree; if you treat one tree out of three or four, use three or four times that amount per tree. Then select the measuring spoon size that holds the amount closest to the amount you need to put on a tree. Empty the spoonful on the tree while wearing chemically resistant gloves. Do not use measuring spoons or spatulas afterward for any food or feed.

Another method of application is the use of a pressurized backpack sprayer equipped with an extension tube attached to a metering device. The metering device can consist either of a timer/solenoid valve or a positive displacement system. Using the timer/solenoid valve, the trigger activates a timer to open a solenoid valve for a fixed interval of time. A positive displacement system consists of a piston in a cylinder delivering a fixed volume of material whenever the trigger is pulled. The meterjet gun (a TeeJet® product manufactured by Spraying Systems Co.) works for this application. In either delivery system, calibrate the applicator by adjusting either the internal application pressure, or the interval of time the solenoid valve remains open, or the volume of material that is displaced. Never exceed two applications of 1,000 grams first and then 400 grams second of SPLAT Cydia per acre per year.

Do not handle other types of insect pheromones or lures before or after SPLAT Cydia use without thoroughly washing gloves and hands. Monitor treatments carefully, especially for other primary and secondary pests, which may need other control measures.

Cleaning equipment: Clean equipment with water and household-type detergent.

8/15

Do not handle other types of insect pheromones or lures before or after SPLAT use without thoroughly washing gloves and hands.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store sealed container in a cool well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

Container Disposal: Do not reuse empty container. Completely empty the container. Then, dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



Manufactured by:
ISCA Technologies, Inc.
2060 Chicago Ave, Suite C2
Riverside, CA 92507
(951) 686-5008
info@iscatech.com

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. ISCA Technologies warrants that this material conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label. To the extent allowable by state law, ISCA Technologies makes no other express or implied warranty. Read all directions carefully.

9/15

SPLAT Cydia

For Mating Disruption of the Codling Moth, *Cydia pomonella* and Hickory Shuckworm, *Cydia caryana*

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained release of insect pheromones. SPLAT Cydia provides control by disrupting mating behavior.

ACTIVE INGREDIENT:

Codling Moth Technical Pheromone	
(E,E)-8,10-Dodecadien-1-ol:.....	10.0%
OTHER INGREDIENTS:	90.0%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side/back panel for Additional Precautionary Statements

**EPA Reg. No. 80286-
EPA Est. No. 80286-CA-003**

Net Contents: ___ lbs ___ kg

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

If on skin or clothing:


- Take off contaminated clothing.
- Wash skin immediately with soap and water and rinse with plenty of water for 5-10 minutes.
- Call poison control center or doctor for treatment advice.

If swallowed:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless advised to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Hotline Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR MEDICAL EMERGENCY INFORMATION CALL 1-800-222-1222.**



10/15

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals:

CAUTION. Causes moderate eye irritation. Harmful if swallowed, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before use.

Personal Protective Equipment (PPE): Applicators and handlers must wear long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves, and protective eyewear. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should wash hands thoroughly with soap and water after handling, before eating, drinking, chewing gum, using tobacco, or using the toilet.
Users should remove clothing/PPE immediately if pesticide gets inside. Then shower cleaning skin thoroughly with soap and water and put on clean clothing.
Users should remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. Remove and wash contaminated clothing before reuse.
As soon as possible, wash body and scalp thoroughly with soap and water and change into clean clothing.

ENVIRONMENTAL HAZARDS:

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE:

It is a violation Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State and Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI), which for this product is 4 hours. For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemically resistant gloves, shoes plus socks, and protective eyewear.

SPLAT Cydia is an insect pheromone mating disruptant for control of many moth species of the genus *Cydia*, principally the codling moth, *Cydia pomonella*, on pome fruit, walnut, quince, apple, pear, crabapple and cherry and the hickory shuckworm, *Cydia caryana*, on pecan. The disruption of adult moth mating will reduce larval damage to fruit and nuts through reduction of larval pest densities. The application rate of SPLAT Cydia is 400 grams up to 1,000 grams (1 kg) of undiluted product per acre. This is equivalent to 40 to 100 grams of the pheromone active ingredient (AI) per acre. Never exceed

11/15

2 applications of 100 grams first and then 40 grams second of the pheromone AI per acre per year. Do not apply more than 140 grams of the active ingredient per acre per year.

Size and Spacing of SPLAT Cydia dollops: SPLAT is unlike other pheromone dispensing products because the drop or dollop size can be made small or large depending on pest population density, the amount of pheromone applied, and the duration of control desired. A high pest infestation is more quickly and effectively controlled by many small dollops that emit more pheromone in a shorter time than fewer large dollops. The larger dollops emit the same amount of pheromone as small dollops, but are longer lasting making them more useful for long-term suppression of moderate to low density *Cydia* populations. High pest populations require two applications; the first with a high number of smaller point sources or dollops and a second application a month later with fewer larger dollops for more persistent control. For a maximum of two applications per year, apply 100 grams AI per acre first followed by 40 grams AI per acre in the second application.

Method of Application: Do not open containers until ready to use. Do not use SPLAT from damaged or punctured containers. Apply SPLAT Cydia early in the season or immediately after the first moth is detected in pheromone-baited insect traps, because the pheromone has to be applied before most mating is expected. The early or first application is effective for about 100 days. Reapply SPLAT Cydia, based on monitoring result and field scouting.

Apply SPLAT Cydia evenly over the acreage to be treated. If insect scouts are employed to monitor fruit pests, they may advise on the size, timing, and number of deposits or dollops to use per acre after they have read this label. Apply dollops of the product on branches, tree trunks, bark, or foliage as high as can be reached by an adult and out of reach of children. The dollop size can vary depending on the number of pheromone sources per acre. To place 100 sources on one acre, evenly deposit dollops of 10 ± 2 grams. To place 1,000 sources on one acre, deposit dollops of 1 ± 0.2 grams.

Use a small spatula, small putty knife, or household-measuring spoon to apply SPLAT Cydia directly out of its container without diluting it. The same size dollop or same amount of product should be applied on each tree. If measuring spoons are used for application, one tablespoon contains 14 grams of product; one teaspoon 4.6 grams; $\frac{1}{2}$ teaspoon 2.3 grams; and $\frac{1}{4}$ teaspoon 1.15 grams. If 1000 grams are to be used for the application, divide 1,000 grams by the approximate number of trees per acre to find out how many grams of product to use per tree. Next, decide if you want to treat every tree, every second tree, or one tree out of three or four. If you treat every tree, use that amount per tree; if you treat every second tree, use twice that amount per tree; if you treat one tree out of three or four, use three or four times that amount per tree. Then select the measuring spoon size that holds the amount closest to the amount you need to put on a tree. Empty the spoonful on the tree while wearing chemically resistant gloves. Do not use measuring spoons or spatulas afterward for any food or feed.

Another method of application is the use of a pressurized backpack sprayer equipped with an extension tube attached to a metering device. The metering device can consist either of a timer/solenoid valve or a positive displacement system. Using the timer/solenoid valve, the trigger activates a timer to open a solenoid valve for a fixed interval of time. A positive displacement system consists of a piston in a cylinder delivering a fixed volume of material whenever the trigger is pulled. The meterjet gun (a TeeJet® product manufactured by Spraying Systems Co.) works for this application. In either delivery system, calibrate the applicator by adjusting either the internal application pressure, or the interval of time the solenoid valve remains open, or the volume of material that is displaced. Never exceed two applications of 1,000 grams first and then 400 grams second of SPLAT Cydia per acre per year.

Do not handle other types of insect pheromones or lures before or after SPLAT Cydia use without thoroughly washing gloves and hands. Monitor treatments carefully, especially for other primary and secondary pests, which may need other control measures.

Cleaning equipment: Clean equipment with water and household-type detergent.

12/15

Do not handle other types of insect pheromones or lures before or after SPLAT use without thoroughly washing gloves and hands.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store sealed container in a cool well-ventilated area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

Container Disposal: Do not reuse empty container. Completely empty the container. Then, dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



TECHNOLOGIES
Pest Management Tools & Solutions

Manufactured by:
ISCA Technologies, Inc.
2060 Chicago Ave, Suite C2
Riverside, CA 92507
(951) 686-5008
info@iscatech.com

WARRANTY AND LIMITATION OF DAMAGES

ISCA Technologies, Inc. (ISCA) warrants that this material conforms to the chemical description on the label. ISCA neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material. To the extent allowable by state law, ISCA's maximum liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this material, whether in accordance with directions or not.

13/15

ACCEPTED
AUG 14 2007
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 80286-1

SPLAT

OFM 30M-1

For Mating Disruption of the Oriental Fruit Moth, *Grapholita molesta*

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained release of semiochemicals. SPLAT OFM 30M-1 provides control of the oriental fruit moth via mating disruption.

ACTIVE INGREDIENT:

Oriental Fruit Moth Technical Pheromone

(Z)-8-Dodecenyl acetate	2.790%	}3.0%
(E)-8-Docecenyl acetate	0.179%	
(Z)-8-Dodecenol	0.030%	

OTHER INGREDIENTS:	97.0%
Total	100.0%

Net Contents: _____ lbs _____ kg

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back panel for Additional Precautionary Statements

EPA Reg. No. 80286-1
EPA Est. No. 80286-CA-003

Net Contents: _____ lbs _____ kg

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call poison control center or doctor for treatment advice.

If swallowed:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Hotline Number

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **FOR MEDICAL EMERGENCY INFORMATION CALL 1-800-222-1222.**

14/15

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Do not apply this product in such a manner as to directly expose workers or other persons. Unprotected persons must vacate the area being treated.

Personal Protective Equipment (PPE): Applicators and handlers must wear long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves, and protective eyewear. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State and Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment, restricted entry interval, and notification to workers (as applicable). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI of 4 hours). For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemically resistant gloves, shoes plus socks, and protective eyewear.

SPLAT OFM 30M-1 is an insect pheromone used to control the oriental fruit moth (*Grapholita molesta*) by disrupting adult moth mating and thus reducing larval damage to crops. It is for use on peaches, nectarines, apricots, plums, apples, quince, almond, macadamias, and other tree crops where this pest occurs. The application rate of **SPLAT OFM 30M-1** is 1,000 grams of undiluted product per acre. This is equivalent to 30.0 grams of oriental fruit moth pheromone active ingredient (AI) per acre. Do not exceed four applications per year to insure a total application rate below 150 grams of the AI per acre. Two applications per year are usually enough to provide effective control.

15/15

Method of Application: Do not open containers until ready to use. Do not use **SPLAT OFM 30M-1** from damaged or punctured containers. Apply **SPLAT OFM 30M-1** early in the season or immediately after the first moth is detected in pheromone baited insect traps. The early application is effective for about 100 days. Reapply **SPLAT OFM 30M-1** as needed, based on local conditions, monitoring result, and field scouting.

Apply **SPLAT OFM 30M-1** evenly over the acreage to be treated. If insect scouts are employed, they may advise on the size and number of deposits or dollops to use per acre after they have read this label. Apply dollops of the product on branches, tree trunks, bark, or foliage as high as can be reached by an adult and out of reach of children. The dollop size can vary depending on the number of pheromone sources per acre. To place 100 sources on one acre, evenly deposit dollops of 10 ± 2 grams. To place 1,000 sources on one acre, deposit dollops of 1 ± 0.2 grams.

Use a small spatula, small putty knife, or household-type measuring spoon to apply the **SPLAT OFM 30M-1** directly out of its container without diluting it. The same size dollop or same amount of product should be applied on each tree. If measuring spoons are emptied, one tablespoon contains 14 grams of product; one teaspoon 4.6 grams; ½ teaspoon 2.3 grams; ¼ teaspoon 1.15 grams. Divide 1,000 grams by the approximate number of trees per acre to find out how many grams of product to use per tree. Next, decide if you want to treat every tree, every second tree, or one tree out of three or four. If you treat every tree simply use that amount per tree; if you treat every second tree, use twice that amount per tree; if you treat one tree out of three or four, use three or four times that amount per tree. Then select the measuring spoon size that holds the amount closest to the amount you need to put on a tree. Empty the spoonful on the tree while wearing chemically resistant gloves. Do not use measuring spoons or spatulas afterward for any food or feed.

Do not handle other types of insect pheromones or lures before or after **SPLAT OFM 30M-1** use without thoroughly washing gloves and hands. Monitor treatments carefully, especially for other primary and secondary pests such as leafrollers.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool well-ventilated area. Store sealed and refrigerated to prevent loss of product.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

Container Disposal: Do not reuse empty container. Completely empty the container. Then, dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



Manufactured by:
ISCA Technologies, Inc.
2060 Chicago Ave, Suite C2
Riverside, CA 92507
(951) 686-5008
info@iscatech.com

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. ISCA Technologies warrants that this material conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label. To the extent allowable by state law, ISCA Technologies makes no other express or implied warranty. Read all directions carefully.