UNITED STATES - JON BOR	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs	EPA Reg. Number: 80286-6	Date of Issuance:
ROMANNAL PROTECTION	Biopesticides and Pollution Prevention Division (7511C) 1200 Pennsylvania Avenue NW Washington, DC 20460	Term of Issuance:	Conditional
		Name of Pesticide Prod	
	NOTICE OF PESTICIDE: <u>X</u> Registration <u>Reregistration</u> (under FIFRA, as amended)	Splat L	BAM HD
lame and Address of Regis		· · · · · · · · · · · · · · · · · · ·	
SCA Technolog	jies, Inc.		
•	venue, Suite C2		
Riverside, CA 9	2507		
Note: Changes in labeli Biopesticides and Pollut bove EPA registration	ng differing in substance from that accepted in connection with this regist ion Prevention Division prior to use of the label in commerce. In any corr number	tration must be submitted to respondence on this produ	o and accepted by the ct always refer to the
nvironment, the Admini- acceptance of any name exclusive use of the name This product is unc 1. Submit and/o	to be construed as an endorsement or recommendation of this product I istrator, on his motion, may at any time suspend or cancel the registration in connection with the registration of a product under this Act is not to be ne or to its use if it has been covered by others. onditionally registered in accordance with FIFRA Sec. 3(c) r cite all data required for registration/ reregistration of you b) and section 4 when the Agency requires all registrants of lata.	n of a pesticide in accordan e construed as giving the re)(7) provided you: ur product under FIFR	ce with the Act. The egistrant a right to
submit such c			
submit such c	PA Registration Number to read, "EPA Reg. No. 80286-6."))	
submit such of 2. Revise the EF			shipment.
 submit such of Revise the EF Submit three Submit storage interim reportione year of the 	A Registration Number to read, "EPA Reg. No. 80286-6."	elease the product for n one year of the Date ance. Failure to submi	of Issuance. An the study within
 submit such of 2. Revise the EF 3. Submit three 4. Submit storage interim reportione year of the Date of Is 	PA Registration Number to read, "EPA Reg. No. 80286-6." (3) copies of the revised final printed labeling before you r ge stability study (OPPTS 830.6317) for this product within t for these data is required 6 months from the Date of Issua he registration notice date will cause this product registration scuance of this registration.	elease the product for n one year of the Date ance. Failure to submi	of Issuance. An the study within
 submit such of Revise the EF Submit three Submit storage interim reportione year of the Date of Is A stamped copy of Signature of Approvi 	PA Registration Number to read, "EPA Reg. No. 80286-6." (3) copies of the revised final printed labeling before you r re stability study (OPPTS 830.6317) for this product within t for these data is required 6 months from the Date of Issua re registration notice date will cause this product registration suance of this registration.	elease the product for n one year of the Date ance. Failure to submi on to expire on or abo	of Issuance. An the study within
 submit such of Revise the EF Submit three Submit storag interim reportione year of the the Date of Is Stamped copy of 	PA Registration Number to read, "EPA Reg. No. 80286-6." (3) copies of the revised final printed labeling before you r re stability study (OPPTS 830.6317) for this product within t for these data is required 6 months from the Date of Issua re registration notice date will cause this product registration suance of this registration.	elease the product for n one year of the Date ance. Failure to submi on to expire on or abo	of Issuance. An the study within



'JUN - 8 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 30236-6

ACTIVE INGREDIENT:

(E)-11-Tetradecen-1-yl acetate	
(E,E)-9,11-Tetradecadien-1-yl acetate	
OTHER INGREDIENTS	
TOTAL	

Net Contents: _____ Lbs _____ Batch Number: _____

EPA Reg. No. 80286-XXX EPA Est. No. 80286-CA-003

LBAM HD

For Mating Disruption of the Light Brown Apple Moth, *Epiphyas postvittana*

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained controlled release of insect pheromones. SPLAT LBAM provides control of the light brown apple moth by disrupting mating behavior.



KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID STATEMENT

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Call poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing.

• Wash skin immediately with soap and water, then rinse with plenty of water for 15-20 minutes.

IF SWALLOWED:

- · Call poison control center or doctor immediately for treatment advice.
- · Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by poison control center or doctor.
- · Do not give anything by mouth to an unconscious person.

Hotline Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR MEDICAL EMERGENCY INFORMATION CALL 1-800-222-1222.

See side/back panel for additional precautionary statements

PAROMINICALINASI PAREMEN

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION. Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, and clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves and protective eyewear.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and handlers must wear long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves, and protective eyewear. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: For terrestrial uses: Except under the forest canopy, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Spray valves must be shut off when passing over surface water not under the canopy. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Users should remove clothing/PPE immediately if product gets underneath clothing. Then, shower cleaning skin thoroughly with soap and water and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. Remove and wash contaminated clothing before reuse. As soon as possible, wash thoroughly and change into clean clothing.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is for both agricultural crop use and for wide-area public pest control programs sponsored by governmental entities. For use in light brown apple moth wide-area control programs on all parks and forests, residential and municipal areas, recreational, and non-crop areas. When used just for agricultural crops, follow the precautions in the Agricultural Use Requirements box directly below. For any requirement specific to your State and Tribe, consult the State/Tribe agency responsible for pesticide regulation.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted entry interval and notification to workers (as applicable). Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers handlers may be in the area during application. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, wear: coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

	NGNEAGRICEDU DI PARISERE QUI REMENTS SA
	e box apply to uses of this product that are NOT within the scope of the Worker Protection
Standard for agricultur	al pesticides (40 CFR Part 170). The WPS applies when this product is used to produce
agricultural plants on fa	arms, forests, nurseries, or greenhouses.
GENERAL	SPLAT LBAM HD uses the pheromone of the light brown apple moth (<i>Epiphyas postvittana</i>) to disrupt adult moth mating and thus reduce larval damage to crops.
CROPS	SPLAT LBAM HD is for use on all crops and non-crop areas where the light brown apple moth is detected.
TIMING	Apply early in the season, prior to and at the first sign of male flight. Reapply as needed based on trap capture and scouting programs. Consult your own pest control advisor, distributor or ISCA Technologies to obtain correct timing for application.
PLACEMENT	Apply SPLAT LBAM HD dollops at upper level of plant foliage, tree canopy, or as high in trees as possible.
RATE	The application rate of SPLAT LBAM HD is 150 - 1500 grams of undiluted product per acre. This is equivalent to 15 - 150 grams of pheromone active ingredient (AI) per acre. Never exceed more than 150 grams of active ingredient (1500 grams of SPLAT LBAM HD) per acre per year.
POINT SOURCE SIZE, NO. PER ACRE, AND SPACING	Droplet or dollop size can be varied (large or small) depending on 1) amount of active ingredient pheromone applied per acre, and 2) duration of control required. A higher number of small dollops will emit more pheromone in a shorter time interval than fewer large dollops. While the pheromone emission rate per unit area remains constant for large and small dollops, the larger dollops have a greater volume/reservoir, which increases its field life making it useful where long-term suppression of LBAM populations is required. Dollop size can vary depending on number of pheromone point sources required. To apply 300 g of SPLAT as 3,000 point sources per acre, distribute 0.1 gram (100 mg) dollops uniformly throughout the area. To apply 300 g as 10,000 point sources, distribute 0.03 gram dollops (30 mg) uniformly throughout the area.
	When applied by aircraft, SPLAT will be dispersed in droplets of various sizes. While most of the

3/ч

	volume of SPLAT LBAM HD applied aerially is represented by droplets 1/8" (3 mm) in diameter,
	equivalent to a 30 mg, some droplets will be larger and some smaller. Actual droplet size can be controlled by the nozzles type and the rate of product applied. Typically, 100 mg droplets will
	have a diameter of 0.2-0.24" (5-6 mm); 30 mg droplets will have a diameter of 0.1-0.125" (2.5-3.0
	mm) and a 5 mg droplet will have a diameter of 0.02-0.03" (0.5-0.75 mm).
EXTERNAL	Use the mating disruption strategy on areas greater than 10 acres for best results. Monitor for
	pests on a regular basis, so that timely interventions with pesticides or other control measures are
FACTORS	possible.
PREPARATION	Always check SPLAT LBAM HD 24 hours before using. If SPLAT LBAM HD has been stored in
FOR USE	a refrigerated area and has hardened, allow at least 48 hours at room temperature to achieve a
	workable consistency. Always mix SPLAT LBAM HD thoroughly before use. A drill with paint
	mixing paddle or similar device works best. Avoid mechanical mixing with sharp blades, which come into contact with the plastic container as this can introduce shards of plastic into the SPLAT
	LBAM HD mixture, which might interfere with product flowability and application.
APPLICATION	SPLAT LBAM HD can be applied manually with items such as a thin spatula, small caulking gun-
1	style dosing gun, pressurized backpack sprayer, or mechanically with a tractor, all terrain vehicle
METHOD	(ATV) truck, aircraft or appropriate mechanical device. Do not apply this product through any
	type of irrigation system.
MECHANICAL	All these application methods typically involve the use of a hopper, pump (positive displacement
APPLICATION	or diaphragm type) boom and output nozzles. For mechanical ground applications, the flow from
	each nozzle is controlled by a standard on-off solenoid valve positioned at each nozzle. These
	solenoids are regulated both by a timer and a simple on-off override switch. The timer is adjusted to open and close the solenoid valve at a rate that is related to the speed of the application
	equipment; the faster the applicator, the shorter the valve closed time. When using mechanical
	application equipment, calibrate the system to deliver the intended amount of material per acre
	(e.g. 150 grams of SPLAT LBAM HD). For unusual situations, always consult your dealer,
	distributor or ISCA Technologies to ensure correct and optimum application of product is
	achieved.
MONITORING	Monitor the target pest(s) with pheromone-baited traps, such as the ISCAlure® LBAM pheromone
	lure and ISCA delta trap monitoring system. Place monitoring traps in the treated area before applying SPLAT LBAM HD to determine the infestation level. To prevent contaminating the
	monitoring traps with SPLAT LBAM HD, either cover the traps or remove the traps before
	treating the area, then following application, uncover or replace the traps. Continue monitoring
	throughout the season to evaluate efficacy and longevity of the application. Reapply as needed
	based on trap capture and field scouting programs.
WEATHER	Apply SPLAT LBAM HD when the ambient temperature is above 55°F and below 95°F. SPLAT
FACTORS	dollops typically cure within 2-3 hours following application, after which they will become rainfast
	and UV resistant. Do not apply if rain is expected within 2-3 hours of application or the
	temperature is outside these ranges. Clean equipment with water and citrus or limonene detergents.
CLEANING	Clean equipment with water and claus of innonene detergents.
EQUIPMENT	
	SIGRAGE/AND DISPOSAL
	Do not contaminate water, food or feed by storage and disposal.
PESTICIDE	Store product sealed and refrigerated, if possible. If refrigerated, do not store with food. If refrigeration is not available, store product in a cool dry place, out of direct sunlight. Do not
STORAGE	exceed 75 F for long-term storage. Avoid freezing. In case of leak or spill, wipe with paper towel
· ·	and dispose of waste when product hardens.
PESTICIDE	Wastes resulting from the use of this product may be disposed of on site or at an approved waste
DISPOSAL	facility.
CONTAINER	Nonrefillable container. Do not refill or reuse this container. Triple rinse (or equivalent) promptly
1	after emptying. Then offer for recycling if available, or puncture and dispose of in a sanitary
DISPOSAL	landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay
	out of smoke.
	WAREAND AND HARDER HAD AND AND AND AND AND AND AND AND AND A
Read all directions car	efully. All statements concerning the use of this product apply only when used as directed. ISCA
Technologies warrants	that this material conforms to the specifications on the label and is reasonably fit for the intended
purpose referred to on	the label. To the extent consistent with applicable law, ISCA Technologies makes no other express
or implied warranty of	merchantability or fitness for a particular purpose or any other express or implied warranty.

4/4