

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs

Biopesticides and Pollution Prevention Division (7511P)

1200 Pennsylvania Avenue NW Washington, DC 20460

EPA Reg. Number:

80286-8

Date of Issuance:

SEP 1 1 2009

Term of

Issuance:

Unconditional

Name of Pesticide Product:

SPLAT LBAM LD

NOTICE OF PESTICIDE:

X Registration

_ Re-registration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

ISCA Technologies, Inc.

2060 Chicago Ave., Suite C2

Riverside, CA 92507

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This registration does not eliminate the need for continual reassessment of the pesticide. If the EPA determines at any time, that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under section 3(c)(2)(B) of FIFRA.

The product is registered in accordance with FIFRA section 3(c)(5) and is subject to the following terms and conditions:

- 1. Submit and/or cite all data required for registration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
- 2. Submit within 1 year of the date of registration data packages for Guideline Studies: OPPTS 830.6320 (Corrosion Characteristics) and OPPTS 830.6317 (Storage Stability).
- 3. Make the following label change before you release the product for shipment: Revise the EPA Registration Number to read, "EPA Reg. No. 80286-8."
- 4. Submit two (2) copies of the revised final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of the final printed label.

A stamped copy of the label is enclosed for your record.

Signature of Approving Official:

W. Michael McDat

Date:

9-11-09

W. Michael McDavit, Associate Director Biopesticides and Pollution Prevention Division

SPLATTM LBAM LD

For Mating Disruption of the Light Brown Apple Moth, Epiphyas postvittana

SPLAT (Specialized Pheromone and Lure Application Technology) is an amorphous polymer matrix for the sustained controlled release of insect pheromones. SPLAT LBAM LD provides control of the Light Brown Apple Moth by disrupting mating behavior.

ACCEPTED



For Organic Production

SEP 1 1 2009

ACTIVE INGREDIENT:

Light Brown Apple Moth Technical Pheromone

(E)-11-Tetradecen-1-yl acetate:

(E,E)-9,11-Tetradecadien-1-yl acetate:

(E,E)-9,11-Tetradecadien-1-yl

CAUTION

FIRST AID STATEMENT

IF IN EYES:

• Hold eye open and rinse slowly and gently with water for 15-20 minutes.

TOTAL:.....100.00%

- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Wash skin immediately with soap and water, then rinse with plenty of water for 15-20 minutes.

IF SWALLOWED:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu).

See side/back panel for additional precautionary statements.

Net Contents:	Lbs	Kg	EPA Reg. No. 80286-
Batch Number:			EPA Est. No. 80286-CA-003

3066

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION. Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, and clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and handlers must wear long-sleeved shirt and long pants, socks, shoes, chemical resistant gloves, and protective eyewear. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Users should remove clothing/PPE immediately if product gets underneath clothing. Then, shower cleaning skin thoroughly with soap and water and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. Remove and wash contaminated clothing before reuse. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Incidental applications to water under the forest canopy are not considered to be a direct application to water, but should be avoided if practicable. Spray valves must be shut off when passing over surface water not under the canopy. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is for both agricultural crop use and for wide-area public pest control programs sponsored by governmental entities. For use in light brown apple moth wide-area control programs on all parks and forests, residential and municipal areas, recreational, and non-crop areas. When used just for agricultural crops, follow the precautions in the Agricultural Use Requirements box directly below. For any requirement specific to your State and Tribe, consult the State/Tribe agency responsible for pesticide regulation. Always consult your pest control advisor, distributor or ISCA Technologies, Inc. for advice regarding the use of **SPLAT LBAM LD**.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted entry interval and notification to workers (as applicable). Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers handlers may be in the area during application. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow any worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, water, wear: coveralls, chemical resistant gloves, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets out of treated areas until sprays or other wet applications have dried.

PRODUCT	SPLAT LBAM LD uses the pheromone of the light brown apple
INFORMATION	moth (<i>Epiphyas postvittana</i>) to disrupt adult moth mating and thus
	reduce larval damage to crops.
CROPS	SPLAT LBAM LD is for use on all crops or areas where the light
	brown apple moth is detected.
TIMING	Apply early in the season, prior to and at the first sign of male flight.
	Reapply as needed based on trap capture and field scouting
	programs. Consult your own pest control advisor, distributor or ISCA
	Technologies to obtain correct timing for application.
PLACEMENT	Apply SPLAT LBAM LD dollops to the upper portions of the canopy
	or as high in tree as possible.
RATE	The application rate of SPLAT LBAM LD is 5 to 5000 grams of
	undiluted product per acre. This is equivalent to .05 - 50 grams of
	pheromone active ingredient (AI) per acre. Never exceed more than
	150 grams of active ingredient (15 kg of SPLAT LBAM LD) per acre
	per year.
POINT	Droplet or dollop size can be varied (large or small) depending on 1)
SOURCE SIZE,	amount of active ingredient pheromone applied per acre, and 2)
NO. PER	duration of control required. A higher number of small dollops will
ACRE, AND	emit more pheromone in a shorter time interval than fewer large
SPACING	dollops. While the pheromone emission rate per unit area remains
	constant for large and small dollops; the larger dollops have a

<u> </u>	
	greater volume / reservoir which increases its field life making it useful where long-term suppression of LBAM populations is required. Dollop size can vary depending on number of pheromone point sources required. To apply 300 g of SPLAT as 3,000 point sources per acre, distribute 0.1 gram (100 mg) dollops uniformly throughout the area. To apply 300 g as 10,000 point sources, distribute 0.03 gram dollops (30 mg) uniformly throughout the area. When applied by aircraft, SPLAT will be dispersed in droplets of various sizes. While most of the volume of SPLAT LBAM LD applied aerially is represented by droplets 1/8" (3 mm) in diameter, equivalent to a 30 mg; some droplets will be larger and some smaller. Actual droplet size can be controlled by the nozzles type and the rate of product applied. Typically, 100 mg droplets will have a diameter of 0.2-0.24" (5-6 mm); 30 mg droplets will have a diameter of 0.1-0.125" (2.5-3.0 mm) and a 5 mg droplet will have a
E)/TED1:::	diameter of 0.02-0.03" (0.5-0.75 mm).
EXTERNAL	Use the mating disruption strategy on areas greater than 10 acres for
FACTORS	best results. Monitor for pests on a regular basis, so that timely interventions with pesticides or other control measures are possible.
PREPARATION	Always check SPLAT LBAM LD 24 hours before using. If SPLAT
FOR USE	LBAM LD has been stored in a refrigerated area and has hardened,
, or ose	allow at least 48 hours at room temperature to achieve a workable
	consistency. Always mix SPLAT LBAM LD thoroughly before use. A
	drill with paint mixing paddle or similar device works best. Avoid
	mechanical mixing with sharp blades, which come into contact with
	the plastic container as this can introduce shards of plastic into the
	SPLAT LBAM LD mixture, which might interfere with product
	flowability and application. If SPLAT LBAM LD is combined with
	other products, make sure the resultant mix it is uniform before
	application. Observe the mix for 30 minutes to ensure that no
	adverse change in composition (e.g. separation, coagulation) has
	taken place.
APPLICATION	SPLAT LBAM LD can be applied manually with items such as a thin
METHOD	spatula, small caulking gun-style dosing gun, pressurized backpack
	sprayer, or mechanically with a tractor, all terrain vehicle (ATV)
	truck, aircraft, or appropriate mechanical device. Do not apply this
MECHANICAL	product through any type of irrigation system. All these application methods typically involve the use of a hopper,
APPLICATION	pump (positive displacement or diaphragm type) boom and output
	nozzles. For mechanical ground applications, the flow from each
	nozzle is controlled by a standard on-off solenoid valve positioned at
i	each nozzle. These solenoids are regulated both by a timer and a
	simple on-off override switch. The timer is adjusted to open and
	close the solenoid valve at a rate that is related to the speed of the
	application equipment; the faster the applicator, the shorter the
	valve closed time. When using mechanical application equipment,
	calibrate the system to deliver the intended amount of material per

	acre (e.g. 150 grams of SPLAT LBAM LD). For unusual situations,	
	always consult your dealer, distributor or ISCA Technologies to	
	ensure correct and optimum application of product is achieved.	
MONITORING	Monitor the target pest(s) with pheromone-baited traps, such as the	
	ISCAlure™ LBAM pheromone lure and ISCA delta trap monitoring	
	system: Place monitoring traps in the treated area before applying	
	SPLAT LBAM LD to determine the infestation level. To prevent	
	contaminating the monitoring traps with SPLAT LBAM LD either	
	cover the traps or remove the traps before treating the area; then	
	following application, uncover or replace the traps. Continue	
	monitoring throughout the season to evaluate efficacy and longevity	
	of the application. Reapply as needed based on trap capture and	
	field scouting programs.	
WEATHER	SPLAT LBAM LD should be applied when the ambient temperature	
FACTORS	is above 55°F and below 95°F. SPLAT dollops typically cure within 2-	
	3 hours following application, after which it will become rainfast and	
	UV resistant. Do not apply if rain is expected within 2-3 hours of	
	application or the temperature is outside these ranges.	
CLEANING	Clean equipment with water and citrus or limonene detergents.	
EQUIPMENT		
	STORAGE AND DISPOSAL	
. Do n	ot contaminate water, food or feed by storage or disposal.	
STORAGE	Store product sealed and refrigerated, if possible. If refrigerated, do	
	not store with food. If refrigeration is not available, store product in	
	a cool dry place, out of direct sunlight. Do not exceed 75 F for long-	
	term storage. Avoid freezing. In case of leak or spill, wipe with paper	
	towel and dispose of waste when product hardens.	
PESTICIDE	Wastes resulting from the use of this product may be disposed of on	
DISPOSAL	site or at an approved waste facility.	
CONTAINER	Nonrefillable container: Do not refill or reuse this container. Triple	
DISPOSAL	rinse (or equivalent) promptly after emptying. Then offer for	
	recycling if available, or puncture and dispose of in a sanitary landfill,	
	or by incineration, or, if allowed by state and local authorities by	
	burning. If burned, stay out of smoke.	
WARRANTY AND LIMITATION OF DAMAGES		
Read all directions carefully. All statements concerning the use of this product apply		
only when used as directed. ISCA Technologies warrants that this material conforms to		
the specifications on the label and is reasonably fit for the intended purpose referred to		
on the label. To the extent consistent with applicable law, ISCA Technologies makes no		
other express or implied warranty of merchantability or fitness for a particular purpose		
or any other express or implied warranty		

ISCA Technologies, Inc. 1230 Spring Street Riverside, CA 92507 (951) 686-5008

or any other express or implied warranty.