



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

August 19, 2015

Jonathan A. Janis  
Isagro USA, Inc.  
430 Davis Dr., Suite 240  
Morrisville, NC 27560

Subject: Label Amendment – Formatting changes to directions for use on soybeans and minor label revisions  
Product Name: DOMARK 230 ME FUNGICIDE  
EPA Registration Number: 80289-7  
Application Date: 05/05/2015  
Decision Number: 504969

Dear Mr. Janis:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 80289-7  
Decision No. 504969

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, you may contact Maryam K. Muhammad at 703-347-0301 or via email at [Muhammad.maryam@epa.gov](mailto:Muhammad.maryam@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written over a circular scribble.

Hope Johnson, Product Manager 21  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure: Accepted product label for Domark 230 ME Fungicide

**DOMARK® 230ME FUNGICIDE**

**For Control and / or Suppression of Certain Diseases in Corn and Soybean**

**Active Ingredient:**

Tetraconazole\* ..... 20.5%

**Other Ingredients:** ..... 79.5%

Total ..... 100.0%

\*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2,-tetrafluoroethoxy)propyl]1H-1,2,4-triazole

DOMARK 230 ME FUNGICIDE is a micro emulsion containing 1.9 pounds of tetraconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION / PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

See booklet for additional Precautions and Directions for Use

<b>FIRST AID</b>	
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have affected person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Health Emergency or Spill (24 hr) call 800-892-0099 or for Transportation Chemical Emergency Spill Leak Fire Exposure or Accident Call CHEMTREC Day or Night 800-424-9300 or 202-483-7616</p>	


EPA Registration No.: 80289-7

EPA Establishment No.:

Batch Code will be placed on the container

**NET CONTENTS: \_\_\_\_\_ Gallons**

Manufactured by Isagro SpA for:  
Isagro USA, Inc.,  
430 Davis Drive, Suite 240  
Morrisville, NC 27560



Distributed by:  
Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366-5569



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, and viton  $\geq$  14 mils.

**Applicators and other handlers must wear:**

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl rubber, and Viton) made of any water proof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise caution when making applications of DOMARK 230 ME FUNGICIDE, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment wash waters or rinsate.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.  
Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Do not** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all activities with the exception of 20 days for detasseling corn grown for seed.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any water proof material
- Shoes plus socks

**PRODUCT INFORMATION**

DOMARK 230 ME FUNGICIDE is formulated as a 1.9 pound active ingredient per gallon micro emulsion (ME). The active ingredient in DOMARK 230 ME FUNGICIDE is tetraconazole, a triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. Tetraconazole is absorbed quickly into the plant tissue and like all triazoles can move up, but not down the plant. Optimal disease control is achieved when

DOMARK 230 ME FUNGICIDE is applied in a regularly scheduled spray program. Preventive applications optimize disease control, and may result in improved plant health [and] [beneficial physiological effects].

When using DOMARK 230 ME FUNGICIDE in combination and/or rotation with other fungicides, it is important to use fungicides that have different modes of action (i.e. non Group 3 fungicides).

**MODE OF ACTION:** The active ingredient in DOMARK 230 ME FUNGICIDE is tetraconazole, which belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the Fungicide Resistance Action Committee (FRAC) as Target Site of Action Group 3 fungicides.

**CROP USE RATES AND TIMING OF APPLICATIONS**

Field Corn, Popcorn, Corn Grown For Seed Production				
Disease	Dosage Rate		When to Apply	Specific Use Instructions
	Fl. Oz./A	GPA		
Gray leaf spot ( <i>Cercospora zeae-maydis</i> )	Single application 4.0 to 6.0 fl oz/a (0.060 to 0.090 lb. ai./A)	Ground: Minimum of 10 GPA	Early Application (V4 – V8)  OR  Late Application VT – R3  Apply prior to disease onset when conditions favor disease development.  Curative applications are most effective when disease incidence does not exceed 5% of the plants at time of application	DOMARK 230 ME FUNGICIDE may be applied for early or late season disease control and may result in [improved plant health] [and] [beneficial physiological effects]. If mixing with herbicides other than solo glyphosate products, Yukon®, Halex® GT, Callisto®, Ignite®, Laudis®, Lexar®, Lumax®, Status or Resolve® Q, consult your local Gowan representative. If disease pressure develops later in the season, an application of an alternate corn fungicide should be made at VT – R3 to provide season-long control.  Use DOMARK 230 ME FUNGICIDE as part of an integrated pest management program (IPM).  Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of plants. To limit the potential for resistance development, do not apply more than twice per year or exceed 6 fl oz per acre per year.
Rust, common ( <i>Puccinia sorghi</i> )		Aerial: Minimum of 2 GPA		
Rust, southern ( <i>Puccinia polysora</i> )				
Anthracnose leaf blight ( <i>Colletotrichum graminicola</i> )				
Eye spot ( <i>Aureobasidium zeae</i> )				
Northern corn leaf blight ( <i>Exserohilum turcicum</i> )				
Northern corn leaf spot ( <i>Bipolaris zeicola</i> )				
Physoderma brown spot ( <i>Physoderma maydis</i> )				
Southern corn leaf blight ( <i>Bipolaris maydis</i> )				
Yellow leaf blight ( <i>Phyllosticta maydis</i> )				

- RESTRICTIONS AND LIMITATIONS**
1. Do not make more than (2) applications per year.
  2. Do not apply more than 6 fl oz (0.090 lb ai tetraconazole) of DOMARK 230 ME FUNGICIDE per acre per year.
  3. Do not apply DOMARK 230 ME FUNGICIDE after corn growth stage R3 (brown silk/milk).
  4. Do not use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage. A compatibility agent, another fungicide, or an insecticide may be included if needed and labeled for use in corn. Refer to adjuvant product label for specific use directions and restrictions. Always follow the more restrictive label.

<b>Soybean</b>				
Disease	Dosage Rate		When to Apply	Specific Use Instructions
	Fl. Oz./A	GPA		
Asian Soybean Rust <i>(Phakopsora pachyrhizi)</i> Cercospora Blight <i>(Cercospora kikuchii)</i> Purple Seed Stain <i>(Cercospora kikuchii)</i> Frogeye Leaf Spot <i>(Cercospora sojina)</i> White Mold/Sclerotinia Stem Rot <i>(Sclerotinia sclerotiorum)</i> Powdery Mildew <i>(Microsphaera diffusa)</i> Brown Spot <i>(Septoria glycines)</i> Anthracnose <i>(Colletotrichum spp.)</i>	4.0 to 5.0 (0.060 to 0.075 lb. ai./A)	Ground: Mimimum of 10 GPA  Aerial: Minimum of 2 GPA; (5 GPA for White Mold and Asian Soybean Rust)	<u>Asian Soybean Rust:</u> Apply prior to disease development when rust infections are likely to occur. If necessary repeat with a second application before growth stage R-6. <u>All Other Soybean Diseases:</u> Make application at soybean growth stage R-1 (early pod fill) or when conditions are favorable for disease development. Repeat application 15 to 21 days after first application if disease pressure is heavy. Under severe disease conditions the higher rate and shorter spray intervals should be used. Curative applications are most effective when disease incidence does not exceed 5% of the soybean plants at time of application.	Use DOMARK 230 ME FUNGICIDE as part of an integrated pest management program (IPM). Apply as a foliar spray or via chemigation in sufficient water to obtain thorough coverage of soybeans.
<b>RESTRICTIONS AND LIMITATIONS</b> <ol style="list-style-type: none"> <li>1. Do not make more than two (2) applications per year.</li> <li>2. Do not apply more than 10 fl oz (0.150 lb ai tetraconazole) of DOMARK 230 ME FUNGICIDE per acre per year.</li> <li>3. Do not graze or feed DOMARK 230 ME FUNGICIDE-treated forage or hay to livestock.</li> <li>4. Do not apply DOMARK 230 ME FUNGICIDE after soybean growth stage R5 (beginning seed).</li> <li>5. Do not harvest immature soybeans for consumption once plants are treated with DOMARK 230 ME FUNGICIDE.</li> <li>6. Do not use on vegetable soybean varieties grown for their immature pods.</li> </ol>				

## RESISTANCE MANAGEMENT

DOMARK 230 ME FUNGICIDE contains tetraconazole, a Group 3 fungicide (sterol biosynthesis inhibitors), and is effective against labeled pathogens resistant to fungicides with modes of action different from those of target site Group 3, such as dicarboximides, strobilurins, benzimidazoles, or phenylamides. However, fungal isolates resistant to Group 3 fungicides may eventually dominate the fungal population if Group 3 fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species, especially if resistance to Group 3 fungicides is already present in the pathogen population. This may result in reduced disease control by Group 3 fungicides. To maintain the performance of DOMARK 230 ME FUNGICIDE in the field, do not exceed the total number of sequential applications of DOMARK 230 ME FUNGICIDE and the total number of applications of DOMARK 230 ME FUNGICIDE per year stated in "CROP USE RATES AND TIMING OF APPLICATIONS". Adhere to the label instructions regarding the consecutive use of DOMARK 230 ME FUNGICIDE or other target site of action Group 3 fungicides that have a similar site of action on the same pathogens. Consider the following to delay the development of fungicide resistance:

1. **Tank mixtures:** If DOMARK 230 ME FUNGICIDE is used in tank mixtures with fungicides from different mode of action Groups that are registered for the same use and that are effective against the pathogens of concern, use at least the minimum labeled rates of each fungicide in the tank mix.
2. **IPM:** Integrate DOMARK 230 ME FUNGICIDE into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or [Gowan] [Isagro] representative for additional IPM strategies established for your area. Use DOMARK 230 ME FUNGICIDE in Agricultural Extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
4. **Reporting:** If a Group 3 target site fungicide appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact your [Gowan] [Isagro] representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

## RAINFASTNESS

DOMARK 230 ME FUNGICIDE is rainfast 2 hours after application. Do not apply if rain is expected within 2 hours of application or disease control may be reduced.

## SPRAYER PREPARATION

Before applying DOMARK 230 ME FUNGICIDE, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply DOMARK 230 ME FUNGICIDE. If two or more products were tank mixed prior to DOMARK 230 ME FUNGICIDE application, follow the most restrictive cleanup procedure.

## MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the DOMARK 230 ME FUNGICIDE to the spray tank. Agitation should create a rippling or rolling action on the water surface.
3. If tank-mixing DOMARK 230 ME FUNGICIDE with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Stickers, spreaders, etc., should be added last.

When tank mixing this product with other pesticides observe the more restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

4. Fill spray tank to desired level with water. Agitation should continue until all spray solution has been applied.
5. Mix only the amount of spray solution that can be applied the day of mixing. DOMARK 230 ME FUNGICIDE should be applied within 24 hours of mixing.
6. Do not combine DOMARK 230 ME FUNGICIDE in a sprayer tank with pesticides or fertilizers, unless your prior use has shown the combination to be physically compatible, effective and non-injurious under your conditions of use.

## **JAR TEST TO DETERMINE COMPATIBILITY OF DOMARK 230 ME FUNGICIDE**

Perform a jar test before mixing commercial quantities of DOMARK 230 ME FUNGICIDE when using DOMARK 230 ME FUNGICIDE for the first time, or when a new water source is being used.

1. Add 1 pt. of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 ml of DOMARK 230 ME FUNGICIDE to the quart jar; gently mix until product goes into suspension.
3. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
4. An ideal tank-mix combination will be uniform and free of suspended particles.

### **APPLICATION EQUIPMENT**

Application equipment must be clean and in good condition. Frequently check nozzles for accuracy.

### **SPRAYER CLEANUP**

Clean spray equipment each day following DOMARK 230 ME FUNGICIDE application. After DOMARK 230 ME FUNGICIDE is applied, use the following steps to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

Thoroughly clean spray equipment, including all tanks, hoses, booms, screens and nozzles, before it is used to apply foliar pesticides.

### **SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, and chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Avoiding spray drift at the application site is the responsibility of the applicator.

#### **Droplet Size**

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### **Wind Speed**

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

#### **Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### **Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

#### **Equipment**

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

### **AERIAL APPLICATION**

To avoid drift, apply the largest droplet size possible that will provide uniform coverage and result in satisfactory disease control. To obtain satisfactory application and avoid drift, the following directions must be observed:

Do not apply during low-level inversion conditions, when winds are gusty or under other conditions that favor drift. Application should be avoided when wind velocity is less than 2 mph and more than 15 mph.

#### **Carrier Volume and Spray Pressure:**

- For aerial application use a minimum of 2 gallons per acre for all diseases except rust and white mold/Sclerotinia stem rot of soybeans for which a minimum of 5 gallons per acre must be used. Increasing the spray volume to 7 gallons or more per acre generally provides better coverage and more consistent disease control.



- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

#### **Nozzle Selection and Orientation:**

Minimize formation of very small drops by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles, such as diaphragm type nozzles, to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft, at an angle between 0 and 15° downward. **Do not** place nozzles on the outer 25% of the wings or rotors.

### **CHEMIGATION INSTRUCTIONS**

DOMARK 230 ME FUNGICIDE may be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

- Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other irrigation experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

#### **Requirements for Chemigation Systems Connected to Public Water Systems**

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favor drift beyond the area intended for treatment.

#### **Sprinkler Chemigation**

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid

from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

**DOMARK 230 ME FUNGICIDE TANK MIX INFORMATION**

**Use Restrictions:**

1. Always read and follow all label directions when using any pesticide alone or in tank-mix combinations.
2. The most restrictive labeling applies when using a tank-mix.

**DOMARK 230 ME FUNGICIDE may be tank-mixed with the following products, including but not limited to:**

Herbicides-Corn	Herbicides-Soybeans	Fungicides-Corn and Soybean	Insecticides-Corn and Soybean	Miticides-Corn
Callisto®	Liberty® (LibertyLink®)	Headline®	Asana®	Onager®
Halex® GT	Roundup® (Roundup Ready®)	Quadris®	Baythroid®	
Laudis®	Targa®		Justice® (Soybeans Only)	
Lexar®			Lorsban®	
Liberty® (LibertyLink®)			Mustang Max®	
Lumax®			Orthene®	
Roundup® (Roundup Ready®)			Pounce®	
Yukon®			Proaxis®	
			Warrior®	

**ROTATIONAL CROP RESTRICTIONS**

Use the time intervals listed below to determine the minimum required time interval between the last DOMARK 230 ME FUNGICIDE application and new crop planting.

Rotational Crop Guideline	
Crop	Time Interval in Days Before Planting
Corn, grape, peanut, pecan, soybean, strawberry and sugarbeet	0
Small Grains: Barley, buckwheat, millet, oats, rice, rye, triticale, and wheat	45
Sugarcane	45
All Other Crops	120

## STORAGE AND DISPOSAL

**Do not** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry, temperature-controlled, secure, place.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### **CONTAINER HANDLING:**

**For rigid, non-refillable containers (2.5 to 5 gallons):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of resulting smoke.

**For rigid, non-refillable containers that are too large to shake (with capacities greater than 5 gallons):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of resulting smoke.

### **PRESSURE RINSE PROCEDURE (all sizes):**

Pressure rinse as follows: Empty the remaining contents into application equipment or a tank mix and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For rigid, refillable containers:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## LIMITATION OF WARRANTY AND LIABILITY

Read the entire label before using this product, including this Limitation of Warranty and Liability.

If the terms are not acceptable, return the product at once unopened for a refund of the purchase price.

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks described below, when used in accordance with the Directions for Use under normal conditions.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ISAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Buyers and Users of this product must be aware that there are inherent unintended risks associated with the use of this product, independent from the control of Isagro. These risks include, but are not limited to, weather conditions, soil factors, moisture conditions, irrigation practices, condition of the crop at the time of application, materials which are present in the tank mix with this product or prior to the application of it, cultural practices or the manner of use or application, all risks which are impossible to eliminate. The Buyers and Users must be aware that these factors may cause: ineffectiveness of the product, reduction of harvested yield of the crop (entirely or partially), or crop injury.

If the Buyer does not agree with the acceptance of these risks, then **THE PRODUCT SHOULD NOT BE APPLIED**. To the extent consistent with applicable law, the Buyer acknowledges and accepts these inherent unintended risks and **AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER**.

To the extent consistent with applicable law, ISAGRO or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise). To the extent consistent with applicable law, the exclusive remedy of the User or Buyer and the exclusive Liability of Isagro or Seller shall be the return of the purchase price of the product, or at the election of Isagro or Seller, the replacement of the product.

To the extent that applicable law allows such requirements, Isagro or its Seller must have prompt notice of any claim so that an immediate inspection of Buyer's or User's growing crops can be made. To the extent consistent with the applicable law, if Buyer and User do not notify Isagro or Seller of any claims, in proper time, it shall be barred from obtaining any remedy.

To the extent consistent with applicable law, Buyers and Users are deemed to have accepted the terms of this Limitation of Warranty and Liability, which may not be modified by any verbal or written agreement.

Domark is a registered trademark of Isagro SpA Corp.

Asana<sup>®</sup>, Baythroid<sup>®</sup>, Callisto<sup>®</sup>, Halex<sup>®</sup> GT, Headline<sup>®</sup>, Ignite<sup>®</sup>, Justice<sup>®</sup>, Laudis<sup>®</sup>, Lexar<sup>®</sup>, Liberty<sup>®</sup>, LibertyLink<sup>®</sup>, Lorsban<sup>®</sup>, Lumax<sup>®</sup>, Mustang Max<sup>®</sup>, Onager<sup>®</sup>, Orthene<sup>®</sup>, Pounce<sup>®</sup>, Proaxis<sup>®</sup>, Quadris<sup>®</sup>, Resolve<sup>®</sup> Q, Roundup<sup>®</sup>, Status<sup>®</sup>, Targa<sup>®</sup>, Warrior<sup>®</sup> and Yukon<sup>®</sup> are trademarks of their respective owners.

AMEND5May15