

80289-12

02/04/2009

1/25



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

80289-12

Date of Issuance:

FEB 4 2009

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

Badge X2

Name and Address of Registrant (include ZIP Code):

ISAGRO USA, Inc.
430 Davis Drive, Suite 240
Morrisville, NC 27560

Attention: Mel Graben

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA Section 4.

2. Submit the following conditional data before the due date of 2/4/2010 – One-year GLP storage stability (830.6317) and corrosion characteristics (830.6320) study.

3. Not all the required information to support the claim "For Organic Production" was submitted with the application. This claim is conditionally approved pending (A) receipt within 45 days of the following information: Confidential Statements of Formula (CSF) and detailed descriptions of the manufacturing process for each source of active ingredient; basic CSF, stamped label and this Reg Notice for Badge X2; and (B) that EPA finds the information supports the claim.

3. Submit two copies of a final printed label within 30 days from the date of this notice which makes the following changes:

Signature of Approving Official:

Tony Kish
Product Manager, Team 22
Fungicide Branch
Registration Division (7505P)

Date:

FEB 4 2009

A. On page 2 and 3 after "Protective eyewear" add "(goggles, safety glasses, or face shield)"

B. For potato, grape, atemoya, the RED allows higher maximum application rates than listed on this label. You can change these rates to match the RED if desired.

C. On page 17, delete the entire Live Oak (ball moss) section because it is covered by the "Ball Moss" section on page 22.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with comments" is enclosed for your records.

Enclosure -stamped label

Badge[®] X₂

Dry Flowable Fungicide/Bactericide

For Agricultural Use

ACCEPTED
with COMMENTS
In EPA Letter Dated

FEB 4 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

803 99-12

ACTIVE INGREDIENTS:

Copper Oxychloride (CAS No. 1332-40-7)* 24.6%

Copper Hydroxide (CAS No. 20427-59-2)* 22.9%

OTHER INGREDIENTS: 52.5%

TOTAL: 100.0%

*Metallic Copper (Cu²⁺) Equivalent is 28% by weight

KEEP OUT OF REACH OF CHILDREN

WARNING – AVISO

See Label for Additional Precautions and Directions for Use

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if unable to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>NOTE TO PHYSICIAN: Possible mucosal damage may contraindicate use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.</p> <p>For Chemical Emergency Spill Leak Fire Exposure or Accident Call CHEMTREC Day or Night Domestic North America 800-424-9300 International 703-527-3883 (collect calls accepted)</p>	

EPA Registration No.: 80289-12

EPA Establishment No.: 79558-ITA-1



Manufactured by Isagro SpA for:

Isagro USA, Inc.

430 Davis Drive, Suite 240, Morrisville, NC 27560



FOR ORGANIC PRODUCTION

NET CONTENTS:

4
25**PRECAUTIONARY STATEMENTS****WARNING – AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if inhaled. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Remove and wash contaminated clothing before reuse. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as Natural Rubber, Selection Category A). Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, and other handlers must wear the following:

- long-sleeved shirt and long pants
- shoes plus socks
- protective eyewear
- chemical-resistant gloves such as Natural Rubber

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA. Do not contaminate water when disposing of equipment washwaters or rinsate.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Certain water conditions including low pH (≤ 6.5), low dissolved organic carbon (DOC) levels (3.0 mg/L or lower), and "soft" waters (i.e. alkalinity less than 50 mg/L), increases the potential acute toxicity to non-target aquatic organisms.

5
25**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers, adults, children or pets, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **48** hours without required PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear if overhead exposure
- Protective eyewear
- Chemical-resistant apron when mixing, loading, cleaning equipment or spills, or otherwise exposed to the concentrate

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not allow adults, children, or pets to enter until dusts have settled.

6
25

GENERAL INSTRUCTIONS

BADGE X₂ may be applied as an aerial, ground dilute or ground concentrate spray unless specifically directed otherwise in the specific crop use directions.

The per acre use rate of BADGE X₂ is applicable for both dilute and concentrate spraying. Depending upon the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to Minimum Recommended Spray Volume Table. Complete spray coverage is essential to assure optimum performance from BADGE X₂. When treating by aerial application or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

Consult the BADGE X₂ label for specific rates and timing of application by crop. **Where application rates and intervals are provided in a range (e.g. 4 to 12 pounds and 7 to 10 days), the higher rates and shorter spray intervals are recommended when rainfall is heavy and/or disease pressure high. Use the higher rates for large mature tree crops.**

SPECIAL PRECAUTIONS

- BADGE X₂ should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.
- Do not tank mix BADGE X₂ with Alette® fungicide for use on any registered crops or ornamentals unless appropriate precautions have been taken to buffer the spray solution because severe phytotoxicity may result. Use in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.
- This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
- Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of BADGE X₂ resulting in possible phytotoxicity or loss of effectiveness.

•Agricultural chemicals may perform in an unpredictable manner when tank mixed, especially where several products are involved. Reduced effect on pests or crop injury may occur. Unless recommended on this label or by a state/local expert, it is advisable to test for compatibility and potential crop injury prior to commercial use of a new tank mix; otherwise tank mixing should not be undertaken.

•It must be determined if proper application equipment is available and if waste associated with its use can be properly handled. Agricultural chemicals are often reactive with the materials used in the construction of application equipment such as aluminum, rubber and some synthetic materials. This factor should be taken into consideration when selecting proper application equipment. It is necessary that all application equipment be thoroughly flushed with clean water after each day's use.

•Do not apply this product through any irrigation (chemigation) system using aluminum parts or components as damage to the system may occur. Such application is prohibited regardless of whether the irrigation system is flushed with water after use of this product.

•Apply this product only through one or more of the following types of systems: sprinkler, including center pivot, lateral move, traveler, big gun, or plastic pipe solid set system(s) which contain no aluminum parts or components. Do not apply this product through any other type of irrigation system.

•While volume is important in obtaining full spray coverage, often factors such as foliage density, environmental conditions and sprayer calibration have a greater impact. Always be sure that sprayers are calibrated to spray equipment manufacturer's specifications and environmental conditions are within those recommended by State and local regulatory authorities.

•When mixing, fill the spray tank one-half full with water. Add BADGE X₂ slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. **DO NOT PREMIX or SLURRY** BADGE X₂. Spreaders, stickers, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the Compatibility Jar Test before mixing a whole tank or contact your chemical supplier. Observe all precautions and limitations on the labels of all products used in mixtures.

FROST INJURY PROTECTION (Bacterial Ice Nucleation Inhibitor)

Application of BADGE X₂ made to all crops listed on this label at the rates and stages of growth indicated, at least 24 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola* and *Pseudomonas fluorescens*) and may therefore provide some protection against light frost. Not recommended for those geographical areas where weather conditions favor severe frost.

RECOMMENDED CROP USES

CITRUS: Grapefruit, Kumquat, Lemon, Lime, Orange, Pummelo, Tangelo and Tangerine.

FIELD CROPS: Alfalfa, Barley, Corn*, Oats, Peanut, Potato, Sugar Beet and Wheat.

SMALL FRUITS: Blackberry, Blueberry*, Cranberry, Currant, Gooseberry, Raspberry and Strawberry.

TREE CROPS: Almond, Apple, Apricot, Avocado, Banana, Cacao, Cherry, Coffee, Filbert, Mango*, Nectarine, Olive, Peach, Pear, Pecan, Pistachio, Plum, Prune, Quince* and Walnut.

VEGETABLES: Bean, Beet, Beet Greens, Broccoli, Brussels Sprout, Cabbage, Cantaloupe, Carrot, Cauliflower, Celeriac*, Celery, Cucumber, Eggplant, Greens (Collard, Mustard and Turnip), Honeydew, Muskmelon, Onion/Garlic, Pea, Pepper, Pumpkin, Spinach, Squash, Tomato, Watercress* and Watermelon.

VINES: Grape, Hops and Kiwi.

MISCELLANEOUS: Atemoya*, Carambola*, Chives*, Dill*, Ginseng, Guava, Litchi*, Live Oak*, Macadamia, Mamey Sapote*, Papaya*, Parsley*, Passion Fruit*, Sugar Apple* and Sycamore.

GREENHOUSE AND SHADEHOUSE CROPS: BADGE X₂ may be used in greenhouses and shadehouses to control diseases on any crop on this label where physiology allows greenhouse or shadehouse culture. While specific directions are presented for Citrus, Cucumber, Eggplant, Pepper, and Tomato; general use may occur for any crop on this label where physiology allows greenhouse or shadehouse culture.

ORNAMENTALS: Specified as listed.

*Except California

MINIMUM RECOMMENDED SPRAY VOLUME (GALLONS PER ACRE) WHEN APPLYING BADGE X₂

USE	AERIAL	DILUTE	CONCENTRATE
Vegetables	3	20	—
Field Crops	3	20	—
Small Fruits	5	150	50
Vines	5	150	50
Tree Crops	10	400	50
Miscellaneous crops	10	150	50
Citrus	10	800	100*
Ornamentals	10	100	50

*When using pesticide application equipment such as Curtec® or other similar sprayers which are capable of obtaining thorough coverage at low volumes, application rates as low as 20 gallons per acre of spray volume may be used.

GENERAL USE RECOMMENDATIONS

The following specific instructions are based on general application procedures. The recommendations of your local State Agricultural Extension Service should be closely followed as to timing, frequency and number of sprays per season.

CITRUS

BADGE X₂ may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. BADGE X₂ per acre rates in these mixes must not exceed the maximum recommended label rates for disease control. Adding foliar nutritionals or other products to spray mixtures containing BADGE X₂ and applying to citrus during the post-bloom period when young fruit are present may result in spray burn.

DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MIN. RETREATMENT INTERVAL (DAYS)	COMMENTS
Algal Spot, Melanose, Scab	1.75-5	3.15	12.6	7	Apply as pre-bloom and post-bloom sprays. Use the higher rates when conditions favor disease development.
Greasy Spot, Pink Pitting	0.75-2.5	2.4	12.6	7	Apply in summer on expanded new flush. Repeat on subsequent flushes where disease pressure is severe. Use the higher rates when conditions favor disease development.
Alternaria Brown Spot	1.75-3.5	3.15	12.6	21	On susceptible varieties apply when the first spring flush appears and each flush thereafter. Application to fruit should start after two thirds of the petals have fallen and be repeated on a 21 day schedule. Use the higher rates when conditions favor disease development.
Phytophthora Brown Rot, Septoria Spot	1.75-3.5	3.15	12.6	7	Begin application in fall before or just after the first rain and continue as needed. For brown rot only, apply to skirts of trees to a height of at least 4 feet. For control of septoria spot or where fruit have already been infected with brown rot, apply to entire tree. Apply also to bare ground 1 foot beyond skirt. Use the higher rates when conditions favor disease development. NOTE: In California, in areas subject to copper injury, add 1/3 to 1 pound of high quality lime per pound of BADGE X ₂ .

8
25

DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MIN. RETREATMENT INTERVAL (DAYS)	COMMENTS
Phytophthora Foot Rot	0.5	0.4	12.6	7	Mix with 1 quart of water, Tre-Hold® or latex paint. Paint trunks of trees from the soil surface to the lowest scaffold limbs. Apply in May prior to summer rains and/or in the fall prior to wrapping trees for freeze protection. Treatment serves as protection for up to 1 year, but does not cure existing infections. NOTE: Areas where microjet or low volume irrigation hit the tree trunk may require retreatment due to wash off.
Citrus Canker (Suppression)	2-5	3.15	12.6	7	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require an additional application. Number and timing of applications will be dependent upon disease pressure. Under heavy pressure, each flush of new growth should be sprayed.

NOTE: Phytotoxicity may occur on young tender flush when BADGE X₂ is applied to citrus seedlings grown in greenhouses or shadehouses.

CITRUS (FIELD NURSERY GROWN)

To control Melanose, Scab, Pink Pitting, Greasy Spot and Brown Rot and for suppression of Citrus Canker, apply 1.75 to 3.5 pounds of product per acre. Apply BADGE X₂ at 28 day intervals or as needed depending on disease severity and rainfall. The maximum single application rate is 3.15 pounds of Cu²⁺ per acre. The maximum annual application rate is 12.6 pounds of Cu²⁺ per acre. The minimum retreatment interval is 7 days.

FIELD CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	0.75	0.53	1.12	30	Apply 10 to 14 days before each harvest or earlier if disease threatens. NOTE: Spray injury may occur with sensitive varieties such as Lahontan.
Corn* (Field Corn, Popcorn, Sweet Corn)	Bacterial Stalk Rot	0.5-1.75	1.05	4.2	7	Begin treatment when disease first appears and repeat every 7 to 10 days or as needed. Use the higher rates and shorter spray intervals when conditions favor disease development.
Peanut	Cercospora Leaf Spot	0.75-1.25	0.79	4.74	7	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 10 to 14 day intervals or as needed. Reduce sprays to 7 day intervals during humid weather. Use the higher rates when conditions favor disease development. Flowable sulfur may be added.

9
25**FIELD CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Potato	Early Blight, Late Blight	0.5-1.75	1.6	25	5	Apply 0.5 to 1.75 pounds at 7 to 10 day intervals or as needed starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 1.75 pounds per acre when disease is more severe. Under conditions of severe disease, control with BADGE X ₂ will be improved by tank mixing with other compatible fungicides registered for use on potatoes. Read and follow all label instructions of tank mix partners. Use the higher rates when conditions favor disease development.
Sugar Beet	Cercospora Leaf Spot	0.75-2	1.31	7.86	10	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals or as needed. Use the higher rates when conditions favor disease development. Addition of a spreader/sticker is recommended.
Wheat, Barley, Oats	Helminthosporium Spot Blotch, Septoria Leaf Blotch	0.5-0.75	0.53	1.06	10	Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease development. BADGE X ₂ can be applied as a foliar application for early season disease control and again at early heading and followed with another application 10 days later.

*Except California

SMALL FRUITS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Brambles (Aurora, Blackberry, Boysen, Cascade, Chehalem, Logan, Marion, Raspberry, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	1.75	2.0	10	7	Make fall application after harvest. Apply delayed dormant spray after pruning/training in the spring. If needed, agricultural-type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	0.75	0.8	10	7	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural-type spray oil may be added. NOTE: Crop injury may occur if applied to foliage under certain environmental conditions such as hot or prolonged moist periods. Discontinue application if signs of crop injury appear.

10
25**SMALL FRUITS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Blueberry*	Bacterial Canker	1.75-3.5	2.1	8.4	28	Make first application before rain falls and a second application 4 weeks later. Use the higher rates when conditions favor disease development.
	Fruit Rot, Phomopsis Twig Blight	1.0-2.25	2	8.4	7	Dormant Application: Begin applications when bloom buds begin to swell. Make additional applications at 10 to 14 day intervals or as needed before blooms open. Use the higher rates when conditions favor disease development
Cranberry	Fruit Rot	3.5	2.1	6.3	7	Make first application in late bloom. Apply one or two additional applications at 10 to 14 day intervals or as needed depending on disease severity.
	Rose Bloom	3.5	2.1	6.3	7	Apply three sprays on 10 to 14 day schedule or as needed as soon as symptoms are observed.
	Bacterial Stem Canker	3.5	2.1	6.3	7	Apply postharvest and again in spring at bud swell. Apply one or two additional applications at 10 to 14 day intervals or as needed depending on disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight (<i>Monilinia</i>)	3.5	2.1	6.3	7	Apply delayed dormant spray in the spring. Repeat at 10 to 14 day intervals or as needed through pre-bloom.
Currant, Gooseberry	Anthrachnose, Leaf Spot	4.25	2.5	10	10	Make initial application after first leaves have expanded. Continue on a 10 to 14 day schedule or as needed during wet conditions in the spring. Make an additional application after harvest.
Strawberry	Angular Leaf Spot, (<i>Xanthomonas</i>), Leaf Blight, Leaf Scorch, Leaf Spot	0.75-1.25	1.2	8.19	7	Begin application when plants are established and continue on a weekly schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease development. NOTE: Discontinue applications if signs of crop injury appear.

* Except California

TREE CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Almond, Apricot, Cherry, Plum, Prune	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Coryneum Blight (Shot Hole)	3.5-7.0	6.4	18	7	<p>Make first application before fall rains and a second at late dormant. Use the higher rates when conditions favor disease development. If needed, agricultural-type spray oil may be added.</p> <p>For cherries: Where disease is severe, an additional application shortly after harvest may be required. Use the higher rates when conditions favor disease development.</p> <p>Almond only: For bacterial blast control in sprinkler irrigated orchards or where disease is severe, apply 0.5 pounds per acre post-bloom at 2 week intervals or as needed or just before sprinkling. NOTE: Foliar injury may occur from post-bloom sprays on almonds, especially on NePlus varieties.</p>
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	2.5-3.5 (Almond) 3.5-5.0 (all others)	1.5	18	5	<p>Apply during early bloom. Do not apply after full bloom or injury may occur. Use the higher rates when rainfall is heavy and disease pressure is high.</p>
	Black Knot (Plum)	1.75-3.5	1.5	18	5	<p>Make an application at bud swell up to early bloom for early season disease suppression. Apply before full bloom. Use the higher rates when rainfall is heavy and disease pressure is high.</p> <p>NOTE: To avoid plant injury, do not use after full bloom. Use the higher rates when conditions favor disease development.</p>
	Cherry Leaf Spot (Sour Cherries Only)	2.25-3.5	1.5	18	5	<p>Apply at petal fall as well as one to two times after petal fall. Use the lower rates where disease infection is light and use the higher rates for a dormant application or where disease infection is moderate to heavy. Do not apply to sweet cherry or the English Morello variety as severe injury will result. The addition of 1 to 3 pounds of hydrated lime per pound of BADGE X₂ may reduce crop injury.</p> <p>NOTE: Moderate to severe injury such as leaf spotting and defoliation may occur from post-bloom applications.</p>
Apple	Anthrachnose, Blossom Blast, European Canker (<i>Nectria</i>), Shoot Blast (<i>Pseudomonas</i>)	5.25-7	6.4	16	n/a	<p>Apply before fall rains. Use the higher rate when conditions favor disease development. NOTE: Use on yellow varieties may cause discoloration. To avoid discoloration, pick before spraying.</p>

12
25**TREE CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
	Apple Scab, Fire Blight	1.75	0.5	16	5	Make application between silver-tip and green-tip. Apply as a full-cover spray for early season disease suppression. NOTE: Moderate to severe crop injury may occur from late application; discontinue use when green-tip reaches ½ inch.
	Apple Scab	1.25	0.5	16	5	Extended spray schedule where fruit finish is not a concern: Continued application may be made at 5 to 7 day intervals or as needed between ½ inch green-tip and first cover spray. NOTE: Moderate to severe crop injury may result from this extended spray schedule. It is not intended for fresh market apples or fresh apples where fruit finish is a concern as it is likely to cause fruit russetting. The addition of 1 to 3 pounds of hydrated lime per pound of BADGE X ₂ may reduce crop injury.
	Fire Blight	1-1.25	0.5	16	5	
	Collar Rot, Crown Rot	1.75	0.5	16	5	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply in early spring or in fall after harvest for best results. Do not apply to foliage or fruit. NOTE: Do not use if soil pH is below 5.5 since copper toxicity may result.
Avocado	Anthrachnose, Blotch, Scab	3.5-5.25	3.15	18.9	14	Apply when bloom buds begin to swell and continue application at monthly intervals for five to six applications. Use the higher rates when conditions favor disease development.
Banana	Sigatoka (Black and Yellow)	0.75	0.8	18.9	7	Apply by air in 3 gallons of water. If needed, agricultural-type spray oil may be added. Apply on a 14 day schedule or as needed throughout the wet season. Apply at 21 day intervals or as needed during dry periods.
	Black Pitting	1.75	1.05	18.9	7	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Cacao	Black Pod	0.75-3.75	2.25	15.75	14	Begin applications at the start of the rainy season and continue while infection conditions persist. Apply 0.75 to 2 pounds at 14 to 21 day intervals or as needed depending on disease severity. For drier areas, make two to four applications using 2.5 to 3.75 pounds per acre according to disease incidence and planting density. Use the higher rates when conditions favor disease development.

13
25**TREE CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Coffee	Coffee Berry Disease (<i>Colletotrichum coffeanum</i>)	2.5-3.5	2.1	12.6	14	Apply first spray after flowering and before onset of long rains and then at 21 to 28 day intervals until picking. Use the higher rates when conditions favor disease development.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	2.5-3.5	2.1	12.6	14	Begin spray program before the onset of long rainy periods and continue throughout the rainy season at 14 to 21 day intervals or as needed. The critical time of spraying to control this disease is just before, during and after flowering(s) especially when coinciding with wet weather. Use the higher rates when rainfall is heavy and disease pressure is high.
	Leaf Rust (<i>Hemileia vastatrix</i>)	0.75-1.75	1.6	12.6	14	Apply before the onset of rain and then at 21 day intervals or as needed while the rains continue. Use the higher rates when rainfall is heavy and disease pressure is high.
	Iron Spot (<i>Cercospora coffeicola</i>), Pink Disease (<i>Corticium salmonicolor</i>)	0.75	0.8	2.4	28	Use concentrate or dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
Filbert	Bacterial Blight	7-10.5	6	24	14	Apply as a postharvest spray. In seasons of heavy rainfall apply a second spray when three-fourths of the leaves have dropped. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil may be added. Use only in the states of Oregon and Washington.
	Eastern Filbert Blight	7-10.5	6	24	14	Apply as a dilute spray in adequate water for thorough coverage. Make applications starting at bud swell to bud break and continue at 2 week intervals or as needed until early May. Thorough coverage is essential. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil may be added. Use only in the states of Oregon and Washington.
Mango*	Anthraxnose	2-4	2.6	18.2	30	Apply monthly after fruit set until harvest. Use the higher rates when rainfall is heavy and disease pressure is high.
Olive	Olive Knot, Peacock Spot	3.5-5.25	3.15	6.3	30	Make first application before winter rains begin. A second application in early spring should be made if disease is severe. Apply the higher rates for heavy disease pressure or when conditions favor disease development.

14
25**TREE CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Peach, Nectarine	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot, (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	3.5-7	6.4	18	7	Make first application before fall rains and a second at late dormant. For peach leaf curl, late dormant application must be made before leaf buds swell. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural-type spray oil may be added.
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	3.5-5.25	1.5	18	5	Full cover spray at pink bud. Use the higher rates when conditions favor disease development.
	Bacterial Spot	0.5	0.4	18	5	Post-bloom application applied at first and second cover sprays. NOTE: Do not spray 3 weeks prior to harvest. Use only recommended rates. Spotting of leaves and defoliation may occur from use in cover sprays.
Pear	Fire Blight	0.5	0.4	16	5	Apply at 5 day intervals or as needed throughout the bloom period. NOTE: Russetting may occur in copper sensitive varieties. Excessive dosages may cause fruit russet on any variety.
	Blossom Blast (<i>Pseudomonas</i>)	5.25-7	6.4	16	n/a	Apply before fall rains and again during dormancy before spring growth starts. Use the higher rates when disease pressure is high or when conditions favor disease development.
Pecan	Kernel Rot, Shuck Rot (<i>Phytophthora cactorum</i>), Zonate Leaf Spot (<i>Cristulariella pyramidalis</i>)	0.75-1.75	1.6	8.4	14	For suppression, apply in sufficient water to ensure complete spray coverage at 2 to 4 week intervals or as needed starting at kernel growth and continue until shucks open. Use the higher rates and shorter spray intervals if frequent rainfall occurs.
	Ball Moss* Spanish Moss*	2.5-3.5	2.1	8.4	365	Apply in 100 gallons of water in the spring when ball moss is actively growing, using 1 ½ gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. The addition of a nonionic surfactant will improve control. A second application may be required after 12 months.
Pistachio	Botryosphaeria Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria alternata</i>), Septoria Leaf Blight	1.75-3.5	2.1	8.4	14	Make initial application at bud swell and repeat on a 14 to 28 day schedule or as needed. If disease conditions are severe, use the higher rates and shorter spray intervals.
Quince*	Fire Blight	0.5	0.4	16	5	Apply at 5 day intervals or as needed throughout the bloom period. Apply in adequate water for thorough coverage.

15
28**TREE CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Walnut	Walnut Blight	3.5-5.25	3.15	25.2	7	Apply first spray at early pre-bloom prior to or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage or as needed when frequent rainfall or extended periods of moisture occur. Thorough coverage of catkins, leaves and nutlets is essential for effective control. Use the higher rates when conditions favor disease development. NOTE: Adequate control may not be obtained when copper tolerant species of <i>Xanthomonas</i> bacteria are present.

* Except California

VEGETABLES

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Bean (Dry, Green)	Brown Spot, Common Blight, Halo Blight	0.5-1.25	0.79	4.74	7	For protective sprays, make first application when plants are 6 inches high; repeat on a 7 to 14 day schedule or as needed depending on environmental conditions. Use the higher rates for more severe disease pressure.
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot	0.75-2	1.31	7.86	10	Begin applications when conditions first favor disease development and repeat at 10 to 14 day intervals or as needed. Use the higher rates when conditions favor disease development.
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	0.75-1.5	0.8	5	7	Begin applications when disease first threatens and repeat at 7 to 14 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease development.
Celery, Celeriac*	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	0.75	0.8	5.3	7	Begin applications as soon as plants are first established in the field, repeating at 5 to 7 day intervals or as needed depending on disease severity and environmental conditions.

16
25**VEGETABLES**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMEN T INTERVAL (DAYS)	COMMENTS
Crucifers (Broccoli, Brussels Sprout, Cabbage, Cauliflower, Collard Greens, Mustard Greens, Turnip Greens)	Black Leaf Spot (<i>Alternaria</i>), Black Rot (<i>Xanthomonas</i>), Downy Mildew	0.5-0.75	0.53	2.65	7	Apply at 7 to 10 day intervals or as needed. Begin application after transplants are set in the field or shortly after emergence of field seeded crops or when conditions favor disease development. Use the higher rates when conditions favor disease development. NOTE: Reddening of older leaves may occur on broccoli and a flecking of wrapper leaves may occur on cabbage.
Cucurbits (Cantaloupe , Cucumber, Honeydew, Pumpkin, Squash, Muskmelon, Watermelon)	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Powdery Mildew, Watermelon Bacterial Fruit Blotch (Suppression)	0.5-1.25	1.05	5.25	5	Begin applications prior to disease development and continue while conditions are favorable for disease development. Repeat sprays at 5 to 7 day intervals or as needed. Use the higher rates when conditions favor disease development. NOTE: Crop injury may occur from application at higher rates and shorter intervals. Discontinue use if injury occurs.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	0.75	0.79	7.9	7	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals or as needed depending on disease severity.
Onion, Garlic	Bacterial Blight, Downy Mildew, Purple Blotch	0.75	1	6	7	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals or as needed depending on disease severity. Can cause phytotoxicity to leaves.
Pea	Powdery Mildew	0.5-1.25	0.79	3.95	7	Begin applications when disease symptoms first appear and repeat at weekly intervals or as needed. Use the higher rates when conditions favor disease development.
Pepper	Anthracnose, Bacterial Spot, Cercospora Leaf Spot	0.75-1.25	0.79	11.85	3	Begin applications when conditions first favor disease development and repeat at 7 to 10 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease development.
Spinach	Anthracnose, Blue Mold, Cercospora Leaf Spot, White Rust	0.75-1.25	0.79	3.95	7	Begin application when disease first appears or when conditions favor disease development. Repeat at 7 to 10 day intervals or as needed. Use the higher rates when conditions favor disease development. NOTE: Flecking may occur on spinach leaves.

17
2.5**VEGETABLES**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Tomato	Anthrachnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.75-1.75	0.53	17.4	3	Begin application when disease first threatens and repeat at 5 to 10 day intervals or as needed depending on disease severity. Use the higher rates when conditions favor disease development
Watercress*	Cercospora Leaf Spot	0.75	0.53	2.12	7	Begin applications when plants are first established in the field, repeating at 7 to 14 day intervals or as needed depending on disease severity. Do not exceed four applications per crop. Apply using ground spray equipment at no less than 50 gallons of spray solution per acre.

* Except California

VINES

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Grape	Black Rot, Downy Mildew, Phomopsis, Powdery Mildew	0.75-1.75	1.6	20	3	Begin applications at bud break with subsequent applications throughout the season depending on disease severity. Use the higher rates when conditions favor disease development. NOTE: Foliage injury may occur on copper sensitive varieties such as Concord, Delaware, Niagara and Rosette. Either test for sensitivity or add 1 to 3 pounds of hydrated lime per pound of BADGE X ₂ .
Hops	Downy Mildew	0.75	0.53	2.65	10	Make crown treatment after pruning, but before training. After training, additional treatments are needed at about 10 day intervals. NOTE: Discontinue use 2 weeks before harvest.
Kiwi	<i>Erwinia herbicola</i> , <i>Pseudomonas fluorescens</i> , <i>Pseudomonas syringae</i>	3.5	2.1	6.3	30	Apply in 200 gallons of water per acre. Make applications on a monthly basis. A maximum of three applications may be made.

MISCELLANEOUS CROPS

18
25

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Atemoya*	Anthrachnose	1.5-2	1.6	12.6	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.
Carambola *	Anthrachnose	2.5-3.5	2.1	10.5	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease.
Chives*	Downy Mildew	0.75	0.53	2.65	7	Begin applications when plants are established in the field. Repeat applications every 7 to 10 days or as needed depending on disease conditions.
Dill*	Phoma Leaf Spot, Rhizoctonia Foliage Blight	0.75-1.25	0.79	3.95	7	Begin applications when plants are first established in the field and repeat at 7 to 10 day intervals or as needed depending upon disease severity and environmental conditions. Use the higher rates when conditions favor disease development.
Ginseng	Alternaria Leaf Blight, Stem Blight	1-1.75	1.05	5.25	7	Use as a tank mix with 2 pounds Rovral® 50W in 100 gallons of water. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Begin BADGE X ₂ -Rovral applications as soon as plants have emerged in spring. Application should be repeated every 7 days or as needed until plants become dormant in fall. Apply fungicides at least 8 hours before rain. Use of a spreader-sticker or sticker is advised. NOTE: Alternaria Leaf and Stem Blight is most severe in humid conditions such as those found in the dense canopies of 2 to 4 year old Ginseng. It is very important that the stems be thoroughly covered with fungicide; therefore, use a spray apparatus which distributes the fungicide throughout the canopy.
Guava	Anthrachnose, Red Algae	1.25-2	1.23	4.92	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease pressure.

MISCELLANEOUS CROPS

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Live Oak*	Ball Moss	2.5-3.5	2.1	8.4	365	Apply in 100 gallons of water in the spring when ball moss is actively growing using 1.5 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months. NOTE: BADGE X ₂ may be injurious to ornamentals grown under Live Oaks or Pecans. This product may be reactive on metal and masonry surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.
Litchi*	Anthracnose	1.25-2	1.23	4.92	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease pressure.
Macadamia	Anthracnose	2.5-4	2.36	9.44	7	Initiate sprays at first sign of flowering and repeat on weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates for severe disease pressure.
	Phytophthora Blight (<i>P. capsici</i>), Raceme Blight (<i>Botrytis cinerea</i>)	1.25-2.5	2.36	9.44	7	Apply during raceme development and bloom periods. Apply in sufficient water for thorough coverage. Use the higher rates when conditions favor disease development.
Mamey Sapote*	Algal Leaf Spot, Anthracnose	2.5-3.5	2.1	8.4	14	Apply when conditions favor disease development. Repeat on 14 to 30 day schedule as disease severity and environmental conditions dictate. Use the higher rates when conditions favor disease development.
Papaya*	Anthracnose	1.75-4.25	2.63	21.2	14	Apply before disease appears. Apply at 14 day intervals or as needed. The addition of an approved spreader is desirable. Use the higher rates when conditions favor disease development.
Parsley*	Bacterial Blight (<i>Pseudomonas sp.</i>)	1.25	1	2	10	Begin applications when plants are first established in the field and repeat at 10 day intervals as needed depending on disease severity and environmental conditions.
Passion Fruit*	Anthracnose	2.5-4	2.36	9.44	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates when conditions favor disease development.

20
25**MISCELLANEOUS CROPS**

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE (LBS Cu ²⁺ /A)	MAX. ANNUAL RATE (LBS Cu ²⁺ /A)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Sugar Apple* (<i>Annona</i>)	Anthracnose	5.25-7.75	3.15	12.6	7	Make initial application just before flowering and repeat on a weekly schedule until just before harvest. Apply in sufficient water for thorough coverage. Use the higher rates when conditions favor disease development.
Sycamore	Anthracnose	0.75-1.25	1.2	3.6	7	Apply as a full cover spray in 100 gallons of water or sufficient volume for thorough coverage. Make first application at bud crack and second application 7 to 10 days later at 10% leaf expansion. Use the higher rates when conditions favor disease development.

* Except California

GREENHOUSE AND SHADEHOUSE CROPS

Notice to User: BADGE X₂ may be used in greenhouses and shadehouses to control disease on crops which appear on this label and specific instructions have been developed for the crops listed. The grower should bear in mind that the sensitivity of crops grown in greenhouses and shadehouses differs greatly from crops grown under field conditions. Neither the manufacturer nor seller has determined whether or not BADGE X₂ can be used safely on all greenhouse and shadehouse grown crops. The user should determine if BADGE X₂ can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, i.e., foliage, fruit, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use.

Apply BADGE X₂ according to specific rates given for those crops in pounds per acre. **One and a half (1.5) level tablespoons of BADGE X₂ per 1000 square feet is equivalent to 2 pounds per acre.** BADGE X₂ should be applied in adequate water for thorough coverage of plant parts. Begin application at first sign of disease and repeat at 7 to 14 day intervals or as needed; use shorter spray intervals during periods when severe disease conditions persist.

CROP	DISEASE	APP. RATE (TBSP PRODUCT/1000 SQ. FT.)	MAX. APP. RATE (TBSP Cu ²⁺ /1000 SQ. FT.)	MAX. ANNUAL RATE (TBSP Cu ²⁺ /1000 SQ. FT.)	MINIMUM RETREATMENT INTERVAL (DAYS)	COMMENTS
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	1.2	1.2	9.5	7	Begin applications when conditions favor disease development. Repeat sprays at 30 day intervals or as needed depending on disease severity.
Cucumber	Angular Leaf Spot, Downy Mildew	0.5-0.8	0.8	4	5	Apply weekly when plants begin to vine. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	0.5	0.6	6	7	Begin applications prior to development of disease symptoms. Repeat sprays at 7 to 10 day intervals or as needed depending on disease severity.
Pepper	Bacterial Spot	0.6	0.6	9	3	Begin applications when conditions first favor disease development and repeat at 3 to 10 day intervals or as needed depending on disease severity.

21
25

CROP	DISEASE	APP. RATE (TBSP PRODUCT/1000 SQ. FT.)	MAX. APP. RATE (TBSP Cu ²⁺ /1000 SQ. FT.)	MAX. ANNUAL RATE (TBSP Cu ²⁺ /1000 SQ. FT.)	MINIMUM RETREATMEN T INTERVAL (DAYS)	COMMENTS
Tomato	Anthrachnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	0.4	0.4	13	3	Begin applications when disease first threatens and repeat at 3 to 10 day intervals or as needed depending on disease severity.

CONIFERS

For use on conifers, including Douglas Fir, Fir*, Juniper, Leyland Cypress*, Pine* and Spruce*, in Christmas tree plantings, forest stands and silviculture nurseries. For control of foliar diseases, apply BADGE X₂ as a thorough cover spray at rates ranging from 0.75 to 1.75 pounds per acre. Begin applications in the spring at the initiation of new growth and repeat at 2 to 4 week intervals or as needed. Use the higher rates when disease pressure is severe or when environmental conditions favor disease development. **There is a maximum application rate of 2.0 lbs Cu²⁺/A with a maximum annual rate of 20 lbs Cu²⁺/A with a minimum retreatment interval of 7 days.**

BADGE X₂ is recommended for use on the listed conifers for control of the following diseases:

CROP	LATIN NAME	DISEASE
Douglas Fir	<i>Pseudotsuga menziesii</i>	Rhabdocline Needlecast
Fir*	<i>Abies</i> spp.	Needlecasts
Juniper	<i>Juniperus</i> spp.	Anthrachnose, Phomopsis Twig Dieback*
Leyland Cypress*	<i>X Cupressocyparis leylandii</i>	Cercospora Needle Blight
Pine*	<i>Pinus</i> spp.	Needlecasts
Spruce*	<i>Picea</i> spp.	Needlecasts

Lichens*: To control lichens on any of the conifers above, apply 3.5 pounds of BADGE X₂ per acre as a dormant application before new growth emerges in the spring. The addition of a non-ionic surfactant (NIS) will improve control. A second application may be required after 12 months. **NOTE:** Do not buffer or combine with emulsifiable concentrate insecticides.

* Except California

ORNAMENTALS

Use BADGE X₂ for control of bacterial and fungal diseases of foliage, flowers and stems on ornamentals in greenhouses, shadehouses, outdoor nurseries, and outdoor landscape plantings:

For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 1.5 to 5 pounds per acre of BADGE X₂. When new growth is present, apply as a thorough cover spray at rates ranging from 1.5 to 2 pounds per acre of BADGE X₂. **One and a half (1.5) level tablespoons of BADGE X₂ per 1000 square feet is equivalent to 2 pounds per acre.** Begin application at first sign of disease and repeat at 7 to 14 day intervals or as needed; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist.

Unless otherwise noted, the maximum single application rate is 2 pounds of Cu²⁺ per acre and the maximum annual rate is 20 pounds of Cu²⁺ per acre. The minimum retreatment interval is 7 days.

BADGE X₂ may be used alone or in combination with other fungicides registered for use on ornamentals as a maintenance spray. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Notice to User: Plant sensitivities to BADGE X₂ have been found to be acceptable for the specific genera and species listed on this label under the

conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants and the wide range of growing conditions, it is impossible to test every one for sensitivity to BADGE X₂. Neither the manufacturer nor the seller has determined whether or not BADGE X₂ can be safely used on ornamental or nursery plants not listed on this label. The user should determine if BADGE X₂ can be used safely prior to commercial use. In a small area, apply the recommended rates to the plants in question, (bedding plants, foliage, etc.), and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use. **NOTE:** This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

CROP	SCIENTIFIC NAME	DISEASE
Aglaonema*	<i>Aglaonema</i> spp.	Bacterial Leaf Spot

Althea (Rose of Sharon)	<i>Hibiscus syriacus</i>	Bacterial Leaf Spot
Andromeda, Japanese*	<i>Pieris japonica</i>	Leaf Spots, Twig Blight
Aralia	<i>Dizygotheca elegantissima</i>	Alternaria, Cercospora Leaf Spot, Xanthomonas Leaf Spot
Arborvitae	<i>Thuja</i> spp.	Alternaria Twig Blight, Cercospora Leaf Blight
Aster*	<i>Aster</i> spp.	Downy Mildew, Leaf Spots
Azalea 1/	<i>Rhododendron</i> spp.	Botrytis Blight, Cercospora Leaf Spot, Phytophthora Dieback, Powdery Mildew
Beech*	<i>Fagus</i> spp.	Leaf Spots
Begonia	<i>Begonia semperflorens</i>	Bacterial Leaf Spot (<i>Erwinia</i> spp., <i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp.)
Bougainvillea	<i>Bougainvillea spectabilis</i>	Anthracnose, Bacterial Leaf Spot
Boxwood*	<i>Buxus</i> spp.	Leaf Spots
Camellia	<i>Camellia japonica</i> , <i>C. sasanqua</i>	Anthracnose, Bacterial Leaf Spot
Camphor Tree	<i>Cinnamomum camphora</i>	<i>Pseudomonas</i> Leaf Spot
Canna	<i>Canna</i> spp.	<i>Pseudomonas</i> Leaf Spot
Carnation 1/	<i>Dianthus</i> spp.	Alternaria Blight, Botrytis Blight, <i>Pseudomonas</i> Leaf Spot
Cedar*	<i>Cedrus</i> spp.	Tip Blight
Cherry, Nanking*	<i>Prunus tomentosa</i>	Bacterial Leaf Spot
Chinese Tallow Tree	<i>Sapium sebiferum</i>	Bacterial Leaf Spot (<i>Pseudomonas</i> spp., <i>Xanthomonas</i> spp.)
Chrysanthemum 1/	<i>Chrysanthemum morifolium</i>	Botrytis Blight, <i>Pseudomonas</i> Leaf Spot, Septoria Leaf Spot
Cotoneaster	<i>Cotoneaster</i> spp.	Botrytis Blight
Crabapple*	<i>Malus</i> spp.	Fire Blight
Cypress*	<i>Cupressus</i> spp.	Twig Blight
Dahlia	<i>Dahlia pinnata</i>	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Delphinium*	<i>Delphinium</i> spp.	Leaf Spots
Dianthus	<i>Dianthus</i> spp.	Bacterial Soft Rot, Bacterial Spot
Dogwood, Flowering	<i>Cornus florida</i>	Anthracnose
Dogwood, Kousa*	<i>Cornus kousa</i>	Fungal Leaf Spot
Douglas Fir	<i>Pseudotsuga menziesii</i>	Rhabdocline Needlecast
Dracaena*	<i>Dracaena marginata</i>	Bacterial Leaf Spot
Dumb Cane*	<i>Dieffenbachia</i> spp.	Bacterial Leaf Spot
Dusty Miller	<i>Senecio cineraria</i>	Bacterial Leaf Spot (<i>Pseudomonas cichorii</i>)
Echinacea	<i>Echinacea</i> spp.	Bacterial Leaf Spot (<i>Pseudomonas cichorii</i>)
Elm, Chinese	<i>Ulmus parvifolia</i>	Xanthomonas Leaf Spot
Euonymus	<i>Euonymus</i> spp.	Anthracnose, Botrytis Blight
Fern, Boston*	<i>Nephrolepis exaltata</i>	Bacterial Leaf Spot
Fern, Holly	<i>Cytomium falcatum</i>	<i>Pseudomonas</i> Leaf Spot
Fig, Weeping*	<i>Ficus benjamina</i>	Bacterial Leaf Spot
Filbert (Ornamental)*	<i>Corylus</i> spp.	Filbert Blight
Fir*	<i>Abies</i> spp.	Needlecasts
Gardenia	<i>Gardenia jasminoides</i>	Alternaria Leaf Spot, Botrytis Bud Rot, Cercospora Leaf Spot
Geranium	<i>Pelargonium</i> spp.	Alternaria Leaf Spot, Botrytis Gray Mold, Cercospora Leaf Spot
Gladiola	<i>Gladiolus</i> spp.	Alternaria Leaf Spot, Anthracnose, Bacterial Leaf Blight, Botrytis Gray Mold
Golden Rain Tree	<i>Koelreuteria paniculata</i>	Bacterial Leaf Spot
Grape Ivy*	<i>Cissus</i> spp.	Bacterial Leaf Spot
Hawthorn*	<i>Crataegus</i> spp.	Fire Blight
Hibiscus 4/	<i>Hibiscus</i> spp.	Bacterial Leaf Spot
Holly*	<i>Ilex</i> spp.	Bacterial Blight, Leaf Spots
Honeylocust*	<i>Gleditsia triacanthos</i>	Bacterial Leaf Spot
Honeysuckle, Tatarian*	<i>Lonicera tatarica</i>	Bacterial Leaf Spot
Impatiens	<i>Impatiens sallerana</i>	Bacterial Leaf Spot
Indian Hawthorn 5/	<i>Raphiolepis indica</i>	Anthracnose, Entomosporium Leaf Spot
Iris 6/*	<i>Iris</i> spp.	Bacterial Leaf Spot
Ivy (English, Algerian) 1/	<i>Hedera helix</i> , <i>H. canariensis</i>	Xanthomonas Leaf Spot
Ixora	<i>Ixora coccinea</i>	Xanthomonas Leaf Spot
Juniper	<i>Juniperus</i> spp.	Anthracnose, Phomopsis Twig Dieback*
Lantana	<i>Lantana camera</i>	Bacterial Leaf Spot
Leyland Cypress*	<i>X Cupressocyparis leylandii</i>	Cercospora Needle Blight
Lilac	<i>Syringa</i> spp.	Cercospora Leaf Spot, <i>Pseudomonas</i> Blight*
Lily, Easter 2/	<i>Lilium longiflorum</i>	Botrytis Blight
Linden*	<i>Tilia</i> spp.	Anthracnose, Leaf Blight
Loblolly Bay	<i>Gordonia lasianthus</i>	Anthracnose

Loquat	<i>Eriobotrya japonica</i>	<i>Colletotrichum</i> spp., <i>Entomosporium maculata</i>
Magnolia (Southern)	<i>Magnolia grandiflora</i>	Algal Leaf Spot, Anthracnose, Bacterial Leaf Spot
Magnolia (Sweet Bay)	<i>Magnolia virginiana</i>	Anthracnose
Magnolia (Oriental)	<i>Magnolia soulangiana</i>	Bacterial Leaf Spot
Mandevilla	<i>Mandevilla</i> spp.	Anthracnose
Maple*	<i>Acer</i> spp.	<i>Pseudomonas</i> Leaf Blight
Marigold	<i>Tagetes</i> spp.	<i>Alternaria</i> Leaf Spot, <i>Botrytis</i> Leaf Rot, <i>Cercospora</i> Leaf Spot, Flower Rot
Mountain-Ash*	<i>Sorbus</i> spp.	Fire Blight
Mulberry, Contorted*	<i>Morus bombycis</i>	Bacterial Leaf Spot
Mulberry, Weeping	<i>Morus alba</i>	Bacterial Leaf Spot
Narcissus*	<i>Narcissus</i> spp.	Leaf Blight
Nephtytis*	<i>Syngonium podophyllum</i>	Bacterial Leaf Spot
Oak*	<i>Quercus</i> spp.	Leaf Spots
Oak, Laurel	<i>Quercus laurifolia</i>	Algal Leaf Spot (<i>Cephaleuros virescens</i>)
Oleander	<i>Nerium oleander</i>	Bacterial Leaf Spot, Fungal Leaf Spot
Oregon Grapeholly*	<i>Mahonia aquifolium</i>	Leaf Spots
Pachysandra	<i>Pachysandra procumbens</i>	<i>Volutella</i> Leaf Blight
Palm, Date	<i>Phoenix canaries</i>	<i>Pestalotia</i> Leaf Spot
Palm, European Fan	<i>Chamaerops humilis</i>	<i>Pestalotia</i> Leaf Spot
Palm, Parlor*	<i>Chamaedorea elegans</i>	Bacterial Leaf Spot
Palm, Queen	<i>Arecastrum romanzoffianum</i>	Exosporium Leaf Spot, <i>Phytophthora</i> Bud Rot
Palm, Washingtonia	<i>Washingtonia robusta</i>	<i>Pestalotia</i> Leaf Spot
Peach (Flowering) 3/*	<i>Prunus</i> spp.	Bacterial Blast, Brown Rot, Fire Blight
Pear (Flowering)	<i>Pyrus calleryana</i>	Fire Blight, Leaf Spot
Pentas (Egyptian Star)	<i>Pentas</i> spp.	Bacterial Leaf Spot (<i>Pseudomonas</i> spp.*, <i>Xanthomonas</i> spp.)
Peony	<i>Paeonia</i> spp.	<i>Botrytis</i> Blight
Periwinkle	<i>Catharanthus roseus</i> , <i>Vinca</i> spp.	<i>Phomopsis</i> Stem Blight
Philodendron	<i>Philodendron selloum</i>	Bacterial Leaf Spot
Phlox	<i>Phlox</i> spp.	<i>Alternaria</i> Leaf Spot
Photinia (Red Tip)	<i>Photinia x fraseri</i> , <i>P. glabra</i>	Anthracnose, <i>Entomosporium</i> Leaf Spot
Pine*	<i>Pinus</i> spp.	Needlecasts
Pistachio	<i>Pistacia chinensis</i>	Anthracnose
Plantain Lily 6/	<i>Hosta</i> spp.	Bacterial Leaf Spot
Plum (Flowering) 3/*	<i>Prunus</i> spp.	Bacterial Blast, Bacterial Leaf Spot, Brown Rot, Fire Blight
Pothos*	<i>Scindapsus</i> spp.	Bacterial Leaf Spot
Powder Puff Plant	<i>Calliandra</i> spp.	Bacterial Leaf Spot
Pyracantha	<i>Pyracantha</i> spp.	Fire Blight, Scab
Rhododendron	<i>Rhododendron</i> spp.	<i>Alternaria</i> Flower Spot
Rose 1/	<i>Rosa</i> spp.	Black Spot, Powdery Mildew
Snapdragon	<i>Antirrhinum majus</i>	Anthracnose, Dieback, Downy Mildew
Spathe Flower*	<i>Spathiphyllum</i> spp.	Bacterial Leaf Spot
Spirea*	<i>Spiraea</i> spp.	Fire Blight
Spruce*	<i>Picea</i> spp.	Needlecasts
Sycamore	<i>Platanus</i> spp.	Anthracnose, Leaf Spots*
Tulip	<i>Tulipa</i> spp.	Anthracnose, <i>Botrytis</i> Blight
Umbrella Tree*	<i>Schefflera</i> spp.	Bacterial Leaf Spot
Verbena	<i>Verbena</i> spp.	<i>Xanthomonas</i> Leaf Spot
Viburnum	<i>Viburnum odoratissimum</i> , <i>V. suspensum</i> , <i>V. plicatum</i>	Anthracnose
Viola (Pansy, Violet)	<i>Viola</i> spp.	Downy Mildew
Willow	<i>Salix</i> spp.	Anthracnose
Yew*	<i>Taxus</i> spp.	Needle Blight
Yucca (Adam's Needle)	<i>Yucca</i> spp.	<i>Cercospora</i> Leaf Spot, <i>Septoria</i> Leaf Spot
Zinnia*	<i>Zinnia</i> spp.	Leaf Spots

* Except California

- 1/ Can cause discoloration of foliage and/or blooms on some varieties. To prevent residues on commercial plants, do not spray immediately before selling season.
- 2/ Apply at 4.5 to 7.5 pounds per acre. The maximum single application rate is 2.5 pounds of Cu²⁺ per acre. The maximum amount of metallic copper which may be applied in a 12 month period is 75 pounds of Cu²⁺ per acre. Do not apply any additional copper pesticide to this land for 36 months.
- 3/ Apply dormant through bloom only.
- 4/ Hibiscus - Do not apply to plants in flower.
- 5/ For Indian Hawthorne use 3 to 6 pounds per acre.

24
28

6/ Some cultivars may be sensitive to BADGE X₂.

NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of BADGE X₂, apply the recommended rate to a few plants and observe after 7 to 10 days for symptoms of phytotoxicity.

Control of Ball Moss*, Spanish Moss* and Lichens* on Ornamentals and Shade Trees: Apply BADGE X₂ in early spring when trees are dormant. Apply 9 to 12 pounds of BADGE X₂ in 100 gallons of water, using 1 ½ gallons of spray per foot of tree height. Be sure to thoroughly wet ball moss tufts, Spanish moss or lichens. The addition of a non-ionic surfactant will improve control. A second application may be required after 12 months.

NOTE: BADGE X₂ may be injurious to some ornamental plants growing beneath the trees. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, etc.

Cold Storage Protection for Dormant Rootstock*: To protect bare-root nursery trees from Phytophthora Crown Rot and Botrytis, use 4 to 6 pounds of BADGE X₂ per 100 gallons of water. Apply as a dip or spray to the roots and lower stems of dormant rootstock prior to placing in cold storage. Do not apply to rootstock less than 2 years old.

*Except California

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed through storage and disposal.

Pesticide Storage:	Store under well-vented, cool and dry storage conditions. Do not store under moist conditions.
Pesticide Disposal:	Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
Container Type:	This is a nonrefillable, non-rigid container (bag). Do not reuse or refill this container.
Container Disposal:	Empty the package completely. Then dispose of the empty container according to state and local regulations. Place in trash or offer for recycling if available or return it to the Seller, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire label before using this product, including this Limitation of Warranty and Liability.

If the terms are not acceptable, return the product at once unopened for a refund of the purchase price.

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks described below, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ISAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Buyers and Users of this product must be aware that there are inherent unintended risks associated to the use of this product, independent from the control of Isagro. These risks include, but are not limited to, weather conditions, soil factors, moisture conditions, diseases, irrigation practices, condition of the crop at the time of application, materials which are present in the tank mix with this product or prior to the application of it, cultural practices or the manner of use or application, all risks which are impossible to eliminate. The Buyers and Users should be aware that these factors may cause: ineffectiveness of the product, reduction of harvested yield of the crop (entirely or partially), crop injury or injury to non-target crops or plants or to rotational crops caused by carryover in the soil, resistance of the target diseases to this product. Therefore additional care, treatment and expense are required to take the crop to harvest.

If the Buyer does not agree with the acceptance of these risks, then THE PRODUCT SHOULD NOT BE APPLIED. To the extent consistent with applicable law, by applying this product the Buyer acknowledges and accepts these inherent

25
25

unintended risks and AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

To the extent consistent with applicable law, in no event shall ISAGRO or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise). To the extent consistent with applicable law, the exclusive remedy of the User or Buyer and the exclusive Liability of Isagro or Seller shall be the return of the purchase price of the product, or at the election of Isagro or Seller, the replacement of the product.

To the extent consistent with applicable law, this Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

Isagro or its Seller must have prompt notice (within 7 days of observation) of any claim so that an immediate inspection of Buyer's or User's growing crops can be made. To the extent consistent with applicable law, if Buyer and User do not notify Isagro or Seller of any claims, in proper time, it shall be barred from obtaining any remedy.

To the extent consistent with applicable law, by applying this product the Buyers and Users are deemed to have accepted the terms of this Limitation of Warranty and Liability, which may not be modified by any verbal or written agreement.

Alliete is a registered trademark of Bayer CropScience.
Badge is a registered trademark of Isagro USA
Curtec is a registered trademark of Curtec Corporation.
Kentan is a registered trademark of Isagro S.p.A.
Rovral is a registered trademark of Bayer CropScience.
Tre-Hold is a registered trademark of Amvac Chemical Corporation

ESL

REV

15jan09