



80289-18

5/30/2013

1115

 U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Chemical Safety and Pollution Prevention Office of Pesticide Programs Registration Division (7504P) 1200 Pennsylvania Ave., N.W. Washington, DC 20460	EPA Reg. Number: 80289-18	Date of Issuance: MAY 30 2013
	Term of Issuance: Unconditional	
	Name of Pesticide Product: Eminent® ESP	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Registration Review Under FIFRA, as amended		
Name and Address of Registrant (include ZIP Code): Isagro S.p.A. D/B/A Isagro USA, Inc. 430 Davis Drive, Suite 240 Morrisville, NC 27560		
Mailed to: Mel Graben Regulatory & Technical Manager		
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is registered in accordance with FIFRA section 3(c)(5) provided that you:</p> <ol style="list-style-type: none">1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration/registration review of your product under FIFRA section 4.		
Page 1 of 2		
Signature of Approving Official:  Hope Johnson, Acting Product Manager (21) Fungicide Branch/Registration Division/OPP/OCSPP (7504P)	Date: MAY 30 2013	

2115

MAY 30 2013


2. Make the following changes to the label:
 - a. Change the product registration number to "EPA Reg. No. 80289-18"
 - b. Add Net Contents information
 - c. Add EPA Establishment Number
3. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these requirements are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

The basic Confidential Statement of Formula (CSF) dated May 17, 2013 for the product referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act is acceptable. This basic CSF will be added to the file for this product.


The Agency acknowledges your request for the alternate brand name "Eminent VP".

A copy of the label stamped "Accepted" is enclosed for your records.



Hope Johnson, Acting Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure:
Label stamped "Accepted"



GROUP	3	FUNGICIDE
-------	---	-----------

EMINENT® ESP

(Alternate brand name: Eminent VP)

Active Ingredient:

Tetraconazole*	11.6%
Other Ingredients	88.4%
Total	100.0%

*1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)propyl]1H-1,2,4-triazole

Contains 1 lb active ingredient (tetraconazole) per gallon

EMINENT is a registered trademark of Isagro S.p.A.

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

[If you do not understand this label, find someone to explain it to you in detail.]

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have affected person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
For Chemical Emergency Spill Leak Fire Exposure or Accident Call CHEMTREC Day or Night Domestic, North America 800-424-9300 International 703-527-3883 (collect calls accepted)	

[See inside booklet for additional precautionary statements and use directions]

EPA Registration No. 80289-pending

EPA Establishment No. _____

Manufactured by Isagro SpA for:

Batch code will be placed on the container



Isagro USA, Inc.
430 Davis Drive, Suite 240
Morrisville NC 27560

NET CONTENTS: _____ various

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

CAUTION / PRECAUCION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Some materials that are chemical resistant to this product are barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, and viton ≥14 mils. If you want more options, follow the instructions for category C on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl rubber, Viton, Selection Category C).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.


ENVIRONMENTAL HAZARDS

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise caution when making applications of EMINENT ESP and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all activities with the exception of 7 days for table grape activities of girdling, cane tying and cane turning. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl rubber, Viton, Selection Category C).
- Shoes plus socks

PRODUCT INFORMATION

EMINENT ESP is formulated as a one pound active ingredient per gallon micro emulsion (ME). The active ingredient in EMINENT ESP is tetraconazole, a triazole fungicide (Group 3) that works by inhibiting demethylation and other processes in sterol biosynthesis. Tetraconazole is a systemic, protectant and curative fungicide and is absorbed quickly into the plant tissue. Optimal disease control is achieved when EMINENT ESP is applied in a regularly scheduled spray program.

Pest Management Strategies

1. IPM: Integrate EMINENT ESP into a comprehensive disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, pest control adviser and/or Isagro representative for additional IPM strategies established for your area. Use EMINENT ESP in Agricultural Extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
2. Tank mixtures: EMINENT ESP may be used in tank mixtures with fungicides having a different mode of action which are registered/permitted for the same use and are effective against the target pathogen. Tank-mixing EMINENT ESP with other Group 3 fungicides is not recommended. Follow the more restrictive labeling for any tank mix partner. Do not mix with any product which contains a prohibition on tank mixing.

RAINFASTNESS

EMINENT ESP is rainfast 2 hours after application. **Do not** apply if rain is expected within 2 hours of application or disease control may be reduced.

COMPATIBILITY OF MIXTURES

EMINENT ESP is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, micronutrients and adjuvants. To ensure better results, consult spray compatibility charts available from State Cooperative Extension Service Specialists when comparing tank mixtures and conduct a spray tank compatibility test before mixing this product with other products. To determine the physical compatibility of EMINENT ESP, conduct a simple jar test as follows:

1. Add 1 pt. of water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 ml of EMINENT ESP to the quart jar; gently mix until product goes into suspension.

3. Add the proportionate amount of the mix product(s), with agitation. Then dry formulations, then flowables, then emulsifiable concentrates, and then adjuvants.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An ideal tank-mix combination will be uniform and free of suspended particles. The following conditions indicate potential problems with the mixture and it should not be used:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin.
6. For best results, use combinations on a small number of plants before treating large areas.

SPRAYER PREPARATION

Before applying EMINENT ESP start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply EMINENT ESP. If two or more products were tank mixed prior to EMINENT ESP application, follow the most restrictive cleanup procedure.

Frequently check all application equipment (pressure, nozzles) to ensure complete coverage of the target crop and accurate rate of pesticide application.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the EMINENT ESP to the spray tank. Agitation should create a rippling or rolling action on the water surface.
3. If tank-mixing EMINENT ESP with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions.
4. Adjuvants should be added to the spray solution as required.
5. Fill spray tank to desired level with water. Continue agitation until all spray solution has been applied.
6. Mix only the amount of spray solution that can be applied the day of mixing. Apply EMINENT ESP within 24 hours of mixing.

SPRAYER CLEANUP

Clean spray equipment each day following EMINENT ESP application. After EMINENT ESP is applied; use the following steps to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

SPRAY DRIFT MANAGEMENT

The interaction of many factors including equipment and weather during application determines the potential for spray drift. Applicators are responsible for considering all of these factors when making application decisions. Where states have more stringent regulations, they must be observed.

When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION".

GROUND APPLICATION

Apply product in sufficient water carrier to obtain adequate coverage of all crop surfaces that are intended to be protected from disease. Increase spray volume as crop growth increases. Spray coverage is affected by nozzle type and spacing, sprayer pressure, gallonage per acre (gpa), applicator speed, and other factors.

Airblast (Air Assist) Specific Recommendations: Airblast sprayers deliver the spray mixture into the canopy through a laterally directed airstream. The following drift management practices should be followed when

using an Airblast sprayer:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy
- Block off upward pointed nozzles when there is no overhanging canopy
- Use only enough air volume to penetrate the canopy and provide good coverage
- Do not allow the spray to go beyond the edge of the cultivated area (i.e. turn off sprayer when turning at end rows)
- Only spray inward, toward the field, orchard or vineyard, for applications to the outside rows.

AERIAL APPLICATION

Apply in a minimum of 10 gallons of water per acre. Do not apply under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.

Aerial Spray Drift Reduction Section

Spray Droplet Size: The best drift management strategy is to apply the largest droplets that provide sufficient plant coverage and pest control. Larger droplets reduce drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Spray Droplet Size Control:

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets.
- **Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than any other orientations and is the recommended practice.
- **Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles orientated straight back produce the largest droplets and the lowest drift.

Boom Length: Reducing the effective overall boom length to 70% of the wingspan of fixed-wing aircraft or 80% of a helicopter rotor width may further reduce drift without reducing swath width.

Application Height: Applications must not be made at a height greater than 10 feet above the top of the largest plants.

Application Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, the applicator must compensate for this displacement by adjusting the path of the aircraft or boom on-off. Increase swath adjustment distances, with increasing drift potential (higher wind, height, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Avoid application below 2 mph due to variable wind direction and high inversion potential. Application is not allowed when wind speeds exceed 10 mph due to risk of direct drift to nontarget sensitive crops or locations. **Note:** Wind patterns can be affected by local terrain. All applicators must be familiar with local wind patterns and how they affect spray drift. **Note:** Follow State and local regulations with regard to minimum and maximum wind speeds during aerial application, as they may be more restrictive. Applicators must be familiar with and comply with State and local regulations.

Temperature and Humidity: Applications made during periods of low relative humidity require set-up of equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is typically greatest when conditions are both hot and dry.

Surface Temperature Inversion: Do not apply this product during a local, low level temperature inversion because drift potential is high. Small droplets can be transported in unpredictable directions due to the light and variable winds common during temperature inversions. Temperature inversions are typically characterized by temperatures that increase with altitude and they are common on nights with limited cloud

cover and light to no wind. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

CHEMIGATION INSTRUCTIONS:

- Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other irrigation experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Prevent the Movement of Eminent ESP into the Soil

- Minimize pesticide contact with the soil surface by chemigating above the crop canopy.
- Stop chemigation when pesticide mixture is observed running off crop surfaces or after 0.25 inches of water has been applied, whichever occurs first.
- Allow for sufficient time after chemigation for crop surfaces to dry prior to expected rainfall or to irrigation applied above the crop canopy.

Requirements for Chemigation Systems Connected to Public Water Systems

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventor (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favor drift beyond the area intended for treatment.

- When mixing, fill nurse tank half full with water. Add EMINENT ESP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.
- EMINENT ESP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

Sprinkler Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- When mixing, fill nurse tank half full with water. Add EMINENT ESP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.
- EMINENT ESP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

ROTATIONAL CROP RESTRICTIONS

Use the time intervals listed below to determine the minimum required time interval between last EMINENT ESP application and new crop planting.

Crop	Replant Interval
Soybean, corn, grape, strawberry, sugarbeet, peanut, pecan, gooseberry, kiwifruit (hardy), maypop, schisandra berry, strawberry, bearberry, bilberry, blueberry (lowbush), cloudberry, lingonberry, muntries, and partridgeberry	0 days
Small grains after sugarbeet application	45 days
All other crops – after application to sugarbeet, peanut and pecan	120 days

RESTRICTIONS AND LIMITATIONS

- **Do not** make more than the specified number of applications of EMINENT ESP to each labeled crop per year.
- A restricted entry interval (REI) of 12 hours is to be followed for all activities. For early entry into treated areas refer to PPE requirements under the AGRICULTURAL USE REQUIREMENTS section.

Crop Specific Application Instructions

SUGARBEET

Crop	Target Diseases	Product Use Rate per Application (fl oz/A)	Application Instructions
sugarbeet	Cercospera leaf spot (<i>C. beticola</i>)	13 (0.102 lb ai/A)	Apply preventively when conditions are favorable for disease development or based on a forecasting system. For powdery mildew, apply at the first sign of disease.
	powdery mildew (<i>Erysiphe polygoni</i>) Ramularia leaf spot (<i>R. beticola</i>)	8 to 13 (0.063 to 0.102 lb ai/A)	After EMINENT ESP application, alternate to a non-triazole (non-Group 3) fungicide which is registered for use on sugarbeet for the target disease(s). Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.
Specific Use Restrictions <ul style="list-style-type: none">Do not apply more than 13 fluid ounces (0.102 lb ai) per season of EMINENT ESP or tetraconazole containing products.Do not apply more than 1 application of EMINENT ESP per season.Do not apply within 14 days of harvest (PHI = 14 days).			

PEANUT

Crop	Target Diseases	Product Use Rate per Application (fl oz/A)	Application Instructions
peanut	<p>early leaf spot (<i>Cercospora arachidicola</i>)</p> <p>late leafspot (<i>Cercosporidium personatum</i>)</p> <p>web blotch (<i>Phoma arachidicola</i>)</p> <p>rust (<i>Puccinia arachidis</i>)</p>	6 to 13 (0.047 to 0.102 lb ai/A)	<p>Begin applications prior to onset of disease when conditions are favorable for disease development, generally around 30 to 40 days after planting.</p> <p>Reapply EMINENT ESP using a 14 day interval.</p> <p>EMINENT ESP may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.</p> <p>Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results.</p> <p>Application may be made by ground, air, or chemigation. Apply in 0.1 to 0.25 inches/A of water for chemigation applications. Chemigation application using excessive water could lead to reduced efficacy.</p>
<p>Specific Use Restrictions</p> <ul style="list-style-type: none"> Do not apply more than 52 fluid ounces (0.812 lb ai) per season of EMINENT ESP or tetraconazole containing products for all diseases. Do not apply more than 4 applications of EMINENT ESP per season. There must be a retreatment interval of at least 14 days between multiple applications of EMINENT ESP. Do not apply within 14 days of digging (PHI = 14 days). Do not feed peanut hay to livestock. 			

PECAN

Crop	Target Diseases	Product Use Rate per Application (fl oz/A)	Application Instructions
pecan	<p>powdery mildew (<i>Microsphaera penicillata</i>)</p> <p>scab (<i>Cladosporium caryigenum</i>)</p> <p>brown leaf spot (<i>Cercospora fusca</i>)</p> <p>downy spot (<i>Mycosphaerella carvigena</i>)</p> <p>leaf blotch (<i>Mycosphaerella dendroides</i>)</p> <p>vein spot/leafspot (<i>Gnomonia nerviseda</i>)</p> <p>liver spot (<i>Gnomonia caryae</i> pv <i>pecanae</i>)</p> <p>zonate leaf spot (<i>Cristulariella moricola</i>)</p>	6 to 16 (0.047 to 0.125 lb ai/A)	<p>Begin applications at bud break and continue through pollination using a 14 day interval. After pollination, during cover sprays use a 14 to 21 day interval.</p> <p>Use the highest labeled rates when disease pressure is high. Make no more than 2 sequential applications of a Group 3 fungicide before alternating to a fungicide with a different mode of action.</p> <p>Lower labeled rates may be used when tank mixing with other fungicides labeled for control of target disease(s).</p> <p>Sufficient water volume must be used to ensure thorough coverage for best disease control. Ground application is recommended for best results. Minimum recommended spray volumes are 20 gallons per acre by ground and 10 gallons per acre by air.</p> <p>Application may be made by ground or air.</p>

Specific Use Restrictions

- Do not apply more than 64 fluid ounces (0.5 lb ai) per season of EMINENT ESP or tetraconazole containing products for all diseases.
- Do not apply more than 4 applications of EMINENT ESP per season.
- There must be a retreatment interval of at least 14 days between multiple applications of EMINENT ESP.
- Do not apply within 30 days of harvest or after shuck split (PHI = 30 days).
- Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed through storage and disposal.

Storage: Store under well-vented, cool and dry storage conditions. Do not store under moist conditions.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: **For up to 5 gallon container:** Nonrefillable container: Do not reuse or refill this container. Empty the package completely and triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

For up to 50 gallon container: Nonrefillable container: Do not reuse or refill this container. Empty the package completely and triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents from this container into application equipment or mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

FOR BULK AND MINI-BULK CONTAINERS

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use, and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call Chem. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

THIS CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire label before using this product, including this Limitation of Warranty and Liability.

If the terms are not acceptable, return the product at once unopened for a refund of the purchase price.

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks described below, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ISAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Buyers and Users of this product must be aware that there are inherent unintended risks associated to the use of this product, independent from the control of Isagro. These risks include, but are not limited to, weather conditions, soil factors, moisture conditions, diseases, irrigation practices, condition of the crop at the time of application, materials which are present in the tank mix with this product or prior to the application of it, cultural practices or the manner of use or application, all risks which are impossible to eliminate. The Buyers and Users should be aware that these factors may cause: ineffectiveness of the product, reduction of harvested yield of the crop (entirely or partially), crop injury or injury to non-target crops or plants or to rotational crops caused by carryover in the soil, resistance of the target weeds to this product. Therefore additional care, treatment and expense are required to take the crop to harvest.

If the Buyer does not agree with the acceptance of these risks, then **THE PRODUCT SHOULD NOT BE APPLIED.** To the extent consistent with applicable law, by applying this product the Buyer acknowledges and accepts these inherent unintended risks and **AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.**

To the extent consistent with applicable law, ISAGRO or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product (including claims based in contract, negligence, strict liability, other tort or otherwise). To the extent consistent with applicable law, the exclusive remedy of the User or Buyer and the exclusive Liability of Isagro or Seller shall be the return of the purchase price of the product, or at the election of Isagro or Seller, the replacement of the product.

To the extent consistent with applicable law, this Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

Isagro or its Seller must have prompt notice of any claim so that an immediate inspection of Buyer's or User's can be made. To the extent consistent with applicable law, if Buyer and User do not notify Isagro or Seller of any claims, in proper time, it shall be barred from obtaining any remedy.

To the extent consistent with applicable law, Buyers and Users are deemed to have accepted the terms of this Limitation of Warranty and Liability, which may not be modified by any verbal or written agreement.

Made in U.S.A.

Clean May 23, 2013

[illegible]