

80967-7

9/30/2010

10441



U.S. Environmental Protection Agency
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:
80967-7
Date of Issuance:
SEP 30 2010

Term of Issuance: **Conditional**

Name of Pesticide Product:
Gly Atra N Go Herbicide

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

MEY Corporation
121 S Estes Drive, Suite 101
Chapel Hill, NC 27514

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you agree in writing to:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Revise the EPA Registration Number from 80967-XXX to 80967-7 on the label.
3. Add an appropriate EPA Establishment Number to the label.
4. Add the Restricted Use Pesticide statement and box to the top of page 1
5. Revise the Co-pack Product name from "GLY ATRA" to either something that includes a reference to the dicamba active ingredient as well, or to something completely different. Submit the proposed new name as a Notification for approval within 30 days of receipt of this Notice.

Signature of Approving Official:

Jim Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:
SEP 30 2010


6. On page 3, revise the Precautionary Statements to the following: "Causes irreversible eye damage. Harmful if inhaled, swallowed, or absorbed through skin. Do not breathe dust or mist. Avoid contact with eyes, skin, or clothing."
7. On page 4, revise "missed" to "mixed" in the second paragraph
8. Move the PPE section and User Safety Recommendations on pages 5 and 6 to before the Environmental Hazards section
9. On page 9, revise the Annual Maximum Use Rate section as follows:
 - a. Revise the first paragraph and "a" section to the following:
 - i. "A) If this product is tank mixed with an additional glyphosate product, or if another glyphosate product is sequentially applied: 1) the combined maximum application rate per year for all applications must not exceed 5.9 lbs a.i. glyphosate per acre per year. 2) Preplant, At-Planting, and Preeemergence applications must not exceed 3.6 lbs a.i. glyphosate per acre per year. 3) Total in-crop applications from emergence through the V8 stage or 30 inches must not exceed 1.5 lbs a.i. glyphosate per year. 4) Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature must not exceed 0.75 lbs a.i. glyphosate per year.
 - ii. Keep items B) and C) as they are.
10. On page 10, revise "16.5 pounds of active ingredient in water soluble packets" to "16.5 pounds of combined active ingredients in water soluble packets"
11. On page 11, revise "0.83 lbs of combined product is equivalent to 0.40 lbs....." to "The 0.83 lb rate of combined product is equivalent to 0.38 lbs glyphosate a.i., 0.30 lbs atrazine a.i., and 0.125 lbs dicamba a.i."
12. On page 11, revise "0.41 lbs of combined product is equivalent to...." To "The 0.41 lb rate of combined product is equivalent to 0.19 lbs glyphosate a.i., 0.15 lb atrazine a.i., and 0.06 lb dicamba a.i."
13. On page 11, move the Spray Solution and Area Treatment Table of Equivalents statements (up until, but not including, the new section Mixing with Water) to immediately following the Weed List table on page 18, and prior to the beginning of the RoundupReady Corn section.
14. On page 14, remove the statement "When applied under conditions described in the following paragraphs for shielded and hooded applications..... "ANNUAL WEEDS RATE TABLE" sections of this label." As this table no longer exists on this label.
15. On page 18, remove "in the "CROPS (ALPHABETICAL)" section of this label."
16. On page 18, revise the statement "Apply 0.41 pounds of this product per acre....." to "Apply the 0.41 lb rate of combined product (1-16.5 lbs packet containing 3 water soluble packs per 40 acres) when weeds are less than 6 inches tall, and the 0.83 lbs rate of combined product ((1-16.5 lbs packet containing 3 water soluble packs per 20 acres) when weeds are over 6 inches tall."
17. On page 19, remove the Maximum Allowable Combined Application Quantities Per Season table, as these were revised and incorporated into the Annual Maximum Use Rate section per item 7 above.
18. On page 19, add "Maximum of 2 applications per year" under the heading "Corn with the Roundup Ready or glyphosate resistant Gene"
19. On page 19, remove "NOTE: For maximum weed control, a postemergence (in-crop) application of this product...residual products listed above"
20. On page 20, revise "Sequential in-crop applications of this product" to "A second in-crop application of this product"

3094

21. On page 20, revise "1.4 pounds per acre per growing season" to "1.6 pounds of combine product per acre per year"
22. On page 20, revise "Allow a minimum of 50 days between application of this product and harvest of corn forage" to "Allow a minimum of 60 days between application of this product and harvest"
23. On page 20, revise "Allow a minimum of 7 days between application and harvest" to "Allow a minimum of 60 days between application and harvest"
24. On page 20, add "Apply a rate 0.41- 0.83 lbs combined product per acre" to the post-harvest Use Instructions. Revise "Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation" to "Allow a minimum of 60 days between treatment and harvest or feeding of treated vegetation"
25. On page 21, add "To the extent consistent with application law" before "For over-the-top uses on Roundup Ready....."
26. On page 22, revise "GLY N GO" to "GLYPHOSATE"
27. On page 23, remove the EPA Reg # and Est # for 80967-6. Also remove the Environmental Hazards and Storage and Disposal sections.
28. On page 25, revise "ASCADERA N GO" to "ATRAZINE" and remove the EPA Reg # and Est # for 80967-9. Also remove the Environmental Hazards and Storage and Disposal sections on pages 27, 28 and 29
29. On page 30, revise "RIOCAMBA N GO" to "DICAMBA" and remove the EPA Reg # and Est # for 80967-8. Also remove the Environmental Hazards and Storage and Disposal sections on pages 31 and 32
30. On page 34, remove the Environmental Hazards section.
31. On page 34, revise "this case treats 20 acres at the lower use rate and 11.5 acres at the higher use rate" to "this case treats 40 acres at the lower use rate and 20 acres at the higher use rate."

The basic formulation CSF [dated 9-27-2010] of the product referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable. The basic CSF will be added to your file.

You will submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records. If you have any questions, please contact Hope Johnson at 703-305-5410.


Jim Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505P)

4 of 4

Master Label containing:
 Sublabel A: Complete Booklet Label
 Sublabel B: Container Label for Glyphosate Solubag
 Sublabel C: Container Label for Atrazine Solubag
 Sublabel D: Container Label for Dicamba Solubag
 Sublabel E: Outer Box Container Label
 Sublabel F: N Go Component Label

GLY ATRA™ N GO™

Active Ingredients:

*N-(phosphonomethyl) glycine..... 95.0%
 Other Ingredients:.....5.0%
 Total..... 100.0%

*Contains 0.95 lb of acid per lb of product.

**Atrazine..... 96.5%
 Related compounds.....1.8%
 Other Ingredients:.....1.7%
 Total..... 100.0%

** This product contains 5.898 lbs a.i. per 6 lb water soluble bag

Dicamba 99.8%
 Related compounds:.....0.2%
 Total..... 100.0%

ACCEPTED
with COMMENTS
in EPA Letter Dated

SEP 30 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

80967-7

DANGER

Keep Out Of Reach Of Children

Read entire label before using this product.

EPA Reg. No.: 80967-XX

EPA Est. No.: XXXXX-XX-XXX

Net Contents: (1x8 lb+1x6 lb+1x2.5 lb); of product per case ** 3 solubags overap
packet, Bulk

Manufactured in China for:
 MEY Corporation
 121 S. Estes Drive, Ste. 101
 Chapel Hill, NC 27514
 USA.
 Tel. (919) 932-5800

5041

Sublabel A: Complete Booklet Label

Restricted Use Pesticide
Ground and Surface Water Concerns
For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

GLY ATRA™

Complete Directions for Use IN ROUNDUP READY or GLYPHOSATE RESISTANT CORN

GLY ATRA™ N GO MUST BE USED WITH N GO™ WATER CONDITIONER/BUFFER. DO NOT USE THIS PRODUCT WITH ANY OTHER SURFACTANTS, CONDITIONERS OR BUFFERING AGENTS. DO NOT USE THIS PRODUCT WITHOUT N GO™ WATER CONDITIONER/BUFFER.

Avoid Contact Of Herbicide With Foliage, Green Stems, Exposed Non-Woody Roots Or Fruit Of Crops (Except As Specified For Individual Roundup Ready® or glyphosate resistant corn Crops. Avoid application to Desirable Plants And Trees, Because Severe Injury Or Destruction May Result.

Read the entire label before using this product. Use only according to label instructions.

Active Ingredients:

*N-(phosphonomethyl) glycine.....	95.0%
Other Ingredients:.....	<u>5.0%</u>
Total.....	100.0%

*Contains 0.95 lb of acid per lb of product.

**Atrazine.....	96.5%
Related compounds.....	1.8%
Other Ingredients:.....	<u>1.7%</u>
Total.....	100.0%

** This product contains 5.898 lbs a.i. per 6 lb water soluble bag

Dicamba	99.8%
Related compounds:.....	<u>0.2%</u>
Total.....	100.0%

DANGER

Keep Out of Reach of Children

Read entire label before using this product.

EPA Reg. No.: 80967-T
EPA Est. No.: XXXXX-XX-XXX

Net Contents: (1x8 lb+1x6 lb+1x2.5 lb); of product per case ** 3 solubags overap packet, Bulk

Manufactured in China for:
MEY Corporation
121 S. Estes Drive, Ste. 101
Chapel Hill, NC 27514
USA.
Tel. (919) 932-5800

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. This product is identified as GLY ATRA N GO, EPA Reg. No. 80967-X. You may also contact 1-800 -222-1222 for emergency treatment information.</p>	
<p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals.

DANGER. Corrosive. Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust or mist. Causes irreversible eye damage. Avoid contact with eyes, skin, or clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Atrazine in GLY ATRA can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Product must not be missed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed/loaded or used within 50 feet of any wells, including abandoned wells, drainage wells, and sinkholes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum of 110% of the capacity of the largest pesticide container or application equipment of the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capabilities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticides shipments to the mixing/loading sites.

Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying atrazine to tile-outletted fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile-outletted fields.
- Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- Apply this product to the entire tile-outletted field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water

when disposing of equipment washwater.

PHYSICAL OR CHEMICAL HAZARDS

Solutions of this product must be mixed and using only stainless steel, fiberglass, plastic and plastic-lined containers.

DO NOT MIX, STORE OR UTILIZE THIS PRODUCT OR SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR TANKS. This product or solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This as mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators using spray equipment mounted on their backs must wear:

1. Coveralls over long-sleeved shirt and long pants,
2. Chemical-resistant footwear plus socks, and
3. Chemical-resistant gloves such as barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils, and
4. Face shield or goggles, and
5. A NIOSH approved respirator equipped with a dust/mist filter with MSHA/ NIOSH approval prefix TC-21C or any R, P, or HE filter.

Mixers, loaders, applicators, flaggers, and other handlers must wear:

1. Long sleeved shirt and long pants,
2. Chemical-resistant gloves such as barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils,
3. Shoes plus socks, and
4. Chemical resistant apron when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to concentrate.
5. Face shield or goggles, and
6. A NIOSH approved respirator equipped with a dust/mist filter with MSHA/ NIOSH approval prefix TC-21C or any R, P, or HE filter.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily

User Safety Recommendations:

1. Users must wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Users must remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Users must remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published MEY Corporation Supplemental Labeling.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.atrazine-watershed.info, or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact MEY Corporation for a refund.

This product is a restricted use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for Atrazine to reach ground and surface water.

Groundwater contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

Do not apply this product through any type of irrigation system.

When tank mixing or sequentially applying atrazine or products containing atrazine to corn, the total pounds of atrazine applied (lbs. ai/A) must not exceed 2.5 pounds active ingredient per year.

Check valves or anti-siphoning devices must be on all mixing equipment to prevent back-siphoning into wells or bulk storage tanks. See the Storage and Disposal section of the label regarding proper disposal of excess pesticide, spray mixtures and rinsate.

All applicable directions, restrictions and precautions are to be followed. This label must be in the possession of the actual product user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Only two applications of this product are allowed per year.

10041

Do not apply this product through aerial application.

THIS IS AN END-USE PRODUCT AND MEY CORPORATION, DOES NOT INTEND AND HAS NOT REGISTERED IT FOR FORMULATION, REFORMULATION, MANUFACTURING OR REPACKAGING USE.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 20 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls worn over short sleeve shirt and short pants,
- chemical resistant footwear plus socks,
- chemical-resistant gloves made of any waterproof material,
- chemical-resistant headgear for overhead exposure,
- protective eyewear,
- face shield or goggles, and
- a NIOSH approved respirator equipped with a dust/mist filter with MSHA/ NIOSH approval prefix TC-21C or any R, P, or HE filter.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticides Storage: Keep container closed to prevent spills and contamination.

Pesticide Disposal: Wastes of this product may be dangerous. Improper disposal of excess pesticide or rinse is a violation of Federal Law. If these wastes cannot be disposed of according to the label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Non-refillable container. Do not reuse outer bag or box. [For bulk product: Triple rinse promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.] Offer for recycling if available. Dispose of outer bag and box in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(How this product works)

Product Description: GLY ATRA N GO Herbicide is a three way herbicide with soil residual activity. It is generally non-selective and gives broad spectrum control of many annual weeds, and perennial weeds in glyphosate resistant corn. **(GLY ATRA N GO) (This product)** is a new (US PATENT 6,936,572 and 7,247,602) three part system where technical glyphosate is used in conjunction with atrazine and dicamba salts with an adjuvant. The **(GLY ATRA N GO) (This product)** MUST be used with the **N GO Water Conditioner/Buffer**. It may be applied through most standard industrial or field-type sprayers after dilution and mixing with water according to label instructions.

Do not add surfactants, additives containing surfactants, buffering agents, pH adjusting agents or ammonium sulfate to the spray solution. Drift control additives or dyes and colorants may be used. See the "MIXING" section of this label for instructions.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth

stages approaching maturity. Refer to the annual, perennial, woody brush and trees rate tables for rate specifications for specific weeds.

Always use the higher rate of this product per acre within the specified range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

Rainfastness: Heavy rainfall within 2 hours after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage must be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action: The multiple active ingredients in this product address an excellent resistance management strategy while placing a very low impact on the environment.

The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids. Micronutrients can be applied as part of a tank mix.

Soil Activity: Emerged weeds and non emerged weeds at the time of application are controlled by this product. Most Weeds germinating from seed after application will not be controlled until canopy is achieved.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Annual Maximum Use Rate:

If this product is tank mixed with an additional glyphosate product, or if another glyphosate product is sequentially applied, the combined maximum application rate per year for all applications must not exceed 6.2 lbs. active ingredient per acre per year.

a) The combined total of all treatments must not exceed 6.2 pounds of this product per acre per year. Preplant, At-planting, Preemergence applications 3.8 pounds per acre.

Total in-crop applications from emergence through the V8 1.6 pounds per acre stage or 30 inches

Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature 0.8 pounds per acre

(black layer formation) until 7 days before harvest

- b) If this product is tank mixed with an additional atrazine product, or if another atrazine product is sequentially applied, the total lbs active ingredient applies must not exceed 2.5 lbs active ingredient per acre per year.
- c) If this product is tank mixed with an additional dicamba product, or if another dicamba product is sequentially applied, the maximum application per year must not exceed 2.0 lbs ae per year.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CORN CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

MIXING

This package contains (1x8 lb+1x6 lb+1x2.5 lb) 16.5 pounds of active ingredient in water soluble packets of **(GLY ATRA N GO) (This product)** and one 2.5 gallon container of **N GO Water Conditioner/Buffer**. Do not handle the water soluble packets of **(GLY N GO) (This product)** with wet gloves or allow them to become wet prior to addition to the spray tank. Do not break open the water soluble packets of **(GLY ATRA N GO) (This product)** use entire contents of packet. If all packets of **(GLY ATRA N GO) (This product)** are not used, close and reseal outer container to protect remaining water soluble packets.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

Maintain sufficient agitation to keep the product in suspension at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed.

140941

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers must be no finer than 50 mesh.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

SPRAY SOLUTION AND AREA TREATMENT
TABLES OF EQUIVALENTS

Spray Solutions

One 16.5 lb total weight packet of GLY ATRA will treat 20 acres of glyphosate resistant corn at the total 0.83 lb per acre rate or 40 acres at the 0.41 lb per acre rate.

Area Treatment Equivalents

Upper and Lower Rates of Gly-ATRA Go (lbs./acre) Combined packet	Acres Treated By One 16.5 lb. packet containing 3 water soluble packs
0.41	40
0.83	20

0.83 lbs of combined product is equivalent to 0.40 lb (0.05 of a pack) glyphosate, 0.3 lb (0.05 of a pack) atrazine, and 0.127 lb (0.05 of a pack) dicamba.

0.41 lbs of combined product is equivalent to 0.2 lb glyphosate, 0.15 lb atrazine, and 0.06 lb dicamba

Mixing With Water

Mix spray solutions of this product as follows:

1. Fill the mixing or spray tank with at least one-half the required amount of water and start sufficient agitation to keep the product in suspension.
2. Add an appropriate amount of **N GO Water Conditioner/Buffer** for a spray solution of 0.5% to 1% by volume and mix well (if applying 10 gal of water volume per acre, mix 6.4 oz (0.5%v/v) or 12.8 oz (1%v/v) of **N GO Water Conditioner/Buffer**. Use upper rate of **N GO** with upper rate of **GLY ATRA™**. Or use 1%v/v rate for water volumes of 15 gal per acre.
3. Add the unopened (**GLY ATRA N GO**) (**This product**) packets to the water/ **N GO Water Conditioner/Buffer** mixture while maintaining sufficient agitation to keep the product in suspension. Add remaining water to spray tank directing the water stream at the floating packet(s) until the required water volume is reached. Water soluble packet(s) must completely dissolve in approximately five minutes.

Dissolution rate may be slowed by cold water, lack of proper agitation, or water containing high concentrations of boron or sulfur.

- 4. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations.
- 5. If during mixing and application, foaming of the spray solution occurs; use an approved anti-foam or defoaming agent.
- 6. Maintain sufficient agitation to keep the product in suspension at all times until spraying is completed.

Tank Mixing Procedure

Mix labeled tank mixtures of this product with water as follows:

- 1 Place a 20 to 35 mesh screen or welling basket over filling port or use an injector.
- 2. Through the screen, fill the spray tank one-half full with water and start sufficient agitation to keep the product in suspension.
- 3. Add an appropriate amount of **N GO Water Conditioner/Buffer** for a spray solution of 0.5% to 1% by volume and mix well.
- 4. Add the unopened (**GLY ATRA N GO**) (**This product**) packets to the water/**GO Water Conditioner/Buffer** mixture while maintaining sufficient agitation to keep the product in suspension in spray tank or injector tank. Add remaining water to spray tank or into injector tank (system,), directing the water stream at the floating packet(s) until the required water volume is reached. Water soluble packet(s) must completely dissolve in approximately five minutes. Dissolution rate may be slowed by cold water, lack of proper agitation, or water containing high concentrations of boron or sulfur.
- 5. If during mixing and application, foaming of the spray solution occurs, use an approved anti-foam or defoaming agent.
- 6. Maintain sufficient agitation to keep the product in suspension at all times until the contents of the tank are sprayed

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Ammonium Sulfate

DO NOT use ammonium sulfate with this product.

Colorants or Dyes

Agriculturally-approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance,

160041

especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's rate specifications.

Drift Control Additives

Drift control additives may be used with all equipment types. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Surfactants

Use **N GO Water Conditioner/Buffer** with this product. Other surfactants approved for food use that are compatible with the active ingredients in this product can be added.

Buffering and pH Adjusting Agents

Use only **N GO Water Conditioner/Buffer** with this product. Any other additional buffering and pH adjusting agents are not required.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Ground Broadcast Spray--Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held Spray Equipment – Handguns, handwands, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. Agitation must be maintained at all time.

Selective Equipment--Recirculating sprayers and shielded and hooded sprayers.

Controlled Droplet Applicator (CDA)--Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

Ground Broadcast Equipment

Use the specified rates of this product in 3 to 20 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume must be increased within the specified range to ensure complete coverage.

Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

Hand-Held Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only. For specified rates and timing, refer to the "ANNUAL WEEDS -- HAND-HELD EQUIPMENT" section of this product label.

Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, and hooded sprayers, after dilution and thorough mixing with water, to listed weeds growing in any non crop site specified on this label.

In cropping systems, hooded sprayers and shielded sprayers in row middles (in between rows of crop plants) where any dripping or leaking will not contact crop foliage. Such equipment must be capable of preventing all crop contact with herbicide solutions and operated without leakage of spray mists or dripping onto crop.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation must be adjusted so that the lowest spray stream contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops must be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Recirculating spray system

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

Shielded and hooded applicators

When applied under the conditions described in the following paragraphs for shielded and hooded applications, this product at specified rates will control those weeds listed in the "ANNUAL WEEDS RATE TABLE" sections of this label. A hooded sprayer is a type of shielded applicator where the spray pattern

180241

is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Keep shields on these sprayers adjusted to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure/low drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is specified. Spray volume must be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- The spray hoods must be operated on the ground or skimming across the ground.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood must be 30 inches.
- Maximum tractor speed: 5 mph to avoid bouncing of the spray hoods.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

CDA Equipment

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount specified in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

CORN USES

For Use on Round Up Ready Corn. Do not use on sweet corn.

TABLE Weeds List

ANNUALS	
Alkanet	Mustard, Black, Blue, Tansy, Treacle, Tumble, Wild
Amaranth, Palmer, Powell, Spiny	Yellowtops
Aster, Slender	Narrow-leaved Hawk's Beard
Bedstraw, Catchweed	Nightshade, Black, Cutleaf
Beggarweed, Florida	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
Broomweed, Common	Pigweed, Redroot, Smooth
Buckwheat, Tartary, Wild	Pineappleweed
Buffalobur	Poorjoe
Burclover, California	Poppy, Red-horned
Burcucumber	Prickly Lettuce
Buttercup, Corn, Creeping, Roughseed	Puncturevine
Western Field	Purslane, Common
Carpetweed	Pusley, Florida
Catchfly, Nightflowering	Radish, Wild
Chamomile, Corn	Ragweed, Common, Giant (Buffaloweed), Lance-Leaf
Chervil, Bur	Rocket, London, Yellow
Chickweed, Common	Rubberweed, Bitter (Bitterweed)
Cleavers	Salsify
Clovers	Senna, Coffee,
Cockle, Corn, Cow, White	Sesbania, Hemp
Cocklebur, Common	Shepherd's Purse
Copperleaf, Hophornbeam	Sicklepod
Cornflower (Bachelor Button)	Sida, Prickly (Teaweed)
Corn Spurry	Smartweed, Green, Pennsylvania
Croton, Tropic, Woolly	Sneezeweed, Bitter
Daisy, English	Sowthistle, Annual, Spiny
Descurania Sophia	Spanish Needles
Dragonhead, American	Spikeweed, Common
Eastern Black Flowering Nightshade	Spurge, Prostrate, Leafy
Eveningprimrose, Cutleaf	Spurry, Corn
Falseflax, Smallseed	Starbur, Bristly
Fleabane, Annual	Starwort, Little
Flixweed	Stinkweed
Fumitory	Sumpweed, Rough
Green Smartweed	Sunflower, Common (Wild), Volunteer
Goosefoot, Nettleleaf	Thistle, Russian
Hempnettle	Velvetleaf
Henbit	Volunteer Canola
Jacobs-ladder	Volunteer Flax
Jimsonweed	Waterhemp
Knawel (German Moss)	Waterprimrose, Winged
Knotweed, Prostrate	Wormwood
Kochia	Wild Buckwheat

Ladys thumb	Wild Mustard
Lambs quarters, Common	Wild Tomato
Lettuce, Miners, Prickly	
Mallow, Common, Venice	
Marestall (Horseweed)	
Mayweed	
Morningglory, Ivyleaf, Tall	

ANNUAL GRASSES	
Barnyard Grass	Persian Darnel
Blue Grass	Volunteer Barley
Crab Grass, Large, (Hairy), Smooth	Volunteer Corn
Downy Brome	Volunteer Wheat
Foxtail, Giant, Green	Wild Oats
Groundcherry	Yellow Foxtail

BIENNIALS	
Burdock, Common	Mallow, Dwarf
Carrot, Wild (Queen Anne's Lace)	Plantain, Bracted
Cockle, White	Ragwort, Tansy
Eveningprimrose, Common	Starthistle, Yellow
Geranium, Carolina	Sweetclover
Gromwell	Teasel
Knapweed, Diffuse, Spotted	Thistle, Bull, Milk, Musk, Plumeless

PERENNIALS	
Alfalfa(1)	Milkweed, Climbing, Common, Honeyvine, Western Whorled
Artichoke, Jerusalem	Nettle, Stinging
Aster, Spiny, White heath	Nightshade, Silverleaf (White Horsenettle)
Bedstraw, Smooth	Onion, Wild
Bindweed, Field, Hedge	Plantain, Broadleaf, Buckhorn
Blueweed, Texas	Poison Ivy
Bursage, Woolly leaf (1) (Bur Ragweed Poverty weed)	Pokeweed
Buttercup, Tall	Purple Loosestrife
Campion, Bladder	Ragweed, Western
Chickweed, Field, Mouseear	Redvine
Chicory(1)	Sericia Lespedeza
Clover(1), Hop	Smartweed, Swamp
Cottontop	Snakeweed, Broom
Dandelion(1),	Sorrel(1), Red (Sheep Sorrel)
Dock(1), Broadleaf (Bitterdock), Curly	Sowthistle(1), Perennial
Dogbane, Hemp	Spurge, Leafy
Dogfennel(1) (Cypressweed)	Sundrop,
Fern, Bracken	Thistle, Canada, Scotch
Garlic, Wild	Toad flax, Dalmation
Goldenrod, Canada, Missouri	Tropical Soda Apple
Goldenweed, Common	Trumpetcreeper (Buckvine)
Hawkweed	Vetch
Henbane, Black(1)	Water hemlock, Spotted
Hoary Cress	Water primrose, Creeping
Horsenettle, Carolina	Wood sorrel(1), Creeping, Yellow
Ironweed	Wormwood, Louisiana

Knapweed, Black, Diffuse, Russian(1) Spotted	Yankee weed
Knotweed	Yarrow, Common(1)
PERENNIAL GRASSES	
Blue Grass, Canada, Kentucky	Foxtail Barley
Brome Grass	Quackgrass
Cattail	Yellow Nutsedge

- (1) Noted perennials may be controlled using lower rates of this product than those recommended for other listed perennial weeds.
- (2) Growth Suppression only.

ROUNDUP READY and Glyphosate resistant Corn

The following instructions include all applications which can be made onto Roundup Ready® and glyphosate resistant corn during the complete cropping season. Do NOT combine these instructions with other rate specifications made for crop varieties which do not contain the Roundup Ready gene, in the "CROPS (ALPHABETICAL)" Section of this label.

MEY CORPORATION RECOMMENDS USE OF THIS PRODUCT FOR POSTEMERGENCE APPLICATION ONLY ON CROP VARIETIES DESIGNATED AS CONTAINING THE ROUNDUP READY or GLYPHOSATE RESISTANT GENE.

Apply 0.41 pounds of this product per acre when weeds are less than 6 inches tall, and 0.83 pounds when weeds are over 6 inches tall.

Applying this product to crop varieties which are not designated as Roundup Ready or glyphosate resistant will result in severe crop injury and yield loss. Avoid contact with foliage, green stems or any desirable plants which do not contain the Roundup Ready or glyphosate resistant gene, since severe injury or destruction will result.

The Roundup Ready or glyphosate resistant designation indicates that the corn variety contains a patented gene which provides tolerance to glyphosate herbicides. Information on Roundup Ready or glyphosate resistant corn varieties may be obtained from your seed supplier.

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment use flat spray nozzles. Check for even distribution of spray droplets.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

See the "MIXING and APPLICATION EQUIPMENT AND TECHNIQUES" sections of this label for additional directions and restrictions on the application of this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT specified for over-the-top applications of this product unless otherwise noted in this product label.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

Corn with the Roundup Ready or glyphosate resistant Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence (in-crop), Spot Treatment, Preharvest, Post-Harvest.

Maximum Allowable Combined Application Quantities Per Season

Combined total per year for all applications	1.4 pounds per acre
Preplant, At-planting, Preemergence applications	0.41 pounds per acre
Total in-crop applications from emergence through the V8 30 inches	0.83 pounds per or acre stage
Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest	0.83 pounds per acre

Preplant, Preemergence and At-Planting

USE INSTRUCTIONS: This product may be applied alone before, during or after planting corn.

NOTE: For maximum weed control, a postemergence (in-crop) application of this product should be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches which ever comes first.

When applied as directed, this product controls labeled annual grass and broadleaf

weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. The postemergent application of 0.41 to 0.83 pounds per acre of this product should be made before the weeds reach a height and/or density that the weeds become competitive with the crop, generally 4 inch tall weeds or less.

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on this label. If new flushes of weeds occur, a sequential application of this product at 0.41 to 0.83 pounds per acre will control the labeled grasses and broadleaf weeds.

PRECAUTIONS, RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for general precautionary instructions for use in Roundup Ready crops. Single in-crop applications of this product are not to exceed 0.83 pounds per acre. Sequential in-crop applications of this product from emergence through the V8 stage or 30 inches must not exceed 1.4 pounds per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage.

Preharvest

USE INSTRUCTIONS: In Roundup Ready corn, up to 0.41 pounds per acre of this product can be applied preharvest. Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest.

PRECAUTIONS, RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

LIMIT OF WARRANTY AND LIABILITY

MEY Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

240441

To the extent consistent with applicable law, buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise. To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

For over-the-top uses on Roundup Ready and glyphosate resistant crop varieties crop safety and weed control performance are not warranted by MEY Corporation when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

TRADEMARKS

GLY ATRA™ is a trademark of MEY Corporation Inc.

ROUND UP READY is a registered trademark of Monsanto Chemical Company

N GO™ is a trademark of MEY Corporation Inc.

GLY ATRA™ AND N GO™ are projected by US patents 7,247,602 and 6,936,572

Sublabel B: Container Label for Glyphosate Solubag

GLY N GO
Herbicide

Active Ingredient:

*N-(phosphonomethyl) glycine.....95.0%

Other Ingredients:5.0%

Total:.....100.0%

*Contains 0.95 lb of acid per lb of product.

Keep Out of Reach of Children

PELIGRO DANGER

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Read entire label before using this product.

Note: This statement will appear on the foil wrapper that holds the soluble package of GLY N GO Herbicide. (REMOVE WATER SOLUBLE PACKET FROM FOIL POUCH BEFORE USING)

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

210041

If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. This product is identified as GLY ATRA N GO™ HERBICIDE, EPA Reg. No. 80967-6. You may also contact 1-800 -222-1222 for emergency treatment information.</p>	
<p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

EPA Reg. No. 80967-6
EPA Est. No.: XXXXX-XX-XXX

Manufactured in China for:
 MEY Corporation
 121 S. Estes Drive, Ste. 101
 Chapel Hill, NC 27514

Net Weight: (bulk), (X soluble bags containing 8 lbs. each)

Hazards to Humans and Domestic Animals.

DANGER. Corrosive. Causes irreversible eye damage. Harmful if inhaled or swallowed. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Personal Protective Equipment (PPE)
 Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, protective eyewear, and chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

See Label Booklet for complete directions for use, warnings and cautions and first aid information.

270941

Do not open.

Use entire contents of this water soluble package at one time following mixing instructions contained in the label booklet.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticides Storage: Keep container closed to prevent spills and contamination.

Pesticide Disposal: Wastes of this product may be dangerous. Improper disposal of excess pesticide or rinse is a violation of Federal Law. If these wastes cannot be disposed of according to the label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Non-refillable container. Do not reuse outer bag or box. [*For bulk product:* Triple rinse promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.] Offer for recycling if available. Dispose of outer bag and box in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Sublabel C: Container Label for Atrazine Solubag

Restricted Use Pesticide
Ground and Surface Water Concerns
For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.
Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

ASCADERA N GO Herbicide

Read the entire label before using this product. Use only according to label instructions.

Active Ingredient:

Atrazine.....	.96.5%
Related compounds.....	1.8%
Other Ingredients:.....	1.7%
Total.....	100.0%

This product contains 5.898 lbs a.i. per 6 lb water soluble bag

Keep Out of Reach of Children

CAUTION

Read entire label before using this product.

Note: This statement will appear on the foil wrapper that holds the soluble package of ASCADERA N GO Herbicide. (REMOVE WATER SOLUBLE PACKET FROM FOIL POUCH BEFORE USING)

EPA Reg. No.: 80967-9
EPA Est. No.: XXXXX-XXX-XX

Net Weight: XX lbs.

Manufactured for:
MEY Corporation
121 S. Estes Drive, Ste. 101

29041

Chapel Hill, NC 27514
USA.
Tel. (919) 932-5800

Net Contents: Net Weight: (bulk), (X soluble bags containing 6 lbs. each)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. This product is identified as ASCADERA N GO™ HERBICIDE, EPA Reg. No. 80967-XX. You may also contact 1-800 –222-1222 for emergency treatment information.</p>	
<p>NOTE TO PHYSICIAN: There is no specific antidote for Atrazine. If this product is ingested, induce emesis or lavage stomach. The use of aqueous slurry of activated charcoal may be considered.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals.

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust or mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils; or viton > 14 mils. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators using spray equipment mounted on their backs must wear:

7. Coveralls over long-sleeved shirt and long pants,
8. Chemical-resistant footwear plus socks, and
9. Chemical-resistant gloves such as barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils.

Mixers, loaders, applicators, flaggers, and other handlers must wear:

5. Long sleeved shirt and long pants,
6. Chemical-resistant gloves such as barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils,
10. Shoes plus socks, and
11. Chemical resistant apron when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed/loaded or used within 50 feet of any wells, including abandoned wells, drainage wells, and sinkholes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest

load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum of 110% of the capacity of the largest pesticide container or application equipment of the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capabilities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticides shipments to the mixing/loading sites.

Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying atrazine to tile-outletted fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile-outletted fields.
- Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- Apply this product to the entire tile-outletted field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticides Storage: Keep container closed to prevent spills and contamination.

Pesticide Disposal: Wastes of this product may be dangerous. Improper disposal of excess pesticide or rinse is a violation of Federal Law. If these wastes cannot be disposed of according to the label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Non-refillable container. Do not reuse outer bag or box. [*For bulk product:* Triple rinse promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.] Offer for recycling if available. Dispose of outer bag and box in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

330041

Sublabel D: Container Label for Dicamba Solubag

RIOCAMBA N GO Herbicide

Read the entire label before using this product. Use only according to label instructions.

Active Ingredient:

Dicamba	99.8%
Related compounds:.....	0.2%
Total.....	100.0%

Keep Out of Reach of Children

WARNING

Read entire label before using this product.

Note: This statement will appear on the foil wrapper that holds the soluble package of RIOCAMBA N GO Herbicide. (REMOVE WATER SOLUBLE PACKET FROM FOIL POUCH BEFORE USING)

EPA Reg. No.: 80967-8
EPA Est. No.: XXXXX-XXX-XX

Net Weight: Net Weight: Bulk, (X solubags containing 2.5 lbs. each)

Manufactured for:
MEY Corporation
121 S. Estes Drive, Ste. 101
Chapel Hill, NC 27514
USA.
Tel. (919) 932-5800

FIRST AID

If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
-------------------	---

34041

If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. This product is identified as RIOCAMBA N GO™ HERBICIDE, EPA Reg. No. 80967-8. You may also contact 1-800 -222-1222 for emergency treatment information.</p>	
<p>NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes eye irritation. **DO NOT** get in eyes, on skin or on clothing. Harmful if swallowed. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are barrier laminate >14 mils, neoprene rubber >14 mils, polyvinyl chloride (PVC) > 14 mils, or viton > 14 mils. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

7. Long sleeved shirt and long pants,
8. Shoes plus socks, and
9. Chemical-resistant gloves (except for applicators using groundboom equipment, pilots and flaggers,

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticides Storage: Keep container closed to prevent spills and contamination.

Pesticide Disposal: Wastes of this product may be dangerous. Improper disposal of excess pesticide or rinse is a violation of Federal Law. If these wastes cannot be disposed of according to the label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Non-refillable container. Do not reuse outer bag or box. [For bulk product: Triple rinse promptly after emptying. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.] Offer for recycling if available. Dispose of outer bag and box in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

310041

Sublabel E: Outer Box Container Label

Restricted Use Pesticide

Ground and Surface Water Concerns

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

GLY ATRA™

For Use In ROUNDUP READY or GLYPHOSATE RESISTANT CORN

370941

GLY ATRA™ N GO™

NO GO Water Conditioner / Buffer MUST BE USED used with GLY ATRA™ N GO™

Read the entire label before using this product. Use only according to label instructions.

Active Ingredients:

N-(phosphonomethyl) glycine.....	95.0%
Other Ingredients:.....	5.0%
Total.....	100.0%

*Contains 0.95 lb of acid per lb of product.

**Atrazine.....	96.5%
Related compounds.....	1.8%
Other Ingredients:.....	1.7%
Total.....	100.0%

** This product contains 5.898 lbs a.i. per 6 lb water soluble bag

Dicamba	99.8%
Related compounds:.....	0.2%
Total.....	100.0%

PRECAUTIONARY STATEMENTS Keep Out Of Reach Of Children

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Hazards to Humans and Domestic Animals

DANGER - CORROSIVE: Causes irreversible eye damage. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Harmful if swallowed. Avoid breathing dust or spray mist. Do not get in eyes or on clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing equipment wash waters or rinsate.

First Aid Information of label Booklet.

Net Contents: Net Contents: (1x8 lb+1x6 lb+1x2.5 lb) in 3 water soluble bags; this case treats 20 acres at the lower use rate and 11.5 acres at the higher use rate

EPA Reg No: 80967-T

Made In China

N GO™

Water Conditioner / Buffer

Water Conditioner / Buffer for use with GLY ATRA™ N GO™

Read the entire label before using this product. Use only according to label instructions.

Monocarbamide dihydrogen sulphate, polyethoxylated amines and esters	82%
Water	18%
Total	100%

Net Contents: 2.5 Gallons

386041

EPA Est. No.: XXXXX-XX-XXX

Manufactured in China for:
MEY Corporation
121 S. Estes Drive, Ste. 101
Chapel Hill, NC 27514
USA.
Tel. (919) 932-5800

390941

Sublabel F: N Go Component Label

**MEY CORPORATION
N GO™**

WATER CONDITIONER / pH ADJUSTER For (GLYPHOSATE) (ATRAZINE) (DICAMBA MIXTURES))

PRINCIPAL FUNCTIONING AGENTS

Monocarbamide dihydrogen sulfate	65%
Proprietary blend of amine phosphates, viscosity reducing agents	
Inerts	35%
Total	100%

All ingredients are approved for use under CFR40, 180, 1001 (c)

DANGER: KEEP OUT OF REACH OF CHILDREN, CAUSES EYE BURNS, CAUSES SKIN IRRITATION. HARMFUL IF SWALLOWED.

PRECAUTIONARY STATEMENTS:

Before using this product, read the entire label, including the conditions of sale. Corrosive: May cause eye, skin and respiratory irritation. May cause allergic skin reactions. May cause severe burns. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators or other handlers of this chemical must wear: Goggles or face shield and chemical resistant gloves when handling. Harmful if inhaled. Do not breathe vapors or spray mist. Wear air purifying mask with appropriate canister. Harmful if swallowed. Thoroughly wash exposed areas of skin after use. Flush with clean water for at least 30 minutes. Seek medical advice immediately. Consult the MSDS for further information.

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

400f41

PRODUCT INFORMATION:

N GO is a specialized utility modifier that acts as a pH adjuster and water conditioner with built in compatibility agents that widen the conditions under which atrazine, dicamba, and glyphosate and other weak acid herbicides can be applied. N GO maintains the integrity of the spray diluent (water) and protects herbicides from deactivation by hard water cations and micronutrient based tank mix partners.

DIRECTIONS FOR USE:

Always mix and apply this product in accordance with the instructions found on the pesticide label regarding adjuvants. In the absence of specific instructions follow the application and mixing guidelines below.

APPLICATION RATES:

Using the guidelines below apply N GO at 2-4 quarts per 100 U.S. gallons of spray solution (2-4 liters per 400 liters of spray solution) or 0.5%-1.0% v/v.

Add an appropriate amount of **N GO Water Conditioner/Buffer** for a spray solution of 0.5% to 1% by volume and mix well (if applying 10 gal of water volume per acre, mix 6.4 oz (0.5%v/v) or 12.8 oz (1%v/v) of **N GO Water Conditioner/Buffer**. **Use upper rate of N GO with upper rate of GLY ATRA™ Or use 1%v/v rate for water volumes of 15 gal per acre**

MIXING INSTRUCTIONS:

Fill the spray tank $\frac{3}{4}$ full of desired water volume. While the water is agitating, add tank mix ingredients in the following order: add 1) N GO at 0.5% – 1% v/v or (2-4 quarts per 100 U.S. gallons of spray solution); 2) GLY ATRA™ 3) Wait until previously added products have completely dissolved, or suspended; Continue agitation and complete filling the spray tank.

STORAGE AND DISPOSAL:

Product does not freeze! Keep container closed tightly when not in use. Prevent water from being introduced to the contents of this container. Do not reuse empty container. Do not store near heat or open flame. Do not store with oxidizing agents. Rinse container thoroughly, disposing of rinsate and the container in accordance with federal, state and local regulations.

CONDITIONS OF SALE:

Read the information contained herein before buying or using this product. If the stated terms are unacceptable, return the product at once, unopened. It is critical that the product be used and mixed only as specified on this label. Neither the manufacturer nor the seller makes any representation or warranty, expressed or implied, with respect to the results from the use of this material. Buyer and user assume all risks of use and/or handling. MEY CORPORATION warrants that this material is reasonably fit for use as specified on this label. No agent or representative is authorized to make any other representations concerning this material. Unforeseen factors beyond MEY'S control prevent elimination of risks in connection with the use of its chemicals. Such risks include, but are not limited to, damage to plants and/or crops to which the material is applied, or lack of complete control and damage caused by drift to other plants or crops. Such risks may occur even though the product is reasonably fit for use as stated heron

410241

and even though label directions are followed. Follow directions carefully. Timing, mixture, method of application, weather and other conditions are influencing factors in the use of this product and are beyond the control of the seller. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of the seller for any and all losses, personal injuries or damages resulting from the use of this product shall be the purchase price paid by the user or buyer for the product involved.

Licensed under U.S. Patent No. 6,936,572 & 7,247,602
MEY CORPORATION
121 S. Estes Drive Ste.01
CHAPEL HILL NC 27514

For 24-hour technical assistance when using this product in the USA, call: 1-800-
For 24-hour emergency service when using this product in the USA, call CHEMTREC: 1-
800-424-9300
N GO is a trademark of MEY CORPORATION
Made in the USA

Net Contents: 2.5 U.S. Gallons (9.46L)