

80967-13

6/12/2014

1/127



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Micah T. Reynolds
MEY Corporation
121 S. Estes Drive, Suite 101
Chapel Hill, North Carolina 27514

JUN 12 2014

Subject: Notification; Per PR-Notice 98-10
GlySupreme 75-WDG
EPA Reg. No. 80967-13
Date Submitted: June 9, 2014

Proposed Alternate Brand Name:
GlySupreme 75-WDG Glyphosate Herbicide

Dear Mr. Reynolds:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated June 9, 2014 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records. In addition, the alternate brand name "*GlySupreme 75-WDG Glyphosate Herbicide*" is acceptable and has been added to the product file.

If you have any questions regarding this letter, please contact me at (703) 306-0415 or davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

2/127

| | | | |
|--|---|--|-----------------------|
|  | United States Environmental Protection Agency Washington, DC 20460 | <input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other | OPP Identifier Number |
|--|---|--|-----------------------|

Application for Pesticide – Section I

| | | |
|--|--|---|
| 1. Company/Product Number <p style="text-align: center;">80967-13</p> | 2. EPA Product Manager <p style="text-align: center;">Kable (Bo) Davis</p> | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) <p style="text-align: center;">GlySupreme 75-WDG</p> | PM# <p style="text-align: center;">25/Herbicide Branch</p> | |
| 5. Name And Address Of Applicant (Include ZIP Code) <p>MEY Corporation 121 S. Estes Drive, Suite 101 Chapel Hill, North Carolina 27514</p> <input type="checkbox"/> Check if this is a new address | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ |

Section II

| | |
|--|---|
| <input type="checkbox"/> Amendment – Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification – Explain below. | <input type="checkbox"/> Final Printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other – Explain Below. |
|--|---|

NOTIFICATION
JUN 12 2014

Explanation: Use additional page(s) if necessary. (For section I and Section II.)


Submission of Label Notification to add label version date (for tracking purposes) and one additional alternate brand name (ABN).

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Please confirm receipt with Micah Reynolds: mreynolds@tsqusa.com, via phone (202) 828-8967, or via fax (202) 872-0745.

Section III

| | | | | | |
|--|--|---|---|---|--|
| 1. Material This Product Will Be Packaged In: | | | | | |
| Child Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No | Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | 2. Type of Container | | |
| * Certification must be submitted | | | If "Yes" Unit Packaging wgt. 20 kg | No. per Container 4 X 5 kg | <input checked="" type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| 3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container | | 4. Size(S) Retail Container 5 kg, 10 kg, 15 kg, 20 kg bag-in-box | | 5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input checked="" type="checkbox"/> On Labeling accompanying product | |
| 6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithographed <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Pager glued <input type="checkbox"/> Stenciled | | | | | |

Section IV

| | | |
|---|---|--|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name Micah T. Reynolds, Technology Sciences Group, Inc. | Title Senior Regulatory Consultant | Telephone No. (Include Area Code) (202) 828-8967 |
| Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | 6. Date Application Received _____ (Stamped) |
| 2. Signature  | 3. Title Regulatory Consultant to MEY Corporation | |
| 4. Typed Name Micah T. Reynolds | 5. Date June 9, 2014 | |

3/127



Technology Sciences Group Inc.
1150 18th Street, Suite 1000
Washington, DC 20036
Direct: (202) 828-8967
Fax: (202) 872-0745
E-Mail: MReynolds@TSGUSA.com

Micah T. Reynolds
Senior Regulatory Consultant

June 9, 2014

MEY Corporation
Submission of Label Notification for
GlySupreme 75-WDG (EPA Reg. No. 80967-13)
NOTIFICATION

JUN 12 2014

Mr. Bo Davis, PM-25
Registration Division (7505P)
Office of Pesticide Programs (NOTIF)
U.S. Environmental Protection Agency
Document Processing Center
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, Virginia 22202

Dear Mr. Davis:

Technology Sciences Group, Inc., on behalf of MEY Corporation, is submitting the enclosed label notification for the above referenced product to add an optional version date or revision date indicator to assist in tracking label changes for future production runs and label campaigns for this product. Also, an alternate brand name for this product is being added. The alternate brand name, "GlySupreme 75-WDG Glyphosate Herbicide" is being added. Please find enclosed the following documents supporting this notification:

- 1) EPA Notification Form (EPA Form 8570-1);
- 2) One (1) Copy of the revised proposed label with changes indicated;
- 3) One (1) Copy of the revised clean copy of the label.

Due to the length of this label, I am happy to provide PDF electronic files of the redline and clean label versions for ease of review and comparison. **Pages of this label affected by the additions noted above include the following pages only: 2, 83, 84, 87, 88, 120, 121, and 124.** If you have any questions or require additional information or clarification, please do not hesitate to contact me by phone at (202) 828-8967 or by e-mail at mreynolds@tsgusa.com.

Sincerely,

Regulatory Consultant to MEY Corporation

Enclosures

CC: K. Martyn, L. Hodges (MEY)

Washington, D.C.
1150 18th St., NW, Suite 1000
Washington, D.C. 20036
Phone: (202) 223-4392

California
712 Fifth St., Suite A
Davis, CA 95616
Phone: (530) 757-1245

Canada
275 Slater St., Suite 900
Ottawa, Ontario K1P 5H9
Phone: (613) 247-6285

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

GLYSUPREME 75-WDG

Sublabel A: Agricultural Use Label
Sublabel B: Industrial, Turf, and Ornamental Use Label

EPA Reg. No.: 80967-13
EPA Establishment No.: [73432-CHN-001]

Manufactured [for][by]:
MEY Corporation
121 S. Estes Drive, Suite 101
Chapel Hill, NC 27514 USA

NOTIFICATION
JUN 12 2014

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

SUBLABEL A: Agricultural Use Label

GlySupreme 75-WDG

[Water Soluble Dry Herbicide] [Water Dispersible Granule [Herbicide]]

(Alternate Brand Names: GlySupreme 75-Water Dispersible Granule, GlySupreme 75-WDG Herbicide, GlySupreme 75-WDG Glyphosate Herbicide)

This product contains surfactants and ammonium sulphate in the granule resulting in superior weed control.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees because severe injury or destruction may result.

Read the entire label before using this product. Use only according to label instructions.

CONTENTS

- 1 1.0 INGREDIENTS AND FRONT PANEL STATEMENTS
- 2 2.0 IMPORTANT PHONE NUMBERS
- 3 3.0 PRECAUTIONARY STATEMENTS
 - 3.1 Hazards to Humans and Domestic Animals
 - 3.2 Environmental Hazards
 - 3.3 Physical or Chemical Hazards
 - 3.4 Directions for Use/WPS
- 4 4.0 STORAGE AND DISPOSAL
- 5 5.0 INFORMATION
 - (How This Product Works)
- 6 6.0 MIXING
 - 6.1 Mixing with Water
 - 6.2 Tank Mixing Procedure
 - 6.3 Mixing for Hand-Held Sprayers
 - 6.4 Ammonium Sulfate
 - 6.5 Colorants or Dyes
 - 6.6 Surfactants
 - 6.7 Drift Reduction Additives
- 7 7.0 APPLICATION EQUIPMENT AND TECHNIQUES
 - 7.1 Aerial Equipment, Spray Drift Management, and Application Restrictions
 - 7.2 Ground Broadcast Equipment
 - 7.3 Hand-Held or High-Volume Equipment
 - 7.4 Selective Equipment
 - 7.5 Injection Systems
 - 7.6 CDA Equipment
- 8 8.0 ANNUAL AND PERENNIAL CROPS (Alphabetical)
 - 8.1 Cereal and Grain Crops

Deleted: 1
Deleted: March 19,

6/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- 8.2 Corn
- 8.3 Cotton
- 8.4 Fallow Systems
- 8.5 Grain Sorghum (Milo)
- 8.6 Herbs and Spices
- 8.7 Oil Seed Crops
- 8.8 Soybeans
- 8.9 Sugarcane
- 8.10 Vegetable Crops
- 8.11 Miscellaneous Crops

- 9 9.0 TREE, VINE, AND SHRUB CROPS (Alphabetical)
- 9.1 Berry Crops
- 9.2 Citrus
- 9.3 Miscellaneous Tree Food Crops
- 9.4 Non-Food Tree Crops
- 9.5 Pome Fruit
- 9.6 Stone Fruit
- 9.7 Tree Nuts
- 9.8 Tropical and Subtropical Trees and Fruits
- 9.9 Vine Crops

- 10 10.0 PASTURE GRASSES, FORAGE LEGUMES, AND RANGELANDS
- 10.1 Alfalfa, Clove, and Other Forage Legumes
- 10.2 Conservation Reserve Program (CRP)
- 10.3 Grass Seed or Sod Production
- 10.4 Pastures
- 10.5 Rangelands

- 11 11.0 ROUNDUP READY CROPS
- 11.1 Alfalfa with the Roundup Ready Gene
- 11.2 Canola with the Roundup Ready Gene
- 11.3 Corn with the Round Ready Gene
- 11.4 Soybeans with the Roundup Ready Gene
- 11.5 Sugar beets with the Roundup Ready Gene
- 11.6 Cotton with the Roundup Ready Gene
- 11.7 Seed Production of Select Crops with the Roundup Ready Gene

- 12 12.0 NON-CROP USES AROUND THE FARMSTEAD
- 12.1 Weed Control and Trim-And-Edge
- 12.2 Greenhouse/Shadehouse
- 12.3 Chemical Mowing
- 12.4 Cut Stump
- 12.5 Habitat Management

- 13 13.0 ANNUAL WEEDS RATE SECTION
(Alphabetically by Species)
- 13.1 Annual Weeds – Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K
- 13.2 Annual Weeds – Hand-Held or High-Volume Equipment
- 13.3 Annual Weeds – Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

- 14 14.0 PERENNIAL WEEDS RATE SECTION

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

(Alphabetically by Species)

- 15 15.0 WOODY BRUSH AND TREES RATE SECTION
(Alphabetically by Species)
- 16 16.0 WEED RESISTANCE MANAGEMENT
 - 16.1 Weed Resistance Management Recommendations
 - 16.2 Glyphosate-Resistant Biotypes Management
 - 16.3 Control and Management of Glyphosate-Resistant Horseweed in Cotton, Corn, and Soybean
- 17 17.0 LIMIT OF WARRANTY AND LIABILITY

1.0 INGREDIENTS AND FRONT PANEL STATEMENTS

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its ammonium salt.....75.7%
 Other Ingredients.....24.3%
 Total.....100.0%

*Equivalent to 68.7% of glyphosate acid

1 pound of [this product] [GlySupreme 75-WDG] contains 0.687 pounds of glyphosate acid.

EPA Reg. No.: 80967-13

EPA Est. No.: [73432-CHN-001]

Net Contents:

Manufactured [for][by]:
 MEY Corporation
 121 S. Estes Drive, Suite 101
 Chapel Hill, NC 27514 USA

[Lot/Batch code/number]

(Note to reviewer: Lot or Batch number may appear on label or printed directly on packaging.)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See attached booklet for [additional precautionary statements] [first aid] [complete directions for use] [and warranty]

2.0 IMPORTANT PHONE NUMBERS

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE (800) 613-0427

IN CASE OF AN EMERGENCY INVOLVING THIS HERBICIDE PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL (800) 262-8200

3.0 PRECAUTIONARY STATEMENTS

| FIRST AID | |
|----------------------|--|
| If swallowed: | Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce |

Deleted: 1
 Deleted: March 19,

8/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | |
|---|--|
| | vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If inhaled: | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. |
| If in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice |
| If on skin or clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-262-8200. | |

3.1 Hazards to Humans and Domestic Animals

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: **Caution.** Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist or vapor.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE): Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

3.2 Environmental Hazards

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

(Note to Reviewer: For products in containers >50 lbs): Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

3.3 Physical or Chemical Hazards

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

3.4 Directions for Use/WPS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not allow worker entry into treated areas during the restricted entry interval (REI) of four (4) hours or until solution has dried.

Exception: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter treated area if there is no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil,

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

or water) is: coveralls, chemical resistant gloves (made of any waterproof material) and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of the product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

4.0 STORAGE AND DISPOSAL

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use to prevent spills and contamination.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal Facility.

Container Handling: Non-refillable container. Do not reuse or refill this container.

(Note to reviewer: For non-refillable plastic containers (≤50 lbs) small enough to shake):
Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable plastic containers (>50 lbs) too large to shake):
Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable paper and plastic bags, large fiber sacks):
Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable fiber drums with liners or bag-in-box packaging):
Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke. If [drum] [box] carton is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Deleted: 1
Deleted: March 19,

11/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

5.0 INFORMATION (HOW THIS PRODUCT WORKS)

Product Description: This product is a post-emergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush, and trees. It is formulated as a water soluble granule. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

Do not add surfactants, additives containing surfactants, buffering agents or pH adjusting agents to the spray solution when [this product] [GlySupreme 75-WDG] is the only pesticide used unless otherwise directed. See the "MIXING" section of this label for instructions regarding other additives.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advance to compete browning of above-ground growth and deterioration of underground plant parts.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOOD BRUSH AND TREES RATE SECTIONS" for directions for specific weeds.

Always use the higher rate of this product per acre within the directed range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the listed stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

Spray Coverage: For best results, spray coverage must be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow.

Deleted: 1
Deleted: March 19.

12/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Tank Mixing: This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly directed in this labeling. Mixing this product with herbicides or other materials may result in reduced performance.

Annual Maximum Use Rate: Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8.7 pounds of this product per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 11.6 pounds of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

6.0 MIXING

Spray solutions of this product must be mixed, stored, and applied using only stainless steel, aluminum, fiberglass, plastic, or plastic-lined steel containers. **DO NOT MIX, STORE, OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.**

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Do not apply when wind or other conditions favor drift. Hand-held applications must be properly directed to avoid spraying desirable plants.

NOTE: Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches that is not clear.

Eliminate any risk of siphoning the contents of the tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local authorities.

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

6.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows:

For hand held or backpack sprayers (less than or equal to 5 gal. capacity): Add the directed amount of this product to the spray tank. If adding Ammonium Sulfate, pre-dissolve in water before adding. Fill the spray tank with water and ensure thorough mixing. Alternatively, the

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

directed amount of this product can be mixed with water in a large container. Fill sprayer with the mixed solution.

For larger tank sprayers (greater than 5 gal. capacity): Fill the mixing or spray tank one-half full with water and start agitation. If adding Ammonium Sulfate, ensure that it is completely dissolved before proceeding. Add the directed amount of this product using a circular motion while pouring. Continue filling the spray tank with water and ensure thorough mixing.

Use caution to avoid siphoning back into the carrier source. Use approved anti-back siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by pass and return lines at the bottom of the tank and, if needed, use an approved foam or defoaming agent.

6.2 Tank Mixing Procedure

Use the following procedure to mix this product when preparing tank mixtures with other labeled products.

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If adding Ammonium Sulfate ensure that it is completely dissolved before proceeding.
4. Add [GlySupreme 75-WDG] [herbicide] using a circular motion while pouring.
5. If second product is a wettable powder, first make a slurry with the water carrier, then add the slurry SLOWLY through the screen into the tank. Continue agitation.
6. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
7. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
8. Continue filling the spray tank with water and add water soluble liquids near the end of the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift reduction additive and water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers must be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "TANK MIXING" section of "INFORMATION" for additional precautions.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

14/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Under no circumstances does MEY Corporation support a reduced rate application from suggested concentrations on this label. This includes instances where a tank mix partner is used. For all ROUNDUP crop applications, the standard use rate is 1.1 lbs. per acre. Deviation from this rate will reduce performance. If other manufacturers suggest reducing the rate to tank mix with their product or other products, MEY Corporation will not support the application.

6.3 Mixing for Hand-Held Sprayers

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table.

Amount of GlySupreme 75-WDG
% Weight to Weight

| | 0.3% | 0.6% | 0.8% | 1.1% | 2.8% | 5.5% | 11.1% |
|----------------|---------|--------|--------|--------|---------|---------|---------|
| Desired Volume | | | | | | | |
| 1 Gal | 0.4 oz | 0.8 oz | 1.1 oz | 1.5 oz | 3.7 oz | 7.3 oz | 14.8 oz |
| 3 Gal | 1.2 oz | 2.4 oz | 3.2 oz | 4.4 oz | 11.2 oz | 1.4 lb | 2.8 lb |
| 25 Gal | 10.0 oz | 1.3 lb | 1.7 lb | 2.3 lb | 5.8 lb | 11.5 lb | 23.1 lb |
| 100 Gal | 2.5 lb | 0 lb | 6.7 lb | 9.2 lb | 23.4 lb | 45.9 lb | 92.6 lb |

Above percentages are on a weight-to-weight basis with water as an 8.34 pound gallon.

For use in knapsack sprayers, it is suggested that the directed amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

6.4 Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides, on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: When using ammonium sulfate, apply this product at rates directed in this label. Lower rates will result in reduced performance.

6.5 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations. Certain blue dyes are not stable in the spray solution in the presence of this product. A jar test may be needed to determine if the desired blue dye is stable. If stability is a problem, consider switching to an alternate color dye.

6.6 Surfactants

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

15/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Although not generally required, surfactant may be added to spray solutions at water carrier volumes above 30 gallons per acre or application rates below 14 ounces of this product per acre.

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, a rate of 0.25 to 0.5 percent surfactant concentration (1 to 2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution for those surfactants containing less than 70 percent active ingredient is needed. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label booklet.

DO NOT add buffering agents or pH adjusting agents to the spray solution when GlySupreme 75-WDG is the only pesticide used. DO NOT ADD SURFACTANT TO THIS PRODUCT FOR APPLICATIONS OVER-THE-TOP OF ROUNDUP READY CROPS AND PREHARVEST COTTON.

6.7 Drift Reduction Additives

Drift reduction additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet applicator (CDA) equipment. When a drift reduction additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label. The use of drift reduction additives can affect spray coverage which may result in reduce performance.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial – Fixed Wing and Helicopter

Ground Broadcast Spray – Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Hand-Held or High-Volume Spray Equipment-Knapsack and backpack sprayers. pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-hand and motorized spray equipment used to direct the spray onto weed foliage.

*This product is not registered in California or Arizona for use in mistblowers.

Selective Equipment – Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

Injection Systems-Aerial or ground injection sprayers.

Controlled Droplet Applicator (CDA) – Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Deleted: 1
Deleted: March 19,

16/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all of these factors when making application decisions.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense sprays as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase nozzle pressure.

7.1 Aerial Equipment, Spray Drift Management, and Application Restrictions

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Use the listed rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 17 ounces per acre. Refer to the individual use area sections of this label for volumes, application rates, and further instructions.

This product plus dicamba and/or 2,4-D tank mixtures may not be applied by air in California.

Ensure uniform application – To avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards, more than 45 degrees. Where states have more stringent regulations, they must be observed.

Importance of droplet size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversions" sections of this label).

Controlling droplet size

- Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- Pressure: Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles: Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation: Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- Boom length: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.
- Application Height: Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporate and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance must increase, with increasing drift potential (higher wind, smaller droplets, etc.)

Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 miles per hour due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

Deleted: 1

Deleted: March 19,

18/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions.

Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The product must only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water.

Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

FOR AERIAL APPLICATIONS IN CALIFORNIA ONLY

Aerial applications of this product are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
2. In alfalfa and pasture renovation applications.
3. Over-the-top applications in Roundup Ready corn and cotton. Refer to further label instructions for Roundup Ready corn and Roundup Ready cotton for specific application instructions for over-the-top applications in these crops.
4. Preharvest in alfalfa, corn, cotton, wheat, Roundup Ready corn and Roundup Ready cotton. Refer to label instructions for Roundup Ready corn and Roundup Ready cotton for specific preharvest application instructions for each individual crop.

Do not plant subsequent crops other than those listed in the label for 30 days following application.

Deleted: 1
Deleted: March 19,

19/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

When applied as directed under the conditions described, this product controls annual and perennial weeds listed in the label.

When tank mixing this product with 2,4-D for aerial applications, only 2,4-D amine formulations may be used. This tank mixture may be used for fallow and reduced tillage systems and alfalfa and pasture renovation applications only.

DO NOT EXCEED MAXIMUM RATE OF 1 LBS. PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR WITH THE FOLLOWING EXCEPTIONS:
DO NOT EXCEED A MAXIMUM RATE OF 2 LBS. PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR IN FALLOW AND REDUCED TILLAGE SYSTEMS, AND PRIOR TO HARVEST IN ROUNDUP READY CORN.

FOR AERIAL APPLICATIONS IN FRESNO COUNTY CALIFORNIA ONLY

From February 15th through March 31st Only

Applicable Area

This supplement only applies to the area contained inside the following boundaries within Fresno County, California.

- North: Fresno County line
- South: Fresno County line
- East: State Highway 99
- West: Fresno County line

Information

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

Written Directions

A written direction MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written direction MUST state the proximity of surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

AERIAL APPLICATOR TRAINING AND EQUIPMENT

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Applications at Night – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

20/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

For additional information on the proper aerial application of this product, call 916-784-1718. Note: For aerial application from April 1 through February 14, refer to the "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" supplemental label.

AERIAL APPLICATIONS IN ARKANSAS ONLY

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, AS SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

7.2 Ground Broadcast Equipment

Use the listed rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume must be increased within the listed range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

7.3 Hand-Held or High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only. For listed rates and timing, refer to the "ANNUAL WEEDS—HAND-HELD OR HIGH VOLUME EQUIPMENT" section of this product label.

7.4 Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, after dilution and thorough mixing with water, to listed weeds growing in any non-crop site specified on this label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants) where any dropping or leaking will not contact crop foliage. Such equipment must be capable of preventing all crop contact with herbicide solutions and operated without leakage of spray mists or dripping onto crop. Wipers over-the-top of crops may be used only when specifically directed in this product's labeling.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applications made above desirable vegetation must be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable

Deleted: 1
Deleted: March 19,

21/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

vegetation. Droplets, mist, foam or splatter of the herbicide solution setting on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops must be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Recirculating Spray System

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

Shielded and Hooded Applicators

When applied under the conditions described in the following paragraphs for shielded and hooded applications, this product at listed rates will control those weeds listed in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION" sections of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from the spray solution. Keep shields on these sprayers adjusted to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary extend the front and rear flaps of the hoods to reach the ground in deep furrows. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods of the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure/low drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is necessary. Spray volume must be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- The spray hoods must be operated on the ground or skimming across the ground.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood must be 30 inches.
- Maximum tractor speed: 5 miles per hour to avoid bouncing of the spray hoods.
- Maximum wind speed: 10 miles per hour.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Wiper Applicators

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

When applied under the conditions described in the following paragraphs, this product CONTROLS many weeds, including volunteer corn, Texas panicum, common rye, shattercane, sicklepod, spanishneedles and bristly starbur; and SUPPRESSES many weeds including Florida beggarweed, bermudagrass, hemp dogbane, dogfennel, guineagrass, johnsongrass, milkweed, silverleaf nightshade, redroot pigweed, giant ragweed, smutgrass, sunflower, Canada thistle, musk thistle, vaseygrass, velvetleaf.

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if two applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators –Prepare solutions by dissolving 1.5 to 3.5 pounds of this product per gallon of water.

For Panel Applicators and Pressure-Feed-Systems-Prepare solutions by dissolving 1.5 to 4.0 pounds of this product per gallon of water.

Note: In preparing these concentrated solutions always allow adequate time for product to dissolve. Use of warm water will shorten dissolution time.

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. This product may be injected into the spray stream after dilution and thorough mixing with water. Do not mix this product with the concentration of other products when using injection systems.

7.6 CDA Equipment

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount listed in this label when applied by conventional broadcast equipment. For vehicle mounted CDA equipment, apply 2 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units – Apply an 11 percent solution of this product (13.7 oz of product per gallon) at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 miles per hour (1 quart per acre). For the control of perennial weeds,

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

apply an 11 to 22 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 miles per hour (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage of any other green tissue of desirable vegetation, as damage or destruction may result.

8.0 ANNUAL AND PERENNIAL CROPS (Alphabetical)

NOTE: THIS SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED CROPS WITHIN SECTION 8 GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the "ROUNDUP READY CROPS" section of this label for instructions for treating Roundup Ready crops.

TYPES OF APPLICATIONS:

Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, At-Planting, Hooded Sprayers in Row Middles, Shielded Sprayers in Row Middles, Wiper Applications in Row Middles, and Post-Harvest treatments.

USE INSTRUCTIONS:

Apply this product during fallow intervals preceding planting, prior to planting or transplanting, at-planting or preemergent to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be according to the rates listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", AND "WOODY BRUSH AND TREES RATE SECTIONS" in this label. Repeat applications may be made up to a maximum of 8.7 pounds per acre per year.

Post-directed hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted below, wipers may also be used above certain crops to control tall weeds. Refer to the "SELECTIVE EQUIPMENT" section of this label for essential precautions when using hooded sprayers or wipers to avoid crop injury caused by leakage of spray mists or dripping onto crops. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

RESTRICTIONS:

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast

Deleted: 1
Deleted: March 19.

24/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

applications made at emergence will result in injury or death to emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.

For broadcast postemergent treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

8.1 Cereal and Grain Crops

LABELED CROPS: Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Rye Quinoa, Teff, Teosinte, Triticale, Wheat (all types), Wild rice.

RESTRICTIONS: Do not treat rice fields or leaves when field contains water.

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following Red Rice Control Prior to Planting Rice, Spot Treatment (except Rice) Over-the-Top Wiper Applications (Feed Barley and Wheat only), Preharvest (Feed Barley and Wheat only).

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting of cereal crops. Applications must be made prior to emergence of the crop.

Red Rice Control Prior to Planting Rice

USE INSTRUCTIONS: Apply 26 ounces of this product in 5 to 10 gallons of water per acre. Flush field prior to application to obtain uniform germination and stand of red rice. Make application when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

RESTRICTIONS: Avoid spraying during low humidity conditions, as reduced control may result. Do not treat rice fields or levees when the fields contain floodwater. Do not re-flood treated fields for 8 days following application.

Over-the-Top Wiper Applications (Feed Barley and Wheat Only)

USE INSTRUCTIONS: Wiper applications may be used in wheat and feed barley. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth, and when the rye is at least 6 inches above the wheat crop.

RESTRICTIONS: Allow at least 35 days between application and harvest. Do not use roller applicators.

Preharvest (Feed Barley and Wheat Only)

Deleted: 1

Deleted: March 19,

25/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of wheat or feed barley. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard-dough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest.

This product may be applied using either aerial or ground spray equipment. For ground applications, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre.

RESTRICTIONS: Do not apply more than 17 ounces of this product per acre. Allow 7 days between application and harvest or grazing. Do not apply preharvest to wheat or barley grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of cereal crops. Higher rates be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

8.2 Corn

TYPES OF CORN: Field corn, Seed corn, Silage corn, Sweet corn and Popcorn.

TYPES OF APPLICATONS: Those listed in Section 8.0 plus the following: Spot Treatment, Preharvest.

For Roundup Ready corn, see the "ROUNDUP READY CROPS" section of this label.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank mixture before, during or after planting corn. Applications must be made prior to emergence of the crop.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

| | | |
|------------------|---------------------|-------------------|
| 2,4-D | Degree XTRA® | Harness XTRA 5.6L |
| Aim | Distinct™ | Lariar® |
| Atrazine | Dual Magnum™ | Lasso®/Alachlor |
| Axiom™ | Dual II Magnum™ | Linex™/Lorox |
| Balance Pro™ | Epic™ | Marksman™ |
| Banvel™/Clarity™ | Frontier™/Outlook™ | Micro-Tech® |
| Bicep li Magnum™ | Fultime™ | Prowl™ |
| Butler | Guardsman™/Leadoff™ | Python™ |
| Degree® | Harness® | Simazine |
| | Harness XTRA | Topnotch™ |
| | | Resource® |

For difficult-to-control annual weeds including fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

to 6 inches tall, apply this product at 17 ounces per acre in these tank mixtures. For other labeled annual weeds, apply 13 to 17 ounces of this product per acre when weeds are less than 6 inches tall, and 17 to 26 ounces when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

RESTRICTIONS: Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses including barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. The area covered by this instruction includes Route 50 South in Illinois and Indiana and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Corn must be at least 12 inches tall, measured without extending leaves. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. To the extent consistent with applicable law such damage shall be the sole responsibility of the applicator. Do not apply more than 17 ounces of this product per acre for each application and no more than 52 ounces per acre per year for hooded sprayer applications.

Spot Treatment

USE INSTRUCTIONS: For spot treatments, apply this product prior to silking of corn.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

Preharvest

USE INSTRUCTIONS: Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). For ground applications, apply up to 52 ounces of this product per acre. For aerial applications, apply up to 35 ounces of this product per acre.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest. Do not apply preharvest to corn grown for seed, as a reduction in germination or vigor may occur.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

Deleted: 1
Deleted: March 19.

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

8.3 Cotton

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following: Selective Equipment, Spot Treatment, Preharvest.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton. Applications must be made prior to emergence of the crop.

Hooded Sprayer, Selective Equipment

USE INSTRUCTIONS: This product may be applied through hooded sprayers, recirculating sprayers, shielded applicators or wiper applicators in cotton. Allow at least 7 days between application and harvest.

RESTRICTIONS: See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for information on proper use and calibration of this equipment.

Spot Treatment

USE INSTRUCTIONS: For spot treatments, apply this product prior to boil opening of cotton.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift spray outside target area for the same reason.

Preharvest

USE INSTRUCTIONS: This product provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "ANNUAL WEEDS", "PERENNIAL WEEDS" and "WOODY BRUSH AND TREES RATE SECTIONS" sections of this label. For cotton regrowth inhibition, apply 8.7 to 35 ounces of this product per acre.

Up to 35 ounces of this product may be applied using either aerial or ground spray equipment. Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

TANK MIXTURES: This product may be tank mixed with DEF™ 6, Folex™, Ginstar, or Prep™ to provide additional enhancement of cotton leaf drop.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of cotton. Do not apply preharvest to cotton grown for seed, as a reduction in germination or vigor may occur.

8.4 Fallow Systems

LABELED CROPS: This product may be applied during the fallow period prior to planting or emergence of any crop on this label.

Deleted: 1
Deleted: March 19,

28/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Aid-to-Tillage.

Chemical Fallow

USE INSTRUCTIONS: This product may be applied during the fallow period prior to planting or emergence of any crop listed on this label. This product may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures with 2,4-D and dicamba may be used. Applications up to 35 ounces per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops.

RESTRICTIONS: For any crop not listed on this label, applications must be made at least 30 days prior to planting. Do not apply dicamba tank mixtures by air in California.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

Preplant Fallow Beds

USE INSTRUCTIONS: This product may be applied to fallow beds prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 30 days prior to planting. This product will control weeds listed in the "ANNUAL WEEDS", "PERENNIAL WEEDS", and "WOODY BRUSH AND TREES RATE SECTIONS" sections of this label.

TANK MIXTURES: In addition, 6.5 ounces of this product plus 2 to 3 fluid ounces of Goal™ 2XL per acre will control the following weeds with the maximum height or length indicated: 3 inches – common cheeseweed, chickweed, groundsel; 6 inches – London rocket, shepherd's purse.

8.2 ounces of this product plus 2 to 3 fluid ounces of Goal 2XL per acre will control the following weeds with the maximum height or length indicated: 6 inches – common cheeseweed, groundsel, marestail (*Conya Canadensis*); 12 inches – chickweed, London rocket, shepherd's purse.

Aid-to-Tillage

USE INSTRUCTIONS: This product may be used in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 6.5 ounces of this product in 3 to 10 gallons of water per acre. Make application before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

RESTRICTIONS: Tank mixtures with residual herbicides may result in reduced performance.

8.5 Grain Sorghum (Milo)

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following: Spot Treatment, Over-the-Top Wiper Applications, Preharvest.

Deleted: 1
Deleted: March 19,

29/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in tank-mixture before, during or after planting grain sorghum. Applications must be made prior to emergence of the crop.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

| | |
|-----------------|------------|
| Aztazome | Lariat |
| Bicep II Magnum | Lasso |
| Bullet | Micro-Tech |
| Dual II Magnum | |

For difficult-to-control annual weeds including fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 17 ounces per acre in these tank mixtures. For other labeled annual weeds, apply 13 to 17 ounces of this product per acre when weeds are less than 6 inches tall, and 17 to 26 ounces when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased or acceptable weed control.

Spot Treatment, Over-the-Top Wiper Applications

USE INSTRUCTIONS: This product may be applied as a spot treatment in grain sorghum. Make spot treatments before heading of milo. This product may be applied with wiper applicators to control or suppress the weeds listed under "Wiper Applicators" in the "SELECTIVE EQUIPMENT" section of this label.

RESTRICTIONS: For spot treatment, do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

For wiper applicators, allow at least 40 days between application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

RESTRICTIONS: Milo must be at least 12 inches tall, measured without extending leaves. Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. To the extent consistent with applicable law such damage shall be the sole responsibility of the applicator. Do not graze or feed milo forage or fodder following applications of this product through hooded

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

sprayers. Do not apply more than 17 ounces of this product per acre per application and no more than 52 ounces per acre per year for hooded sprayer applications.

Preharvest

USE INSTRUCTIONS: Make applications at 30 percent grain moisture or less.

RESTRICTIONS: Do not apply more than 35 ounces of this product per acre. As with other herbicides that cause sudden plant death, avoid preharvest applications of this product to milo infected with charcoal rot as lodging can occur. Allow a minimum of 7 days between application and harvest of sorghum. Do not apply preharvest to sorghum grown for seed, as a reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of grain sorghum. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 17 ounces of this product per acre for control, or 8.7 ounces of this product per acre for suppression.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

8.6 Herb and Spices

LABELED CROPS: Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Camomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Culantro (leaf), Culantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following: Over-the-Top Wiper Applications (Peppermint and Spearmint only), Spot Treatment (Peppermint and Spearmint only).

RESTRICTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler system.

Over-the-top Wiper Applications, Spot Treatment (Peppermint and Spearmint only)

Deleted: 1
Deleted: March 19,

31/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

USE INSTRUCTIONS: This product may be used as a spot treatment or wiper application in spearmint and peppermint. Apply spot treatments on a spray-to-wet basis with hand-held equipment, including back-pack and knapsacks sprayers, pump-up pressure sprayers, hand-guns, hand-wands or any other hand-held or motorized spray equipment used to direct the spray solutions to a limited area. In wiper applications, the applicator must be adjusted so that the wiper contact point is at least 2 inches above the crop. Weeds must be a minimum of 6 inches taller than the crop.

RESTRICTIONS: Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30 day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested must be treated at one time. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for this reason. In wiper applications, contact of the herbicide solution with the crop may result in damage or destruction.

8.7 Oil Seed Crops

LABELED CROPS: Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower. For Roundup Ready canola, see the "ROUNDUP READY CROPS" section of this label.

TYPES OF APPLICATIONS: Those listed in Section 8.0.

USE INSTRUCTIONS: This product may be applied before, during or after planting oil seed crops. Broadcast applications must be made prior to emergence of the listed oil seed crops. Wiper applicators or hooded sprayers may be used between the rows once the crop is established.

TANK MIXTURES: For sunflowers, a tank mixture with Prowl may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue.

RESTRICTIONS: Do not apply more than 35 ounces of this product per acre on canola. Do not apply more than 17 ounces of this product per acre for sunflowers as a single preplant or preemergent application per year. Do not feed or graze sunflower forage following application of this product. Do not apply more than 53 ounces per acre of this product on Safflower for all preemergence, shielded spray and preharvest applications.

Preplant, At Planting, Preemergence

USE INSTRUCTIONS: This product may be applied before, during or after planting oil seed crops listed in this section. Observe the maximum application rates for canola, safflower and sunflower listed at the beginning of this section.

TANK MIXTURES: For sunflowers, a tank mixture with pendimethalin (Prowl) may be applied before, during or after planting into conventionally tilled soil, a cover crop, established sod or previous crop residue.

Selective Equipment

USE INSTRUCTIONS: This product may be applied using a wiper applicator or shielded sprayer in between the rows once the crop is established. See additional instructions on the use of wiper applicators and hooded sprayers in the "APPLICATION EQUIPMENT AND

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TECHNIQUES" section of this label. Observe the maximum application rates for canola, safflower and sunflower listed at the beginning of this section.

Preharvest (Safflower and Sunflower Only)

USE INSTRUCTIONS: This product provides weed control as a harvest aid when applied to a physiologically mature crop of safflower and sunflower prior to harvest. For safflower, apply a maximum of 53 oz of this product when seed has lost its opaque character, approximately 20 to 30 days after the end of flowering of the secondary branches. For sunflower, apply a maximum of 17 oz of this product when the backsides of sunflower heads are yellow and bracts are turning brown, and seed moisture content is less than 35 percent.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or livestock feeding. Application must be made at least 30 days prior to planting any crop not listed on this label.

Post-Harvest

USE INSTRUCTIONS: This product may be applied for weed control after harvest of oil seed crops. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation. Applications must be made at least 30 days prior to planting any crop not listed on this label.

8.8 Soybeans

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following: Spot Treatment, Preharvest, Selective Equipment.

For Roundup Ready soybeans, see the "ROUNDUP READY CROPS" section of this label.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting soybeans. Applications must be made prior to emergence of the crop.

Selective Equipment

USE INSTRUCTIONS: This product may be applied in soybean using a shielded applicator, hooded sprayer, wiper applicator or sponge bar. See additional instructions on the use of selective equipment in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Allow at least 7 days between application and harvest.

Preharvest

USE INSTRUCTIONS: This product may be applied to soybean prior to harvest after pods have set and lost all green color. Apply at rates given in the "ANNUAL WEEDS RATE SECTION" and "PERENNIAL WEEDS RATE SECTION". Take care to avoid excessive seed shatter loss due to ground application equipment.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: Do not apply more than 5.4 lbs of this product per acre for preharvest application. Do not apply more than 35 oz of this product per acre by air. Allow a minimum of 7 days between application and harvest of soybeans. If the preharvest application rate is greater than 17 oz. per acre, do not graze or harvest treated hay or fodder for livestock feed within 25 days of application. If the application rate is 17 oz. per acre or less, the grazing restriction is reduced to 14 days after application. Preharvest application on soybean grown for seed could result in a reduction in germination or vigor.

TANK MIXTURES: Apply these tank mixtures in 10 to 20 gallons of water per acre.

| | | |
|---------------|--------------------|-----------------|
| Aim™ | Dual II Magnum | Micro-Tech |
| Amplify™ | Firstate™ | Prowl |
| Assure li™ | Flexstar™ | Pursuit™ |
| Authority™ | Frontier™/Outlook™ | Pursuit Plus |
| Boundary™ | Fusion™ | Reflex™ |
| Canopy™ | Gauntlet™ | Scepter™ |
| Canopy XL™ | Lasso | Sencor™/Lexone™ |
| Command™ | Linex™ | Squadron™ |
| Command XTRA™ | Lorox/Linuron | Steel™ |
| Domain™ | Lorox Plus™ | Valor |
| Dual Magnum | | Resource® |

This product may be tank-mixed with 2,4-D or 2,4-DB. See the 2,4-D label for intervals between application and planting.

For difficult-to-control annual weeds including fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 17 ounces per acre in these tank mixtures. For other labeled annual weeds, apply 13 to 17 ounces of this product per acre when weeds are less than 6 inches tall, and 17 to 26 ounces when weeds are over 6 inches tall.

RESTRICTIONS: Tank-mixtures with some of the above listed herbicides may result in reduced weed control due to antagonism. Read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental labeling or fact sheets published separately for all herbicides used. Use according to the most restrictive directions for each product in the mixture.

Spot Treatment

USE INSTRUCTIONS: For spot treatment, apply this product prior to initial pod set in soybeans

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

8.9 Sugar Cane

TYPES OF APPLICATION: Those listed in Section 8.0 plus Spot Treatment, Preplant, At Planting, Preemergence

USE INSTRUCTIONS: This product may be applied in or around sugarcane fields, or in fields prior to the emergence of plant cane.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

34/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

Spot Treatment

USE INSTRUCTIONS: This product may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, apply a 1-percent solution of this product in water using a spray-to-wet technique. Best results are obtained on volunteer or diseased sugarcane with at least 7 new leaves.

RESTRICTIONS: Avoid spray contact with healthy sugarcane plants as severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

Fallow Treatment

USE INSTRUCTIONS: This product may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. This product may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4.3 to 5.4 lbs of this product in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Aerial application of up to 52 oz per acre may be made in fallow sites where there is sufficient buffer to prevent drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used. Ensure that the specific product being used is labeled for these applications in sugarcane. Read and follow label directions of all products in the tank mixture.

Hooded Sprayers

USE INSTRUCTIONS: This product may be used through hooded sprayers for weed control between the rows of sugarcane. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional use instructions.

RESTRICTIONS: Do not allow treated weeds to come into contact with the crop. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

8.10 Vegetable Crops

NOTE: THIS "VEGETABLE CROPS" SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED VEGETABLE CROPS WITHIN SECTION 8.10 GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, Prior to Transplanting Vegetables, At-Planting, Hooded Sprayers in Row Middles, Shielded Sprayers in Row Middles, Wiper Applications in Row Middles, and Post-Harvest, Directed Applications (Nonbearing Ginseng), Over-the-Top Wipers (Rutabagas Only).

RESTRICTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler system. Care must be taken to

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

insure that the wash water flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at planting applications, applications must be before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles must be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

8.10.1 Brassica Vegetables

LABELED CROPS: Broccoli, Chinese broccoli (gai lon), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

8.10.2 Bulb Vegetables

LABELED CROPS: Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

8.10.3 Cucurbit Vegetables and Fruits

LABELED CROPS: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Melons (all), Momordica spp (includes balsam apple, balsam pear bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon). Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini). Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash). Watermelon.

RESTRICTIONS: For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Cherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

8.10.4 Leafy Vegetables

LABELED CROPS: Amaranth (Chinese spinach), Arugula (roquette), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

Deleted: 1
Deleted: March 19,

36/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: For Watercress, avoid applications within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of injury.

8.10.5 Fruiting Vegetables

LABELED CROPS: Eggplant, Groundcherry (*Physalis* spp), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato

RESTRICTIONS: For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato, do not make hooded or shielded sprayer applications in row middles because of the potential for crop injury.

8.10.6 Legume Vegetables (Succulent or Dried)

LABELED CROPS: Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin) Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (*fava*), Chickpea (*garbanzo*), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea field pea, garden pea, green pea, snow pea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

8.10.7 Root and Tuber Vegetables

LABELED CROPS: Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish Salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

Directed Applications (Nonbearing Ginseng Only)

USE INSTRUCTIONS: This product may be used for weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high volume wands, lances, and orchard guns or with wiper application equipment.

RESTRICTIONS: Direct applications so that there is no contact of this product with the ginseng plant. Applications must be made at least one year prior to harvest.

Over-the-Top Wiper Applications (Rutabagas Only)

USE INSTRUCTIONS: Wiper applicators may be used over-the-top of rutabagas.

RESTRICTIONS: Allow at least 14 days between application and harvest of rutabagas.

8.11 Miscellaneous Crops

LABELED CROPS: Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Those listed in Section 8.0 plus the following Weed Control, Site Preparation, Spot Treatment (Asparagus).

For Roundup Ready sugar beets, see the "ROUNDUP READY CROPS" section of this label.

RESTRICTIONS: Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch) or fruit of crops because severe injury or destruction may result. When making preemergence and at planting applications, applications must be made before crop emergence to avoid severe crop injury. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. In crops with vines, hooded sprayer, shielded sprayer and wiper applications to row middles must be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in this product's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional information.

Weed Control, Site Preparation

USE INSTRUCTIONS: This product may be applied for weed control or for site preparation prior to planting or transplanting crops listed in this section.

RESTRICTIONS: When applying this product prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of this product, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5 inch application of water, either by natural rainfall or via a sprinkler system. Care must be taken to insure that the wash water flushes off the plastic mulch and does not enter transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Do not apply within a week before the first asparagus spears emerge. Do not feed or graze treated pineapple forage following application.

Spot Treatment (Asparagus)

USE INSTRUCTIONS: This product may be applied immediately after cutting, but prior to the emergence of new spears.

RESTRICTIONS: Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

Post-Harvest (Asparagus)

USE INSTRUCTIONS: This product may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments must be applied as a directed or shielded spray in order to avoid contact of the spray with ferns, stems or spears.

RESTRICTIONS: Direct contact of the spray with the asparagus may result in serious crop injury. Select and use listed types of spray equipment for postemergence post-harvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

9.0 TREE, VINE, AND SHRUB CROPS (Alphabetical)

NOTE: THIS SECTION GIVES DIRECTIONS THAT APPLY TO ALL LISTED TREE, VINE, AND SHRUB CROPS WITHIN SECTION 9 GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGOREIS FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

TYPES OF APPLICATIONS: Preplant (Site Preparation) Broadcast Sprays, Weed Control, Middles (between rows of trees, vines or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (shielded sprayers, wiper treatments), Directed Sprays, Spot Treatment, Perennial Grass Suppression, Cut Stump.

Applications may be made with boom equipment, CDA equipment, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

USE INSTRUCTIONS: This product may be applied in middles (between rows of trees or vines), strips (within rows of trees or vines), and for weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 8.7 ounces to 5.5 pounds per acre according to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE SECTIONS" sections of this label. Utilize rates at the higher end of the rate range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 11.6 pounds per acre per year.

The maximum use rates stated throughout this product's labeling applying to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

RESTRICTIONS: Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of this product with other than matured brown bark can result in serious crop damage or destruction. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) must be used to minimize the potential for leakage or drift of herbicide sprays onto crops. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with crop may be used. See "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for additional directions and precautions.

Allow a minimum of 3 days between application and transplanting.

Middles (between rows)

USE INSTRUCTIONS: This product will control or suppress annual and perennial seeds and ground covers growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

Deleted: 1
Deleted: March 19.

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TANK MIXTURES: A tank mixture of this product plus Goal 2XL may be used for annual weeds in middles between rows of citrus crops, tree fruits, tree nuts and vine crops. Use this mixture when weeds are stressed or growing in dense population. 8.7 to 17 ounces per acre of this product plus 3 to 12 fluid ounces per acre of Goal 2XL will control annual weeds with a maximum height or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's purse, annual sowthistle, filaree (suppression), horseweed/marestail, stinging nettle and common purslane (suppression). 8.7 to 17 ounces per acre of this product plus 3 to 12 fluid ounces per acre of Goal 2XL will control common cheeseweed (malva) or hairy fleabane with a maximum height or diameter of 3 inches.

Strips (in rows)

TANK MIXTURES: This product may be applied in rows of tree or vine crops in tank mixtures with the following products:

- | | |
|--------------------|--------------|
| Devrinol™ 50 DF | Simazine 4L |
| Direx™ 4L | Simazine 80W |
| Goal 2XL | Sim-Trol™ 4L |
| Karmex DF | Solicam™DF |
| Krovar I | Surflan™ AS |
| Prowl | Surflan 75W |
| Princep Caliber™90 | |

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Perennial Grass Suppression

This product will suppress perennial grasses including bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 4.3 ounces of this product in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 3.2 ounces of this product per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply this product 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 3.2 ounces of this product in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 2.8 ounces of this product per acre, followed by an application of 1.9 to 2.8 ounces per acre about 45 days later. Make no more than 2 applications per year.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

For burndown of Bermudagrass, apply 17 to 35 ounces of this product in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 3.2 to 8.7 ounces of this product per acre east of the Rocky Mountains and 8.7 ounces of this product per acre west of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full green-up. If the Bermudagrass is mowed prior to application, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 3.2 to 6.5 ounces of this product per acre must be used in shaded conditions or where a lesser degree of suppression is desired.

Cut Stump

USE INSTRUCTIONS: Cut stump applications of this product may be made during site preparation or site renovation, prior to transplanting tree crops. This product will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

Citrus Trees: Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo, Tangor

Fruit Trees: Apple, Apricot, Cherry (sweet sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince.

Nut Trees: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).

Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a solution prepared by dissolving 2.1 to 3.7 pounds of this product per gallon of water to the freshly cut surface immediately after cutting. Delays in applications may result in reduced performance. For best results, applications must be made during periods of active growth and full leaf expansion.

RESTRICTIONS. DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

9.1 Berry Crops

LABELLED CROPS: Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry Dirksen thornless berry, Himalaya berry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry). Blueberry, Cranberry, Currant, Elderberry Gooseberry, Huckleberry, Loganberry, Raspberry (black, red), Salal.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus Spot Treatment in Cranberry Production and Post-Harvest Treatments in Cranberry Production.

RESTRICTIONS: To avoid damage, herbicide sprays must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest in cranberries. Allow a minimum of 14 days between last application and harvest in other berry crops. Do not make directed sprays within the cranberry bush areas prior to berry harvest.

Spot Treatment in Cranberry Production

USE INSTRUCTIONS: Spot treatments may be used to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in this label may be used. Drop water level to remove standing water in ditches prior to application. In hand-held sprayers, use 0.7 to 1 percent solution of this product. Spray to wet vegetation, not to run-off.

RESTRICTIONS: For treatments after draw down of water in dry ditches, allow 2 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after draw down to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not apply this material through the irrigation system. Do not make applications by air. Do not apply directly to water. Use nozzles that emit medium to large-sized droplets to minimize drift in order to avoid crop injury.

Post-Harvest Treatments in Cranberry Production

USE INSTRUCTIONS: Application of this product may be made after the harvest of cranberries to control weeds growing within the field. Best results will be obtained if applications are made to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed under "APPLICATION EQUIPMENT AND TECHNIQUES" in this label may be used. If using hand-held sprayers, use 0.3 to 0.6 percent solution of this product. Spray to wet vegetation, not to run-off. If using hand-held boom sprayers, apply 35 to 70 ounces of this product per acre.

RESTRICTIONS: Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not apply this product through the irrigation system. Do not make applications by air. Do not apply directly to water. Even though vines appear dormant, contact of the herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

9.2 Citrus

LABELLED CROPS: Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

USE INSTRUCTIONS (The instructions below pertain to applications in Florida and Texas): For burndown or control of the weeds listed below, apply the listed rates of this product in 3

Deleted: 1
Deleted: March 19,

42/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 35 to 52 ounces of this product per acre. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 35 ounces per acre when plants are less than 8 inches tall and 52 ounces per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Krovar™ I or Karmex™ may improve control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

Perennial weeds:

S = Suppression
PC = Partial control
B = Burndown
C = Control

| Weed SPECIES | GlySupreme 75-WDG RATE PER ACRE | | | |
|-------------------------|---------------------------------|----------|----------|---------|
| | 1.1 LBS | 2.1 LBS. | 3.2 LBS. | 4.4 LBS |
| Bermudagrass | B | -- | PC | C |
| Guineagrass | B | C | C | C |
| Texas and Florida Ridge | -- | B | C | C |
| Paragrass | B | C | C | C |
| Torpedograss | S | S | PC | C |

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only.

9.3 Miscellaneous Tree Food Crops

LABELED CROPS: Cactus (fruit and pads), Palm (heart, leaves), Palm (oil).

TYPES OF APPLICATIONS: Those listed in Section 9.0.

9.4 Non-Food Tree Crops

LABELED CROPS: PINE, Poplar, Eucalyptus, Christmas trees, Other non-food tree crops.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

Directed Sprays, Spot Treatment, Wipers

USE INSTRUCTIONS: This product may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

RESTRICTIONS: Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. UNLESS OTHERWISE DIRECTED, THIS PRODUCT IS NOT FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN CHRISTMAS TREES AND OTHER PINE TREES.

Deleted: 1
Deleted: March 19.

43/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

USE INSTRUCTIONS: This product may be used prior to planting labeled crops listed in this section.

RESTRICTIONS: Precautions must be taken to protect nontarget plants during site preparation applications.

9.5 Pome Fruit

LABELED CROPS: Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince.

TYPES OF APPLICATIONS: Those listed in Section 9.0

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in pome crops.

9.6 Stone Fruit

LABELED CROPS: Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), Plumcot.

TYPES OF APPLICATIONS: Those listed in Section 9.0.

RESTRICTIONS: Allow a minimum of 17 days between last application and harvest in stone fruit crops. For olive groves, apply as directed sprays only.

Restrictions on Application Equipment

For cherries, any application equipment listed in Section 9.0 may be used in all states.

Any application equipment listed in Section 9.0 may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states, use wiper equipment only.

For PEACHES grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. **EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.**

9.7 Tree Nuts

LABELED CROPS: Almond, Beechnut, Betelnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (black, English).

TYPES OF APPLICATIONS: Those listed in Section 9.0

Deleted: 1
Deleted: March 19,

44/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut.

9.8 Tropical and Subtropical Trees and Fruits

LABELED CROPS: Ambarella, Atemoya, Avocado, Banana, Barbados cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates, Durian, Feijoa, Figs, Governor's plum, Guava, Ilama, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamy apple, Mango, Mangosteen Marmaladebox (genip), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Surinam cherry, Tamarind Tea, Ti (roots and leaves). Wax jambu.

TYPES OF APPLICATIONS: Those listed in Section 9.0 plus Bananacide (Banana Only).

RESTRICTIONS: Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

Bananacide (Banana only)

USE INSTRUCTIONS: This product may be used to destroy banana plants infected with the Banana Bunchy Top Virus as well as non-infected banana plants to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 0.03 ounces (0.8 grams) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved using 0.07 fluid ounces (2 mL) of a solution prepared by dissolving 2.1 to 3.7 pounds of this product per one gallon of water either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. Make the injection at least one foot above the ground, except for very small plants, which must be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be mechanically destroyed. For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

RESTRICTIONS: Do not apply more than 0.5 ounces (13 grams) of this product per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Following transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying this product for weed control.

9.9 Vine Crops

LABELED CROPS: Grapes (raisin, table, wine), Hops, Kiwi, Passion fruit.

Deleted: 1
Deleted: March 19,

45/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Those listed in Section 9.0

USE INSTRUCTIONS: Applications must not be made when green shoots, canes or foliage are in the spray zone.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury, or make applications with shielded sprayers or wiper equipment.

RESTRICTIONS: Allow a minimum of 14 days between last application and harvest in vine crops. Do not use selective equipment in kiwi.

10.0 PASTURE GRASSES, FORAGE LEGUMES AND RANGELANDS

10.1 Alfalfa, Clover, and other Forage Legumes

LABELLED CROPS: Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types).

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Spot Treatment, Over-the-Top Wiper Applications, Renovation, Preharvest (except kenaf and leucaena).

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting crops listed in this section. Applications must be made prior to emergence of the crop.

RESTRICTIONS: Remove domestic livestock before application. The crop may be fed or grazed as soon as it reaches sufficient maturity.

Preharvest (except Kenaf and Leucaena)

USE INSTRUCTIONS: This product may be used in declining stands or any stand where severe crop injury or destruction is acceptable. This product will control annual and perennial weeds, including quackgrass, when applied prior to crop harvest. Applications may be made at any time of the year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

RESTRICTIONS:

Make only one application to an existing crop stand per year. The treated crop and weeds can be harvested and fed to livestock according to the intervals below.

| | Maximum Single Application Rate | Minimum Interval Between application and harvest/grazing |
|---------------------------------|---------------------------------|--|
| Alfalfa | 2.2 pounds per acre | 36 hours |
| All other labeled Legumes above | 1.6 pounds per acre | 3 days |

This application may destroy an alfalfa stand and may severely injure or destroy other labeled crops such as clover. Do not apply preharvest to alfalfa grown for seed, as a reduction in germination or vigor may occur.

Deleted: 1

Deleted: March 19,

46/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Spot Treatment, Over-the-Top Wiper Applications

USE INSTRUCTIONS: This product may be applied as a spot treatment or with wiper applicators. For wipers, see the "Wiper Applicators" in the "SELECTIVE EQUIPMENT" section of this label. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field are must be treated at one time. Remove domestic livestock before application and wait 3 days after application before grazing livestock or harvesting.

Renovation

USE INSTRUCTIONS: This product may be applied as a broadcast spray to renovate existing stands of alfalfa, clover, and other labeled forage legumes. If the crop is to be grazed or harvested for feed, use up to 2.2 pounds per acre in alfalfa and up to 1.6 pounds per acre in other labeled legumes. For complete removal of established stands of clover, it may be necessary to use the higher treatment rates listed in the "PERENNIAL WEEDS RATE SECTION" in this label.

RESTRICTIONS: When treatment rates of 2.2 pounds per acre for alfalfa or 1.6 pounds per acre for other forage legumes are used, remove domestic livestock before application and wait 3 days after application before reintroduction. If treatment rates above these levels are necessary, do not graze or harvest treated foliage for livestock feed. Crops listed for treatment in this label may be planted into the treated area at any time; for other crops, wait 30 days between application and planting.

Dormant Alfalfa Use

INSTRUCTIONS

This product will control or suppress many weeds, including quackgrass, downy brome and cheatgrass in dormant alfalfa.

Apply 4.8 to 6.5 ounces per acre of this product. Apply in the spring to alfalfa that is dormant. Applications must be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliolate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliolate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa. Do not use this product where a slight yield reduction in the first cutting of alfalfa cannot be tolerated. Do not make more than one application per year. Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist soil conditions as effects of this product wear off.

Application of this product is limited to persons who have attended a Monsanto-approved training program. Application of this product can cause crop injury. Any crop injury is the sole responsibility of the applicator.

10.2 Conservation Reserve Program (CRP)

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Renovation (rotating out of CRP), Site Preparation, Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Applications.

Renovation (Rotation out of CRP), Site Preparation

USE INSTRUCTIONS: This product may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation directions. For any crop not listed for treatment in this label, applications must be made at least 30 days prior to planting.

Postemergence Weed Control in Dormant CRP Grasses, Over-the-Top Wiper Applications

USE INSTRUCTIONS: This product may be used to suppress competitive growth and seed production of undesirable vegetation in CRP acres. Such applications may be made with wiper application equipment or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 6.5 to 8.7 ounces of this product per acre in early spring before desirable CRP grasses, including crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

RESTRICTIONS: Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 3.3 pounds per acre per year onto CRP grasses.

10.3 Grass Seed or Sod Production

LABELED CROPS: Any grass (Gramineae family) except Corn, Sorghum, Sugarcane and those listed in this label under "CEREAL AND GRAIN CROPS".

TYPES OF APPLICATIONS: Preplant, Preemergence, At-Planting, Renovation, Removal of Established Stands, Site Preparation, Shielded Sprayers, Over-the-Top Wiper Applications, Spot Treatments, Creating Rows in Annual Ryegrass.

Preplant, Preemergence, At-Planting, Renovation, Removal of Established Stands, Site Preparation

USE INSTRUCTIONS: This product controls most existing vegetation prior to renovating turf or forage grass seed areas or establishing turf grass grown for sod. It may also be used to destroy remaining undesired grass vegetation when production fields are converted to alternate species or crops. Make applications before, during, or after planting or for renovation. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm season grasses, including Bermudagrass, summer or fall applications provide best control. Broadcast equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. Application rates up to 5.5 pounds per acre may be used to totally remove established stands of tough to kill grass species.

RESTRICTIONS: Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques including vertical mowing, coring or slicing must be delayed for 7 days after application to allow proper translocation into underground plant parts. If application rates total 3.3 pounds per acre or less, no waiting period between treatment and feeding or

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

livestock grazing is required. If the rate is greater than 3.3 pounds per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. For any crop not listed for treatment in this label, applications must be made at least 30 days prior to planting. Applications must be made prior to the emergence of the crop to avoid crop injury.

Shielded Sprayers

USE INSTRUCTIONS: Apply 1.1 to 3.3 pounds of this product in 10 to 20 gallons of water per acre to control weeds between grass seed rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by the protective shields. For additional instructions, see "Shielded Applicators" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. To the extent consistent with applicable law such damage shall be the sole responsibility of the applicator.

Over-the-Top Wiper Applications

USE INSTRUCTIONS: Applicators must be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Weeds must be a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when height of weeds varies so that no all weeds are contacted. In these instances, repeat treatments may be necessary. For additional instructions, see "Wiper Applicators" in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

RESTRICTIONS: Contact of the herbicide solution with desirable vegetation may result in damage or destruction.

Spot Treatments

USE INSTRUCTIONS: Use a 1.0- to 2.0-percent solution.

RESTRICTIONS: Apply this product prior to heading of grasses grown for seed. The crop receiving the spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason. Hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Creating Rows in Annual Ryegrass

USE INSTRUCTIONS: Use 0.6 to 1.1 pounds of this product per acre. Use the higher rate when the ryegrass is greater than 6 inches tall. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height.

RESTRICTIONS: Set nozzle heights to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use low pressure nozzles, or drop nozzles designed to target the application over a narrow band.

To the extent consistent with applicable law grower assumes all responsibility for crop losses from misapplication.

Deleted: 1
Deleted: March 19,

49/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

10.4 Pastures

LABELLED CROPS: Any grass (Gramineae family) except Corn, Sorghum, Sugarcane and those listed in this label under "CEREAL AND GRAIN CROPS". Grasses that may be treated include Bahiagrass, Bermudagrass, Bluegrass Brome, Fescue, Guineagrass, Kikuya grass, Orchard grass, Pangola grass, Ryegrass, Timothy, Wheatgrass.

TYPES OF APPLICATIONS: Preplant, Preemergence, Spot Treatment, Over-the-Top Wiper Applications, Pasture renovation, Postemergent Weed Control (Broadcast Treatment).

Preplant, Preemergence, Pasture Renovation

USE INSTRUCTIONS: This product may be applied prior to planting or emergence of forage grasses. In addition, this product may be used to control perennial pasture species listed on this label prior to re-planting.

RESTRICTIONS: If application rates total 3.3 pounds per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3.3 pounds per acre remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting. Crops listed for treatment in this label may be planted into the treated area at any time; for other crops, wait 30 days between application and planting.

Spot Treatment, Over-the-Top Wiper Applications

USE INSTRUCTIONS: This product may be applied as a spot treatment or with wiper applicators in pastures. Applications may be made in the same area at 30-day intervals.

RESTRICTIONS: For spot treatments or wiper application methods using rates of 3.3 pounds per acre or less, the entire field or any portion of it may be treated. When spot treatments or wiper application are made using rates above 3.3 pounds per acre, no more than 10 percent of the total pasture may be treated at any one time. To achieve maximum performance, remove domestic livestock before application and wait 7 days after application before grazing livestock or harvesting.

Postemergent Weed Control (Broadcast Treatments)

USE INSTRUCTIONS: This product may be used to suppress competitive growth and seed production of annual weeds and undesirable vegetation in pastures. For selective applications with broadcast spray equipment, apply 6.5 to 8.7 ounces of this product per acre in early spring before desirable perennial grasses break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

RESTRICTIONS: Some stunting of perennial grasses will occur if broadcast applications are made when plants are not dormant. No waiting period is required between application and grazing or harvesting for feed. Use of higher application rates will cause stand reductions. Do not apply more than 3.3 pounds per acre per year onto pasture grasses except for renovation uses (see instructions above). If replanting is needed due to severe stand reduction, applications must be made at least 30 days prior to planting any crop not listed for treatment in this label.

10.5 Rangelands

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TYPES OF APPLICATIONS: Postemergence

This product will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands.

Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years must eliminate most of the viable seeds.

Grazing of treated areas must be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

USE INSTRUCTIONS: Apply 6.5 to 8.7 ounces of this product per acre to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are possible, where spring moisture is usually limited and fall germination allows for good weed growth.

For medusahead, apply 8.7 ounces of this product per acre at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Controlled burning may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

RESTRICTIONS: Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off. Do not use ammonium sulfate when spraying rangeland grasses with this product. No waiting period between treatment and feeding of livestock grazing is required. Do not apply more than 3.3 pounds per acre per year.

11.0 ROUNDUP READY CROPS

The following instructions include all applications which can be made onto the specified Roundup Ready crops during the complete cropping season. Do NOT combine these instructions with other directions made for crop varieties that do not contain the Roundup Ready gene, in the "ANNUAL AND PERENNIAL CROPS (ALPHABETICAL)" section of this label.

THIS PRODUCT IS ONLY FOR POSTEMERGENCE APPLICATION ONLY IN CROP VARIETIES DESIGNATED AS CONTAINING THE ROUNDUP READY GENE.

Applying this product to crop varieties that are not designated as Roundup Ready will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruits of crops, or any desirable plants that do not contain the Roundup Ready gene, since severe injury or destruction will result.

The Roundup Ready designation indicates that the crop variety contains a patented gene that provides tolerance to this product. Information on Roundup Ready crop varieties may be

Deleted: 1
Deleted: March 19,

51/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

obtained from your seed supplier. Roundup Ready crop varieties must be purchased from an authorized licensed seed supplier.

NOTE: Roundup Ready seed and the method of selectively controlling weeds in a Roundup Ready crop by applying glyphosate to the weeds and Roundup Ready crop are protected under several U.S. Patents, including 5,352,605 and 5,633,435. A license to use Roundup Ready seed must be obtained prior to use. Monsanto retains ownership of the gene and process technologies, and the Purchaser of the seed receives the right to use the licensed genes and technologies subject to the limited use license conditions. Seed containing the Roundup Ready trait cannot be used for research and demonstration, reverse engineering or in connection with herbicide registration. Progeny seed containing the Roundup Ready trait cannot be saved for replanting or transferred to others for replanting. Contact your Authorized Monsanto Retailer for information on obtaining a limited use license.

For ground applications with broadcast equipment, apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment use flat spray nozzles. Check for even distribution of spray droplets.

For serial applications, apply this product in 3 to 15 gallons of water per acre. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

See the "MIXING" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of this label for additional directions and restrictions on the application of this product.

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT for use in over-the-top applications of this product unless otherwise noted in this product label, supplemental labeling or fact sheets published separately by Monsanto.

Ammonium sulfate may be mixed with this product for applications to Roundup Ready crops. Refer to the "MIXING" section for use instructions for ammonium sulfate.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product. Follow the cleaning procedures specified on the label of the product(s) previously used. **THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.**

NOTE: The following instructions are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of this product can be used to control existing weeds prior to crop emergence. Some weeds, including black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morningglory, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed with multiple germination times or suppressed (stunted) weeds may require a second application of this product for complete control. The second application must be made after some regrowth has occurred and at least 10 days after a previous application of this product.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

11.1 Alfalfa with the Roundup Ready Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence.

| Maximum Allowable Combined Application Quantities Per Year | |
|---|-------------------|
| Combined total per year for all applications, | 8.7 lbs. per acre |
| Including Preplant during year of establishment | |
| Combined total per acre for in-crop applications | 6.5 lbs. per acre |
| For newly established and established stands | |

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting Roundup Ready alfalfa.

Postemergence

USE INSTRUCTIONS: Applications of this product may be made over the top of Roundup Ready alfalfa (in-crop) from emergence until 5 days prior to cutting. To maximize crop yield and quality potential of forage and hay, application of this product must be made after weeds have emerged but before alfalfa growth or re-growth interferes with spray coverage of the target weeds.

Weeds Controlled: For specific rates of application and instructions, refer to the "ANNUAL and PERENNIAL WEEDS RATE SECTIONS" in this booklet. When applied as directed, this product will control these annual and perennial grasses and broadleaf weeds. In addition to those weeds listed in these sections, this product will suppress or control the parasitic weed Dodder (*Cuscuta* spp) in Roundup Ready alfalfa. Repeat applications may be necessary for complete control.

NEW STAND ESTABLISHMENT (Seeding Year) – Due to the biology and breeding constraints of alfalfa, up to 10 percent of the seedlings may not contain the Roundup Ready gene and will not survive after the first application of this product. To eliminate the undesirable effects of stand gaps created by this loss of plants, a single application of at least 0.9 lb. per acre of this product must be applied at or before the 4-trifoliolate growth stage. Refer to the following table for application rates during stand establishment (seeding year).

| New Stand Establishment (Seeding Year) | |
|---|--------------------------|
| <u>Prior to First Cutting</u> | |
| From emergence up to 4 trifoliolate leaves | 0.9 to 1.8 lbs. per acre |
| From 5 trifoliolate leaves up to 5 days before first cutting | Up to 0.9 lb. per acre |
| <u>After First Cutting</u> | |
| In-crop application, per cutting, up to 5 days before cutting | Up to 1.8 lbs. per acre |

ESTABLISHED STANDS (Non-seeding Year) – For in-crop applications, per cutting, up to 5 days before cutting, apply this product up to 1.8 lbs. per acre.

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops. Where Roundup Ready alfalfa is grown with a companion or cover crop, or is overseeded with a second species, in-crop (over the top) applications of this product will eliminate the non-Roundup Ready (non-glyphosate tolerant) species. Any single in-crop application of this product must not exceed 35 ounces per acre. Sequential applications of this product must be at least 7 days apart. The combined total per year for all in-crop applications in both newly established (seeding year) and established stands (non-seeding year) must not exceed 5.6 lbs. per acre. Remove domestic livestock before application. Wait a minimum of 5 days after last application before grazing or cutting and feeding or forage or hay.

11.2 Canola with the Roundup Ready Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence.

DO NOT USE THIS PRODUCT ON CANOLA WITH THE ROUNDUP READY GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA.

| Maximum Allowable Combined Application Quantities Per Year | |
|---|--------------------|
| Total of Preplant, At-Planting, Preemergence applications | 35 ounces per acre |
| Total in-crop application from emergence to 6-leaf usage | 35 ounces per acre |

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting canola.

Postemergence

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

Weeds Controlled: For specific rates of application and instructions, refer to the "ANNUAL and PERENNIAL WEEDS RATE SECTIONS" in this booklet.

Single Application – Apply 8.8 to 17 ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and or growth reduction. Similar injury may result when applications of more than 8.8 ounces per acre are applied after the 4-leaf stage.

Sequential Application – Apply 17 ounces per acre to 1 to 3 leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications can be made to early emerging annual weeds and perennial weeds including Canada thistle and quackgrass or when controlling weeds with multiple application times.

RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops. No more than two over-the-top

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

54/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application must not exceed 35 ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

11.3 Corn with the Roundup Ready Gene

TYPES OF APPLICATION: Preplant, At-Planting, Preemergence, Postemergence (in-crop), Spot Treatment, Preharvest, Post-Harvest.

| Maximum Allowable Combined Application Quantities Per Year | |
|--|---------------------|
| Combined total per year for all applications | 8.7 pounds per acre |
| Total of Preplant, At-planting, Preemergence applications | 5.5 pounds per acre |
| Total in-crop applications from emergence through the V8 stage or 30 inches | 35 ounces per acre |
| Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest | 17 ounces per acre |

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting corn.

TANK MIXTURES: This product may be tank mixed with Aim, Bullet, Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, Lariat, Lasso, Micro-Tech, or Resource at 50 to 100 percent of labeled. Refer to the specific product label and observe all precautions and limitations on the label for any preemergence herbicide application, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines – the more restrictive requirements apply.

NOTE: For maximum weed control, a postemergence (in crop) application of this product must be applied following the use of less than labeled rates of the preemergence residual products listed above.

Postemergence (in-crop)

USE INSTRUCTIONS: This product may be applied postemergence to Roundup Ready corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first.

When applied as directed, this product controls labeled annual grass and broadleaf weeds in Roundup Ready corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of this product. The postemergent application of 13 to 17 ounces per acre of this product must be made before the weeds reach a height and/or density that the weeds become competitive with the crop, generally 4 inch tall weeds or less.

This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on this label. If new flushes of weeds occur, a sequential application of this product at 13 to 17 ounces per acre will control the labeled grasses and broadleaf weeds.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TANK MIXTURES: This product may be applied to tank mixture with Bullet, Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, and Micro-Tech at 50 to 100 percent of labeled rate. This product may be applied in tank mixture with Permit and atrazine at labeled rates. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing, restrictions, soil restrictions, minimum recropping interval and rotational guidelines – the more restrictive requirements apply.

| Tank-mix Partner | Maximum Height of Corn For Application |
|------------------|---|
| Degree | 11 inches |
| Degree Xtra | |
| Harness | |
| Harness Xtra | |
| Harness Xtra 5.6 | |
| Bullet* | 5 inches |
| Micro-Tech* | |
| Permit | 30 inches |
| atrazine | 12 inches |

*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops. Single in-crop applications of this product are not to exceed 17 ounces per acre. Sequential in-crop applications of this product from emergence through the V8 stage or 30 inches must not exceed 35 ounces per acre per growing season. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage.

Preharvest

USE INSTRUCTIONS: In Roundup Ready corn, up to 17 ounces per acre of this product can be applied preharvest. Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

RESTRICTIONS: Allow a minimum of 7 days between application and harvest.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large seeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

POSTEMERGENCE APPLICATIONS TO ROUNDUP READY 603 CORN

INFORMATION

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

56/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Note: The instructions provided below are specific to, and must only be used with, Roundup Ready 603 Corn hybrids. Do not combine these instructions with those in the "Corn With The Roundup Ready Gene" section of this label booklet, or with any other Roundup Ready corn instructions on labeling for this or other glyphosate-containing product. See "Annual Maximum Use Rate" in the "INFORMATION" section of this label booklet for additional information.

The use of the higher in-crop over the top rates on other than Roundup Ready 603 corn may cause crop injury and reduce yields.

Application instructions

For Roundup Ready 603 corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first, this product may be applied over-the-top broadcast or with drop nozzles. When corn height is 24 to 30 inches (free standing), for optimum spray coverage and weed control, drop nozzles are recommended. For corn heights 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles adjusted to avoid spraying into the whorls of the corn plants.

Single in-crop applications of this product must not exceed 26 ounces per acre.

Maximum Allowable Combined Application Quantities Per Year

~~Preplant; At-Planting; Preemergence:~~ Maximum amount of this product which can be applied prior to crop emergence is 5.5 pounds per acre.

Postemergence (in-crop): Maximum combined total of multiple in-crop applications from emergence through the 48 inch stage is 3.2 pounds per acre.

Preharvest: Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) with 35 percent grain moisture or less until 7 days before harvest is 17 ounces per acre. See Restrictions on preharvest applications.

Cropping Season: Combined total per year for all applications may not exceed 8.6 pounds per acre.

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product under hard water conditions, drought conditions or when tank mixed with Bullet® or Micro-Tech® herbicides. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides. Thoroughly rinse the spray system with clean water after use to reduce corrosion. Do not add other additives, including fertilizers and micro-nutrients as this may result in increased potential for crop injury.

ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING THIS PRODUCT.

Deleted: 1
Deleted: March 19,

57/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

For ground applications: Use the listed rates of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Carefully select correct nozzles and spray pressure to avoid spraying a fine mist. Check for even distribution of spray droplets.

For aerial applications: Use the listed rates of this product in 3 to 15 gallons of spray solution per acre. Do not exceed 17 ounces per acre. See "ANNUAL WEEDS RATE SECTION" of the label booklet for specific weed rates. AVOID DRIFT – DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITIONS WHICH FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Weed Control Instructions

Apply 13 to 17 ounces of this product per acre for control of labeled grasses and broadleaf weeds in conventional and no-till corn production systems. Refer to the label booklet for rate directions for specific annual weeds. This product applied at up to 26 ounces per acre will control or suppress the growth of perennial weeds including: bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed, and wirestem muhly. For additional information on perennial weeds, see the "PERENNIAL WEEDS RATE SECTION".

~~Preplant, Preemergence, At-Planting~~

USE INSTRUCTIONS: This product may be applied alone or in a tank-mixture before, during or after planting corn.

TANK MIXTURES: This product may be tank mixed with Bullet, Degree, Degree, Xtra, Harness Xtra, Harness Xtra 5.6L, Lariat, Lasso or Micro-Tech at 50 to 100 percent of labeled rate. Refer to the specific product label and observe all precautions and limitations on the label for any preemergence herbicide application, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines the more restrictive requirements apply.

NOTE: For maximum weed control, a postemergence (in-crop) application of this product must be applied following the use of less than labeled rates of the preemergence residual products listed above.

Preemergence followed by Postemergence Weed Control Program

USE INSTRUCTIONS: This product may be applied postemergence in-crop following any labeled preemergence herbicide application. The post application of this product must be made before the weeds reach a height and/or density that the weeds become competitive with the crop.

An in-crop application of this product at the listed rate will provide control of emerged weeds listed on the label. This product may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready 603 corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are necessary for optimum spray coverage and weed control. For corn heights 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Postemergence Only Weed Control Program

USE INSTRUCTIONS: This product may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on the label. The postemergence application of this product must be made before the weeds reach a height and/ or density that the weeds become competitive with the crop. If new flushes of weeds occur, a sequential application of this product at 13 to 17 ounces per acre will control the labeled grasses and broadleaf weeds. This product may be applied over-the-top broadcast or with drop nozzles postemergence to Roundup Ready 603 corn from emergence through the V8 stage or until corn height reaches 30 inches (free standing), whichever comes first. When corn height is 24 to 30 inches drop nozzles are necessary for optimum spray coverage and weed control. For corn height 30 to 48 inches (free standing), apply this product only using ground application equipment with drop nozzles and avoid spraying into the whorls of the corn plants.

TANK MIXTURES: This product may be applied in tank mixture with a labeled rate of Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, Micro-Tech and Bullet at 50 to 100 percent of labeled rates. This product may be applied in tank mixture with Permit® and atrazine at labeled rates. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum re-cropping interval and rotational guidelines – the more restrictive requirements apply. Tank mixtures with other products may result in increased potential for crop injury and/or weed antagonism. Refer to the table below for height limitation for tank mix partner.

| Tank Mix Partner | Maximum Height Of Corn For Application |
|---|--|
| Degree Degree Xtra Harness Harness Xtra Harness Xtra 5.6L | 11 inches |
| Bullet* Micro-Tech* | 5 inches |
| Permit | 30 inches |
| Atrazine | 12 inches |

* Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

RESTRICTIONS: See the "ROUNDUP READY CROPS" section for precautionary instructions for use in Roundup Ready Crops. Single in-crop applications of this product must not exceed 26 ounces per acre. Allow a minimum of 10 days between in-crop applications of this product. Allow a minimum of 50 days between application of this product and harvest of corn forage or grain. For applications at preharvest timing (see Preharvest section of this supplemental label), allow a minimum of 7 days between application and harvest or feeding of corn stover or grain. There are no rotational crop restrictions following applications of this product.

Preharvest

USE INSTRUCTIONS: A single preharvest application of up to 17 ounces per acre of this product may be made, if no more than a total of 35 ounces of this product has been previously applied in over-the-top or drop nozzle applications. Make a preharvest

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

application at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

RESTRICTIONS: Do not make a preharvest application of this product if more than a combined total of 35 ounces of this product has been previously applied in over-the-top or drop nozzle applications. Allow a minimum of 7 days between a preharvest application and harvest or feeding of corn stover or grain.

Postharvest

USE INSTRUCTIONS: This product may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

RESTRICTIONS: Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

Bullet, Harness and Micro-Tech are registered trademarks of Monsanto Technology LLC Permit is a trademark of, and used under license from, Nissan Chemical Industries, Ltd.

11.4 Soybeans with the Roundup Ready Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence, Preharvest, Post-Harvest.

| Maximum Allowable Combined Application Quantities Per Year | |
|---|---------------------|
| Combined total per year for all applications | 8.7 pounds per acre |
| Total of Preplant, At-planting, Preemergence applications | 5.5 pounds per acre |
| Total in-crop applications from cracking throughout flowering | 52 ounces per acre |
| Maximum preharvest application rate | 17 ounces per acre |

RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting soybeans.

Postemergence

USE INSTRUCTIONS: When applied as directed, this product will control labeled annual grasses and broadleaf weeds in Roundup Ready soybeans. Applications of this product can be made in Roundup Ready soybeans from emergence (cracking) throughout flowering. Refer to the "ANNUAL WEEDS RATE SECTION" in this label for rate instructions for specific annual weeds. Apply 17 ounces per acre on 2 to 8 inch tall weeds. Weeds will generally be 2

Deleted: 1

Deleted: March 19,

60/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

to 8 inches tall, 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of this product. This product may be used up to 35 ounces per acre in any single in-crop application for control of annual weeds and where heavy weed densities exist.

A 17 to 35 ounce per acre rate (single or multiple applications) of this product will control or suppress perennial weeds including: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, maretail (horseweed), nutsedge, quackgrass, rhizome johnsongrass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with this product.

Under adverse growing conditions including drought, hail, wind damage or a poor soybean stand that slows or delays canopy closure, a sequential application of this may be necessary to control late flushes of weeds. **IN THE SOUTHERN STATES A SEQUENTIAL APPLICATION OF THIS PRODUCT WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE ROUNDUP READY SOYBEAN CROP.** To control giant ragweed, apply 17 ounces per acre of this product when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

RESTRICTIONS: The combined total application from emergence through harvest must not exceed 52 ounces per acre. The maximum rate for any single in-crop application is 35 ounces per acre. The maximum combined total of this product that can be applied during flowering is 35 ounces per acre.

Preharvest

USE INSTRUCTIONS: This product provides weed control when applied prior to harvest of soybeans. Up to 17 ounces per acre of this product can be applied by aerial or ground application.

RESTRICTIONS: Care must be taken to avoid excessive seed shatter loss due to ground application equipment. Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

Post-Harvest

USE INSTRUCTIONS: This product may be applied after harvest of Roundup Ready soybeans. Higher rates may be required for control of large weeds which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

11.5 Sugar beets with the Roundup Ready Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Postemergence.

| Maximum Allowable Combined Application Quantities Per Year | |
|---|---------------------|
| Combined total per year for all applications | 8.7 pounds per acre |
| Total of Preplant, At-planting, Preemergence applications | 5.5 pounds per acre |
| Emergence to 8-leaf stage | 44 ounces per acre |
| Between 8-leaf stage and canopy closure | 35 ounces per acre |

Deleted: 1
Deleted: March 19,

61/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

RESTRICTIONS: See the "ROUNDUP READY CROPS" section of this label for precautionary instructions for use in Roundup Ready crops. The combined total application from crop emergence through harvest must not exceed 4.8 pounds per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 26 ounces per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 17 ounces per acre. Allow a minimum of 30 days between last application and sugar beet harvest.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting of Roundup Ready sugar beets.

Postemergence

USE INSTRUCTIONS: This product may be applied postemergent over-the-top to Roundup Ready sugar beets from emergence to 30 days prior to harvest. To maximize yield potential spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of this product may be made with at least 10 days between applications. Refer to the "ANNUAL WEEDS RATE SECTIONS" in this label for rate instructions for specific annual weeds. This product will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

11.6 Cotton with the Roundup Ready Gene

TYPES OF APPLICATIONS: Preplant, At-Planting, Preemergence, Over-the-Top, Selective Equipment, Preharvest.

NOTE: Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 8.8 to 26 ounces per acre of this product.

Maximum Allowable Combined Application Quantities Per Year

| | |
|---|---------------------|
| Combined total per year for all applications | 8.7 pounds per acre |
| Total of Preplant, At-planting, Preemergence applications | 5.5 pounds per acre |
| Total over-the-top applications from cracking to layby | 44 ounces per acre |
| Total precision post-directed for hooded applications through layby | 35 ounces per acre |
| Maximum preharvest application rate | 35 ounces per acre |

RESTRICTIONS: See the "ROUNDUP READY CROPS" section for precautionary instructions for use in Roundup Ready crops. The combined total application of this product from cotton emergence until harvest must not exceed 6.4 pounds per acre. **NO MORE THAN TWO OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THOROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT. NO MORE THAN TWO APPLICATIONS MUST BE MADE FROM THE 5-LEAF STAGE THROUGH LAYBY. SEQUENTIAL IN-CROP OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF THIS PRODUCT MUST BE AT LEAST 10 DAYS APART AND COTTON MUST HAVE**

Deleted: 1
Deleted: March 19,

62/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS.
ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.

Sprayer Preparation: It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care must be taken to thoroughly clean all equipment prior to use.

Tank mixtures with other herbicides may result in reduced weed control, or may cause crop injury for applications where the spray contacts the cotton plant.

Preplant, Preemergence, At-Planting

USE INSTRUCTIONS: This product may be applied before, during or after planting cotton.

Over-the-Top

USE INSTRUCTIONS: This product may be applied by aerial or ground application equipment at rates up to 17 ounces per acre per application postemergence to Roundup Ready cotton from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the 4-leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. There are no rotational crop restrictions following applications of this product.

Over-the-top applications: Up to 44 ounces per sprayed acre of this product may be applied by aerial or ground broadcast application equipment postemergence to Roundup Ready cotton from the ground cracking stage until layby. The "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE" sections in the label must be used to determine application rate. Any single over-the-top application must not exceed 35 ounces per sprayed acre. Sequential applications of this product must be at least 7 days apart.

With ground broadcast equipment: apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spaying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

For aerial applications, apply this product in 3 to 15 gallons of water per acre. DO NOT EXCEED A MAXIMUM RATE OF 17 OUNCES PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Salvage Treatment. This treatment may be used after the 4-leaf stage of development and must only be used where weeds threaten to cause the loss of the crop. 1.7 pounds per acre may be applied either as an over-the-top applications or as a post-directed treatments sprayed higher on the cotton plants and over the weeds. NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT MUST BE USED PER GROWING SEASON.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

NOTE: For specific rates of application and instructions, refer to the "ANNUAL WEEDS" and "PERENNIAL WEEDS RATE" sections of the label. This product applied at 17 to 35 ounces per acre will control or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common Bermudagrass, silverleaf nightshade, trumpet creeper, and redvine. Fall preharvest applications may be required for control of these perennial weeds. Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

RESTRICTIONS: See the "ROUNDUP READY CROPS" section for precautionary instructions for use in Roundup Ready crops.

Selective Equipment

USE INSTRUCTIONS: This product may be applied using precision post-directed or hooded sprayers at rates up to 35 ounces per acre per application to Roundup Ready cotton through layby. At this stage, post-directed equipment must be used which directs the spray to the base of the cotton plants. These application methods may be preferred when there is a need to direct the spray onto weeds that are growing under the crop canopy. Contact of the spray with cotton leaves must be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches). Sequential in-crop applications must be at least 7 days apart from any other in-crop application of this product.

RESTRICTIONS: See the "SELECTIVE EQUIPMENT" part of the "APPLICATION EQUIPMENT AND TECHNIQUES" section for information on proper use and calibration of this equipment.

Preharvest

USE INSTRUCTIONS: This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton any time after layby up to 7 days prior to harvest. This product may be tank mixed with DEF™ 6, Folex™, Ginstar, or Prep™ to enhance cotton leaf drop. Allow a minimum of 7 days between final application and harvest. No more than 35 ounces of this product per acre may be applied preharvest (between layby and seven days prior to harvest). Do not apply more than 17 ounces of this product per acre by air. NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready cotton.

Restrictions: The combined total application from crop emergence until harvest must not exceed 8.7 pounds per acre. Allow a minimum of 7 days between final application and harvest. Do not apply this product to cotton grown for seed, as a reduction in germination or vigor may occur.

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.

11.7 Seed Production of Select Crops with the Roundup Ready Gene

Seed Production of ALFALFA with the Roundup Ready Gene

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT ALFALFA IN PRODUCTION FIELDS OF ALFALFA CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF ALFALFA WILL RESULT IF ALFALFA VARIETIES THAT DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE INSTRUCTIONS

This product will control non-glyphosate tolerant alfalfa in seed production fields of alfalfa containing the Roundup Ready gene. Apply up to 35 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. Subsequent applications of up to 35 ounces per acre each may be applied, if needed to control non-glyphosate tolerant alfalfa plants.

DO NOT EXCEED A MAXIMUM RATE OF 8.7 POUNDS OF THIS PRODUCT PER ACRE PER YEAR.

Application timing – This product can be applied to Roundup Ready alfalfa from emergence to harvest.

Treated alfalfa or the resulting seed may not be used for food or feed. Do not feed or graze treated alfalfa. Do not process treated alfalfa or resulting seed for food or feed.

Seed Production of LETTUCE with the Roundup Ready Gene

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT LETTUCE IN PRODUCTION FIELDS OF LETTUCE CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH OF LETTUCE WILL RESULT IF LETTUCE VARIETIES THAT DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE INSTRUCTIONS

This product will control non-glyphosate tolerant lettuce in seed production fields of lettuce containing the Roundup Ready gene. Apply up to 35 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 35 ounces per acre may be applied, if needed to control non-glyphosate tolerant lettuce plants.

DO NOT EXCEED A MAXIMUM RATE OF 4.3 POUNDS OF THIS PRODUCT PER ACRE PER YEAR.

Application timing – This product can be applied to Roundup Ready lettuce from emergence to harvest.

Treated lettuce may not be used for food or feed. Do not feed or graze treated lettuce. Do not process treated lettuce for food or feed.

Seed Production of RICE with the Roundup Ready Gene

Deleted: 1
Deleted: March 19,

65/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT RICE IN PRODUCTION FIELDS OF RICE CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF RICE VARIETIES THAT DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE INSTRUCTIONS

This product will control non-glyphosate tolerant rice in seed production fields of rice containing the Roundup Ready gene. Apply up to 35 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 35 ounces per acre may be applied, if needed to control non-glyphosate tolerant rice plants.

DO NOT EXCEED A MAXIMUM RATE OF 4.3 POUNDS OF THIS PRODUCT PER ACRE PER YEAR.

Application timing – This product can be applied to Roundup Ready rice from emergence to harvest.

Treated rice may not be used for food or feed. Do not feed or graze treated rice. Do not process treated rice for food or feed.

Seed Production of WHEAT with the Roundup Ready Gene

NOTE: THIS PRODUCT MAY BE USED FOR CONTROL OF NON-GLYPHOSATE TOLERANT WHEAT IN PRODUCTION FIELDS OF WHEAT CONTAINING THE ROUNDUP READY GENE. SEVERE INJURY OR DEATH WILL RESULT IF WHEAT VARIETIES THAT DO NOT CONTAIN THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.

USE INSTRUCTIONS

This product will control non-glyphosate tolerant wheat in seed production fields of wheat containing the Roundup Ready gene. Apply up to 17 ounces of this product in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 17 ounces per acre may be applied, if needed to control non-glyphosate tolerant wheat plants.

DO NOT EXCEED A MAXIMUM RATE OF 35 OUNCES OF THIS PRODUCT PER ACRE PER YEAR.

Application timing – This product can be applied to Roundup Ready wheat from emergence to harvest

Treated wheat may not be used for food or feed. Do not feed or graze treated wheat. Do not process treated wheat for food or feed.

12.0 NON-CROP USES AROUND THE FARMSTEAD

TYPES OF APPLICATIONS: Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stump, Habitat Management.

12.1 Weed Control, Trim-and-Edge

USE INSTRUCTIONS: This product may be used to control annual weeds, perennial weeds and woody brush which are found in any part of the farmstead, including building

Deleted: 1
Deleted: March 19,

66/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

foundations, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, shelterbelts, prior to landscape plantings and equipment storage areas.

TANK MIXTURES: This product may be tank mixed with the following products. Refer to these product labels for approved farmstead sites and application rates. For annual weeds, use 17 ounces per acre of this product when weeds are less than 6 inches tall, 26 ounces per acre when weeds are 6 to 12 inches tall and 35 ounces per acre when weeds are greater than 12 inches tall. For perennial weeds, apply 35 ounces to 5.3 pounds per acre in these tank mixes. For tank mixtures with these products through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "ANNUAL WEEDS- HAND-HELD OR HIGH VOLUME EQUIPMENT" section of this label for listed rates.

- | | |
|-----------------|----------------|
| Arsenal | Plateau |
| Banvel/Clarity | Princep DF |
| Barricade 65WG | Princep Liquid |
| Diuron | Ronstar 50 WP |
| Endurance | Sahara |
| Escort | Simazine |
| Karmex DF | Surflan |
| Krovar I DF | Telar |
| Oust | Vanquish |
| Pendulum 3.3 EC | 2,4-D |
| Pendulum WDG | |

This product plus dicamba tank mixtures may not be applied by air in California.

12.2 Greenhouse/Shadehouse

This product may be used to control weeds in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

12.3 Chemical Mowing

USE INSTRUCTIONS: This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 3.2 ounces of this product per acre when treating Kentucky bluegrass. Use 3.2 ounces of this product when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Use 8.8 ounces of this product per acre when treating bermudagrass. Use 35 ounces of this product per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

RESTRICTIONS: Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

12.4 Cut Stump

TYPES OF APPLICATION: Treating cut stumps in any non-crop site listed on this label

USE INSTRUCTIONS: This product will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a solution prepared by dissolving 2.1 to 3.7 pounds

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

67/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

of this product per gallon of water to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications must be made during periods of active growth and full leaf expansion.

- | | |
|-------------------|-------------|
| Alder | Reed, giant |
| Eucalyptus | Saltcedar |
| Madrone | Sweetgum |
| Oak | Twin oak |
| Pepper, Brazilian | Tan oak |
| Pine, Austrian | Willow |

RESTRICTIONS: Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

12.5 Habitat Management

TYPES OF USES: Habitat Restoration and Maintenance, Wildlife Food Plots.

Habitat Restoration and Maintenance

USE INSTRUCTIONS: This product may be used to control exotic and other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

Wildlife Food Plots

USE INSTRUCTIONS: This product may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage.

13.0 ANNUAL WEEDS RATE SECTION (Alphabetically by Species)

When water carrier volumes are between 16 and 40 gallons per acre for ground applications and between 6 and 15 gallons per acre for aerial applications, the following use rates will control the annual weeds listed in the table below:

- 17 ounces per acre – grass and broadleaf annual weeds less than 6 inches in height or circumference and vines less than 3 inches in length.
- 26 ounces per acre – grass and broadleaf annual weeds 6 to 12 inches in height or circumference and vines 3 to 6 inches in length.
- 35 ounces per acre – grass and broadleaf annual weeds greater than 12 inches in height or circumference and vines greater than 6 inches in length.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

48/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

WHEN WATER CARRIER VOLUMES ARE BETWEEN 3 AND 15 GALLONS PER ACRE FOR GROUND APPLICATIONS AND BETWEEN 3 TO 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS, USE THE RATES SPECIFIED FOR INDIVIDUAL WEEDS AS FOLLOW IN THE "ANNUAL WEEDS RATE TABLE". (Alphabetically by Species).

Apply to actively growing annual weeds. Annual weeds are generally easiest to control when they are small.

Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Maximum size refers to the maximum plant height, length of runners for vines, or circumference of rosette plants in inches.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

This product may be used up to 35 ounces per acre where heavy weed densities exist.

ANNUAL WEEDS RATE TABLE

| WEED SPECIES Maximum size (inches) | RATE (ounces per acre) | | | | |
|---------------------------------------|---------------------------|-------|------|-----|------|
| | 9.5 | 14 | 19 | 24 | 28 |
| Ammannia, purple | 2.8 | 5.6 | 11 | - | 16.7 |
| Annoda, spurred | - | 1.9 | 2.8 | 4.6 | 7.4 |
| Barley | 16.7 | 16.7+ | - | - | - |
| Barnyardgrass | - | 2.8 | 5.6 | 6.5 | 8.4 |
| Bassia, fivehook | - | - | 5.6 | - | - |
| Beggarweed, Florida | - | 4.6 | 7.4 | - | - |
| Bittercress | 11 | 18.6 | - | - | - |
| Bluegrass, annual | 9.3 | - | - | - | - |
| Bluegrass, bulbous | 5.6 | - | - | - | - |
| Brome, downy ^{1,2} | 5.6 | 11 | - | - | - |
| Brome, Japanese | 5.6 | 11 | 22.3 | - | - |
| Browntop panicum | 5.6 | 7.4 | 11 | - | 22.3 |
| Buckwheat, wild ³ | - | 0.9 | 1.9 | - | - |
| Burcucumber | - | 5.6 | 11 | - | 16.7 |
| Buttercup | 11 | 18.6 | - | - | - |
| Carolina geranium | - | - | 3.7 | - | 8.4 |
| Carpetweed | - | 5.6 | 11 | - | - |
| Cheat ² | 5.6 | 18.6 | - | - | - |
| Chervil | 18.6 | - | - | - | - |
| Chickweed | - | 11 | 16.7 | - | - |
| Cocklebur | 11 | 16.7 | 22.3 | - | 33.4 |
| Copperleaf, Hophornbeam | - | 1.9 | 3.7 | - | 5.6 |
| Copperleaf, Virginia | - | 1.9 | 3.7 | - | 5.6 |
| Coreopsis, plains | - | 5.6 | 11 | - | 16.7 |

Deleted: 1
Deleted: March 19,

49/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | | | | |
|--|-----|----------|------|-----|------|
| Corn, volunteer | 5.6 | 11 | 18.6 | - | - |
| Corn Speedwell | 11 | - | - | - | - |
| Crabgrass | 2.8 | 5.6 | 11 | - | - |
| Crowfootgrass | - | - | 5.6 | - | 11 |
| Cutleaf evening primrose | - | - | 2.8 | - | 5.6 |
| Devilsclaw (unicorn plant) | - | 2.8 | 5.6 | - | - |
| Dwarf dandelion | 11 | - | - | - | - |
| Eastern manna grass | 7.4 | 11 | - | - | - |
| Eclipta | - | 3.7 | 7.4 | 11 | - |
| Fall panicum | 3.7 | - | 5.6 | - | 11 |
| Falsedandelion | - | 18.6 | - | - | - |
| Falseflax, smallseed | 11 | - | - | - | - |
| Fiddleneck | - | 5.6 | 11 | - | - |
| Field Pennycrass | 5.6 | 11 | - | - | - |
| Filaree | - | - | 5.6 | - | 11 |
| Fleabane, annual | 5.6 | 18.6 | - | - | - |
| Fleabane, hairy (Conyza bonariensis) | - | - | 5.6 | - | 9.3 |
| Fleabane, rough | 2.8 | 5.6 | 11 | - | - |
| Florida pusley | - | - | 3.7 | - | 5.6 |
| Foxtail, giant, bristly, yellow | 5.6 | 11 | 18.6 | - | - |
| Foxtail, Carolina | 9.3 | - | - | - | - |
| Foxtrail, green | 11 | - | - | - | - |
| Goatgrass, jointed | 5.6 | 11 | - | - | - |
| Goosegrass | - | 2.8 | 5.6 | - | 11 |
| Grain sorghum (milo) | 5.6 | 11 | 18.6 | - | - |
| Groundcherry | - | 2.8 | 5.6 | - | 8.4 |
| Groundsel, common | - | 5.6 | 9.3 | - | - |
| Hemp sesbania | - | 1.9 | 3.8 | 5.6 | 7.4 |
| Henbit | - | - | 5.6 | - | 11 |
| Horseweed/Marestail (Conyza canadensis) | - | 5.6 | 11 | - | 16.7 |
| Itchgrass | 5.6 | 7.4 | 11 | - | 16.7 |
| Jimsonweed | - | - | 11 | - | 16.7 |
| Johnsongrass, seedling | 5.6 | 11 | 16.7 | - | 22 |
| Junglerice | - | 2.8 | 5.6 | 6.5 | 8.4 |
| Knotweed | - | - | 5.6 | - | 11 |
| Kochia ⁴ | - | 2.8to5.6 | 11 | - | - |
| Lambsquarters | - | 5.6 | 11 | - | 18.6 |
| Little barley | 5.6 | 11 | - | - | - |
| London rocket | 5.6 | - | 22.3 | - | - |
| Mayweed | - | 1.9 | 5.6 | 11 | 16.7 |
| Morningglory, annual (Ipomoea spp) | - | - | 2.8 | - | 5.6 |
| Mustard, blue | 5.6 | 11 | 16.7 | - | - |
| Mustard, tansy | 5.6 | 11 | 16.7 | - | - |
| Mustard, tumble | 5.6 | 11 | 16.7 | - | - |
| Mustard, wild | 5.6 | 11 | 16.7 | - | - |
| Nightshade, black | - | 3.7 | 5.6 | - | 11 |
| Nightshade, hairy | - | 3.7 | 5.6 | - | 11 |
| Oats | 2.8 | 5.6 | 16.7 | - | - |
| Pigweed species | - | 11 | 16.7 | 22 | - |
| Prickly lettuce | - | 5.6 | 11 | - | - |
| Purslane | - | - | 2.8 | - | 5.6 |

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | | | | |
|------------------------------------|-----|------|-------|-----|------|
| Ragweed, common | - | 5.6 | 11 | - | 16.7 |
| Ragweed, giant | - | 5.6 | 11 | - | 16.7 |
| Red rice | - | - | 3.8 | - | - |
| Rye, volunteer/cereal ² | 5.6 | 16.7 | 16.7+ | - | - |
| Ryegrass | - | - | 5.6 | - | 11 |
| Sanbur, field | 5.6 | 11 | - | - | - |
| Sanbur, longspine | 5.6 | 11 | - | - | - |
| Shattercane | 5.6 | 11 | 22.3 | - | - |
| Shepherd's purse | 5.6 | 11 | - | - | - |
| Sicklepod | - | 1.9 | 3.7 | - | 7.4 |
| Signalgrass, broadleaf | - | 2.8 | 5.6 | 6.5 | 8.4 |
| Smartweed, Pennsylvania | - | - | 5.6 | - | 8.4 |
| Sowthistle, annual | - | - | 5.6 | - | 11 |
| Spanishneedles | - | - | 5.6 | - | 11 |
| Speedwell, purslane | 11 | - | - | - | - |
| Sprangletop | 5.6 | 11 | 18.6 | - | - |
| Spurge, prostrate | - | 5.6 | 11 | - | - |
| Spurge, spotted | - | 5.6 | 11 | - | - |
| Spurry, umbrella | 5.6 | - | - | - | - |
| Stinkgrass | - | 11 | - | - | - |
| Sunflower | 11 | 16.7 | - | - | - |
| Swinecress | - | 4.6 | 11 | - | - |
| Teaweed/Prickly sida | - | 1.9 | 3.7 | - | 5.6 |
| Texas panicum | 5.6 | 7.4 | 11 | - | 22 |
| Thistle, Russian ⁵ | - | 5.6 | 11 | - | - |
| Velvetleaf | - | - | 5.6 | - | 11 |
| Virginia pepperweed | - | 16.7 | - | - | - |
| Waterhemp | - | - | 5.6 | - | 11 |
| Wheat ² | 5.6 | 11 | 16.7 | - | - |
| Wheat, (overwintered) | - | 5.6 | 11 | - | 16.7 |
| Wild oats | 2.8 | 5.6 | 16.7 | - | - |
| Wild proso millet | - | 5.6 | 11 | - | 16.7 |
| Witchgrass | - | 11 | - | - | - |
| Woolly cupgrass | - | 5.6 | 11 | - | - |
| Yellow rocket | - | 11 | 18.6 | - | - |

¹ For control of downy brome in no-till systems, use 13 ounces per acre.

² Performance is better if application is made before this weed reaches the boot stage of growth.

³ Use 13 ounces per acre of this product to control wild buckwheat in the cotyledon to 2-leaf stage. Use 17 ounces per acre to control 2-to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 17 ounces followed by 17 ounces of this product per acre.

⁴ Do not treat kochia in the button stage.

⁵ Control of Russian thistle may vary based on environmental conditions and spray coverage. Whenever possible, a tank mixture with 2,4-D as described below may improve control.

13.1 Annual Weeds – Tank Mixtures with 2,4-D, Dicamba, or Tordon 22K

Deleted: 1

Deleted: March 19,

71/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

6.5 to 8.8 ounces of this product plus ¼ pound of dicamba or 0.5 pounds of 2,4-D or 1 to 2 fluid ounces of Tordon 22K per acre will control the following weeds with the maximum height or length indicated: 6 inches – prickly lettuce, marehail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Tordon 22K only); 12 inches – cocklebur lambsquarters, pigweed, Russian thistle (2,4-D only).

8.8 ounces of this product plus 0.5 pound of 2,4-D per acre will control the following weeds when they are a maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA

13.2 Annual Weeds – Hand-Held or High-Volume Equipment

For control of weeds listed in the "ANNUAL WEEDS RATE SECTION", apply a 0.3 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 0.6 percent solution.

For best results, use a 1 percent solution on harder-to-control perennials, including Bermudagrass, dock, field, bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 2.6 percent solution for annual and perennial weeds and a 2.6 to 5.1 percent solution for woody brush and trees.

13.3 Annual Weeds – Tank Mixtures with Atrazine for Fallow and Reduced Tillage Systems

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota, and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

13 to 16 ounces of this product plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 16 ounces for control), Downy brome, Green foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass and Kochia (add 0.12 pound of dicamba for control).

14.0 PERENNIAL WEEDS RATE SECTION

(Alphabetically by Species)

Apply to actively growing perennial weeds.

NOTE: If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the listed stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Deleted: 1
Deleted: March 19.

72/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parentetical text) is informational.)

Unless otherwise stated, allow 7 or more days after application before tillage.

Best results are obtained when soil moisture is adequate for active weed growth.

PERENNIAL WEEDS RATE TABLE
(Alphabetically by Species)

| Weed Species | Rate (LB/A) | Water Volume (GPA) | Hand-Held % Solution |
|--------------|-------------|--------------------|----------------------|
|--------------|-------------|--------------------|----------------------|

| | | | |
|---------|-----------|--------|----|
| Alfalfa | 1.1 – 2.2 | 3 – 10 | 1% |
|---------|-----------|--------|----|

Make applications after the last hay cutting in the fall. Allow Alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications must be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

| | | | |
|---------------|-----|--------|------|
| Alligatorweed | 4.4 | 3 – 20 | 0.7% |
|---------------|-----|--------|------|

For partial control, apply when most of the plants are in bloom. Repeat applications will be required to maintain control.

| | | | |
|------------------------------|---|---|----------|
| Anise (fennial) ¹ | - | - | 0.6 – 1% |
|------------------------------|---|---|----------|

For hand held, apply as a spray-to-wet treatment

| | | | |
|-------------------------|-----------|--------|----|
| Bahiagrass ² | 3.2 – 5.5 | 3 – 20 | 1% |
|-------------------------|-----------|--------|----|

| | | | |
|-----------|-----|-------|----|
| Bentgrass | 1.6 | 10-20 | 1% |
|-----------|-----|-------|----|

For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass must have at least 3 inches of growth. Tillage prior to treatment must be avoided. Tillage 7 to 10 days after application is recommended for best results.

| | | | |
|---------------|-----------|--------|----|
| Bermudagrass, | 3.2 – 5.5 | 3 – 20 | 1% |
|---------------|-----------|--------|----|

For control, apply 5.5 pounds of this product per acre. For partial control, apply 52 ounces per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control.

| | | | |
|------------------------------------|-----------|--------|----|
| Bermudagrass, Water (knotgrass) | 1.1 – 1.6 | 5 – 10 | 1% |
|------------------------------------|-----------|--------|----|

Apply 26 ounces of this product in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 17 ounces of this product in 5 to 10 gallons of water per acre. Fallow fields must be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length.

Deleted: 1.

Deleted: March 19,

73/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

This product is not registered in California for use on water Bermudagrass.

Bindweed, field 0.6 – 5.5 3 – 2 1%
Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.

For control, apply 4.4 to 5.5 pounds of this product per acre west of the Mississippi River and 3.2 to 4.4 pounds east of Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late in late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 35 ounces of this product plus 0.5 pound a.i. of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 17 to 35 ounces of this product plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications must be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 8.9 ounces of this product plus 0.5 pound a.i. of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only. Applications must be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

In California only, apply 17 ounces to 5.5 pounds of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 17 ounces of this product in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky 1.1 – 2.2 3 – 40 1%

Apply 35 ounces of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 17 to 26 ounces of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas 4.4 – 5.5 3 – 40 1%

Apply 4.4 to 5.5 pounds of this product per acre west of the Mississippi River and 4.4 pounds per acre east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenform 3.2 – 4.4 3 – 40 0.6 – 0.7%

Apply to fully expanded fronds that are at least 18 inches long

Deleted: 1
Deleted: March 19,

74/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Bromegrass, smooth 1.1 – 2.2 3 – 40 1%

Apply 35 ounces of this product in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 17 to 26 ounces of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf -- 3 – 20 1%

For control, apply 35 ounces of this product plus ½ pound of dicamba per acre. For partial control, apply 17 ounces of this product plus ½ pound of dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed² 2.2 - 3.4 3 – 40 1%

Cattail² 3.2 - 5.5 3 – 40 1%

Clover; red or white¹ 3.2 - 5.5 3 – 20 1%

Also for control, apply 8.9 to 17 ounces of this product plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Cogongrass 3.2 – 5.5 10 – 40 1%

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dallisgrass² 3.2 – 5.5 3 – 20 1%

Dandelion¹ 3.2 – 5.5 3 – 40 1%

Also for control, apply 8.9 ounces of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dock Curly 3.2 – 5.5 3 – 40 1%

Also for control, apply 8.9 to 17 ounces of this product plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dogbane, hemp 4.4 3 – 40 1%

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall.

For suppression, apply 8.8 ounces of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

Deleted: 1
Deleted: March 19,

75/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | | |
|-----------------------------------|-----------|--------|----|
| Fescue (except tall) ² | 3.2 – 5.5 | 3 – 20 | 1% |
|-----------------------------------|-----------|--------|----|

| | | | |
|--------------|-----------|--------|----|
| Fescue, tall | 1.1 – 3.2 | 3 – 40 | 1% |
|--------------|-----------|--------|----|

Apply 52 ounces of this product per acre when most plants have been reached boot-to-early seedhead stage of development.

Fall applications only: Apply 17 ounces of this product in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 8.9 ounces per acre of this product will improve long-term control and control seedlings germinating after fall treatments or the following spring.

| | | | |
|-------------|-----|--------|------|
| Guineagrass | 3.2 | 3 – 40 | 0.6% |
|-------------|-----|--------|------|

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 35 ounces for control. In the flatwoods region of Florida, 52 ounces is required for control.

| | | | |
|--------------------------|-----------|--------|----|
| Horsenettle ¹ | 3.2 – 5.5 | 3 – 20 | 1% |
|--------------------------|-----------|--------|----|

| | | | |
|-------------|-----|--------|----|
| Horseradish | 4.4 | 3 – 40 | 1% |
|-------------|-----|--------|----|

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

| | | | |
|-----------------------|----|----|----------|
| Iceplant ¹ | -- | -- | 0.7 – 1% |
|-----------------------|----|----|----------|

Thorough coverage is necessary for best control.

| | | | |
|----------------------------------|-----------|--------|----|
| Jerusalem artichoke ¹ | 3.2 – 5.5 | 3 – 20 | 1% |
|----------------------------------|-----------|--------|----|

| | | | |
|--------------|-----------|--------|------|
| Johnsongrass | 0.6 – 3.2 | 3 – 40 | 0.6% |
|--------------|-----------|--------|------|

In annual cropping systems apply 17 to 35 ounces of this product per acre. Apply 17 ounces of this product in 3 to 10 gallons of water per acre. Use 35 ounces of this product when applying 10 to 40 gallons of water per acre. In non-crop, or areas where annual tillage (no-till) is not practiced, apply 35 to 52 ounces of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 17 ounces of this product per are.

For burndown of Johnsongrass, apply 8.9 ounces of this product in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

76/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Spot treatment (partial control or suppression) – Apply a 0.6 percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage must be uniform and complete.

Kikuyugrass 2.2 – 3.2 3 – 40 1%

Spray when most kikuyugrass is at least 8 inches in height (3 or 4 leaf stage of growth). Allow 3 or more days after application before tillage.

Knapweed 4.4 3 – 40 1%

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

Lantana -- -- 0.7-1%

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

Lespedeza¹ 3.2 – 5.5 3 – 20 1%

Milkweed, common 3.2 3 – 40 1%

Apply when most plants have reached the late bud to flower stage of growth.

Muhly, wirestem 1.1 – 2.2 3 – 40 1%

Use 17 ounces of this product in 3 to 10 gallons of water per acre. Use 35 ounces of this product when applying 10 to 40 gallons of water per acre or in pasture, sod or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage.

Mullein, common¹ 3.2– 5.5 3 – 20 1%

Napiergrass² 3.2– 5.5 3 – 20 1%

Nightshade, silverleaf 2.2 3 – 10 1%

Applications must be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost.

Nutsedge,
Purple or yellow 0.6 – 3.2 3 – 40 0.6-1%

Apply 52 ounces of this product per acre or apply a 0.6 to 1 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in

Deleted: 1
Deleted: March 19,

77/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.

Sequential applications: 17 to 35 ounces of this product in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3 to 5 leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5 leaf stage. Subsequent applications will be necessary for long-term control.

For partial control of existing plants, apply 8.9 to 35 ounces of this product in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

| | | | |
|--------------|-----------|--------|----|
| Orchardgrass | 1.1 – 2.2 | 3 – 40 | 1% |
|--------------|-----------|--------|----|

Apply 35 ounces of this product in 10 to 40 gallons of water per acre when most plants have received boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 17 to 26 ounces of this product in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn. Apply 17 to 26 ounces of this product in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

| | | | |
|-------------|-----------|---------|----------|
| Pampasgrass | 3.2 – 5.5 | 10 – 40 | 0.7 – 1% |
|-------------|-----------|---------|----------|

Pampasgrass must be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

| | | | |
|------------------------|-----------|--------|----|
| Paragrass ² | 3.2 – 5.5 | 3 – 20 | 1% |
|------------------------|-----------|--------|----|

| | | | |
|------------|-----------|---------|--------|
| Phragmites | 3.2 – 5.5 | 10 – 40 | 0.6-1% |
|------------|-----------|---------|--------|

For partial control, and best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

| | | | |
|----------------|-----------|---------|----------|
| Poison hemlock | 2.2 – 4.4 | 10 – 40 | 0.6 – 1% |
|----------------|-----------|---------|----------|

For hand held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Thorough coverage is necessary for best control.

Deleted: 1
Deleted: March 19,

78/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Pokeweed, common 1.6 3 – 40 1%

Apply to actively growing plants up to 24 inches tall.

Quackgrass 1.1 – 3.2 3 – 40 1%

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 17 ounces of this product in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 35 ounces of this product. Do not tank mix with residual herbicides when using the 17 ounce rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.

In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 35 to 52 ounces of this product in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

Redvine 0.8 – 2.2 5 – 10 1%

For suppression, apply 13 ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 35 ounces per acre. Apply listed rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Reed, giant 4.4 – 5.5 10 – 40 1%

Best results are obtained when applications are made in late summer to fall.

Ryegrass, perennial 1.1 – 3.2 3 – 40 0.6%

In annual cropping systems apply 17 to 35 ounces of this product per acre. Apply 17 ounces of this product in 3 to 10 gallons of water per acre. Use 35 ounces of this product when applying 10 to 40 gallons of water per acre. In non-crop, or areas where annual tillage (no-till) is not practiced, apply 35 to 52 ounces of this product in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1.6 pounds of this product per acre.

Smartweed, swamp¹ 3.2 – 5.5 3 – 40 1%

Also, for control, apply 8.9 ounces of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall.

Sowthistle, perennial 2.4 – 3.2 3 – 40 1%

Deleted: 1
Deleted: March 19,

79/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

Spurge, leafy -- 3 - 10 1%

For suppression, apply 8.9 ounces of this product plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall.

Starthistle, yellow 2.2 10 - 40 1%

Best results are obtained when applications are made during the rosette, bolting and early flower stages.

Sweet potato, wild -- -- 1%

For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.

Thistle, artichoke 2.2 - 3.2 10 - 40 0.6 - 1%

For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.

Thistle, Canada 2.2 - 3.2 3 - 40 1%

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression in the spring, apply 17 ounces of this product, or 8.9 ounces of this product plus 0.5 pound of 2,4-D, in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

Timothy² 2.2 - 3.2 3 - 40 1%

Torpedograss 4 - 5.5 3 - 40 1%

For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Trumpet creeper 2.2 – 3.2 5 – 10 1%

For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Vaseygrass² 3.2 – 5.5 3 – 20 1%

Velvetgrass² 3.2 – 5.5 3 – 20 1%

Wheatgrass, western² 2.2 – 3.2 3 – 40 1%

¹ Apply when most plants have reached the early bud stage of growth.

² Apply when most plants have reached the early heading stage of growth.

15.0 WOODY BRUSH AND TREES RATE SECTION

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Unless otherwise directed, apply broadcast treatments in 3 to 40 gallons of water per acre. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

WOODY BRUSH AND TREES RATE TABLE

| Weed Species | (Alphabetically by Species) | |
|-----------------------------------|-----------------------------|----------------------|
| | Rate (LBS/A) | Hand-Held % Solution |
| Alder | 3.2 – 4.4 | 0.6 – 1% |
| Ash ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Aspen, quaking | 2.2 – 3.2 | 0.6 – 1% |
| Bearmat (Bearclover) ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Beech ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Birch | 2.2 | 0.6% |
| Blackberry | 3.2 – 4.4 | 0.6 – 1% |

Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 0.6 percent solution of this product.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3.2 to 4.4 pounds of this product in 10 to 40 gallons of water per acre.

| | | |
|--------------------------------------|-----------|------------|
| Blackgum | 2.2 - 5.5 | 0.6 - 1% |
| Bracken | 2.2 - 5.5 | 0.6 - 1% |
| Broom, French, Scotch | 2.2 - 5.5 | 0.7 - 1% |
| Buckwheat, California ^{1,2} | 2.2 - 4.4 | 0.6 - 1% |
| Cascara ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Catsclaw ¹ | -- | 0.6 - 0.7% |
| Ceanothus ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Chamise ² | 2.2 - 5.5 | 0.6% |
| Cherry; butter, black, pin | 2.2 - 3.2 | 0.6 - 0.7% |
| Coyote brush | 3.2 - 4.4 | 0.7 - 1% |

Apply when at least 50 percent of the new leaves are fully developed.

| | | |
|----------------------|-----------|----------|
| Dogwood ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Elderberry | 2.2 | 0.6% |
| Elm ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Eucalyptus | -- | 1% |

For control of eucalyptus resprouts apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.

| | | |
|--|-----------|----------|
| Florida holly (Brazilian Peppertree) ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Gorse ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Hasardia ^{1,2} | 2.2 - 4.4 | 0.6 - 1% |
| Hawthorn | 2.2 - 3.2 | 0.6 - 1% |
| Hazel | 2.2 | 0.6% |
| Hickory ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Honeysuckle | 3.2 - 4.4 | 0.6 - 1% |
| Hornbeam, American ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Kudzu | 4.4 - 5.5 | 1% |

Repeat applications may be required to maintain control

| | | |
|--------------------------------|-----------|----------|
| Locust, black ¹ | 2.2 - 4.4 | 0.6 - 1% |
| Madrone resprouts ¹ | -- | 1% |

Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

| | | |
|------------------------|-----------|----------|
| Manzanita ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Maple, red | 2.2 - 4.4 | 0.6 - 1% |

Apply a 0.6 to 0.7 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 35 to 70 ounces of this product per acre.

| | | |
|--------------|--|----------|
| Maple, sugar | | 0.6 - 1% |
|--------------|--|----------|

Apply when at least 50 percent of the new leaves are fully developed.

| | | |
|--------------------------------|-----------|----------|
| Monkey flower ^{1,2} | -- | 0.6 - 1% |
| Oak; black, white ¹ | 2.2 - 4.4 | 0.6 - 1% |
| Oak, post | 3.2 - 4.4 | 0.6 - 1% |
| Oak; northern, pin | 2.2 - 4.4 | 0.6 - 1% |

Apply when at least 50 percent of the new pin leaves are fully developed.

| | | |
|------------------------|-----------|----------|
| Oak; southern red | 2.2 - 3.2 | 0.6 - 1% |
| Persimmon ¹ | 2.2 - 5.5 | 0.6 - 1% |
| Pine | 2.2 - 5.5 | 0.6 - 1% |
| Poison ivy/Poison oak | 4.4 - 5.5 | 1% |

Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

82/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | |
|---|-----------|-----------|
| Poplar, yellow ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Redbud, eastern | 2.2 – 5.5 | 0.6 – 1% |
| Rose, multiflora | 2.2 | 0.6% |
| Treatments must be made prior to leaf deterioration by leaf-eating insects. | | |
| Russian olive ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Sage, black ² | -- | 0.6 % |
| Sage, white ¹ | 2.2 – 5.5 | 0.6 – 1 % |
| Sage brush, California ² | -- | 0.6% |
| Salmonberry | 2.2 | 0.6% |
| Salt-cedar | 2.2 – 5.5 | 0.6 – 1% |
| Sassafras ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Sourwood ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Sumac; poison, smooth, winged ¹ | 2.2 – 4.4 | 0.6 – 1% |
| Sweetgum | 2.2 – 3.2 | 0.6 – 1% |
| Swordfern ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Tallowtree, Chinese ² | -- | 0.6% |
| Tan oak resprouts ¹ | -- | 1% |

Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.

| | | |
|----------------------------------|-----------|----------|
| Thimbleberry | 2.2 | 0.6% |
| Tobacco, tree ¹ | -- | 0.6 – 1% |
| Trumpet creeper | 2.2 – 3.2 | 0.6 – 1% |
| Vine maple ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Virginia creeper | 2.2 – 5.5 | 0.6 – 1% |
| Waxmyrtle, southern ¹ | 2.2 – 5.5 | 0.6 – 1% |
| Willow | 3.2 | 0.6% |

¹ Partial Control

² Thorough coverage of foliage is necessary for best results.

16.0 WEED RESISTANT MANAGEMENT

Based on the mode of action classification system of the Weed Science Society of America, Glyphosate (active ingredient) is a Group 9 herbicide. Group 9 herbicides may contain plants from any weed population that can be naturally resistant to glyphosate. These weed resistant plants can be effectively controlled using a different Group herbicide or by using other means such as cultural or mechanical practices.

16.1 Weed Resistance Management Recommendations

Glyphosate resistant biotypes can be minimized by utilizing the following weed resistance management recommendations:

- 1) Begin by preparing your field using tillage or a burndown herbicide application.
- 2) It is a good practice to scout your fields before and after applications.
- 3) Use new commercial seeds that have fewer weed seeds.
- 4) Control weeds early when they are relatively small.
- 5) Cultural practices such as crop rotation or tillage and the addition of other herbicides such as a selective and/or a residual herbicide where appropriate are suggested.
- 6) Rotating to other Roundup Ready crops is one method for adding other herbicides into a continuous Roundup Ready system.

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- 7) Follow the specified label rate for the most difficult to control weeds. Reject directions that support lower application rates when tank mixing as well as tank mixtures with other herbicides that will reduce product efficacy.
- 8) Follow good agricultural practices by cleaning equipment prior to shifting from field to field preventing weed seed or plant root parts from spreading.
- 9) Any incidence of repeated non-performance of this product on a particular weed must be reported to any MEY Corporation representative, your county extension agent, or to the local retailer.

16.2 Glyphosate-Resistant Biotypes Management

In order to reduce the spread of confirmed glyphosate resistant biotypes, apply the following practices:

- 1) When a naturally occurring resistant biotype(s) is present, tank mix or apply sequentially with an appropriate herbicide with a different mode of action to achieve control.
- 2) Use cultural and mechanical control practices, such as crop rotation or tillage, as appropriate.
- 3) Rotation to other Roundup Ready crops is one method for adding other herbicides into a continuous Roundup Ready system.
- 4) Control escaping weeds including resistant biotypes before they set seed and scout treated fields after herbicide application.
- 5) Clean equipment thoroughly prior to exiting fields known to contain resistant biotypes.

MEY Corporation is not responsible for any losses that may result from the failure of this product to control glyphosate-resistant weed biotypes as the occurrence of new glyphosate-resistant weeds cannot be determined until after product use and scientific confirmation.

16.3 Control and Management of Glyphosate-Resistant Horseweed in Cotton, Corn, and Soybean

For ground applications, use 10 to 20 gallons of water per acre. For aerial applications, use 3 to 15 gallons of water per acre. For tank-mix instructions, read and carefully observe the cautionary statements and all other information appearing on the product labels, supplemental labeling or fact sheets published separately for all herbicides used.

Cotton:

Preplant:

For control of horseweed, apply this product (17 ounces per acre) in a tank-mix with Clarity® (8 fluid ounces per acre). This application must be made 21 to 35 days before planting and before horseweed reaches 6 inches in height. In order to avoid crop injury, a minimum interval of 21 days during which there is at least 1 inch of cumulative rainfall must be observed between Clarity application and planting of cotton.

Post-directed (Roundup Ready® Cotton varieties only):

Management of early season weed competition and the development of a crop height differential between cotton and the horseweed is often achieved by a combination of preplant burndown and postemergent over-the-top and/or directed applications. These measures enhance the development of a height differential that is necessary to successfully make post-directed treatments. In-cop post-directed applications of MSMA (2 pounds active ingredient per acre) tank-mixed with diuron (0.5 to 0.75 pounds active ingredient per acre) must be made when the temperature is 80° F or higher.

Deleted: 1

Deleted: March 19,

84/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Soybeans:

Preplant:

Apply a tank mixture of this product (17 ounces per acre) with 2,4-D (0.5 pounds active ingredient per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting. For areas where 2,4-D cannot be applied due to application restrictions or proximity to a sensitive crop, contact your local retailer and/or crop consultant.

In-crop (Roundup Ready Soybean varieties only):

Horseweed must be controlled prior to planting using preplant burndown treatments. In-crop Roundup Ready soybeans, apply a tank mixture of this product (17 ounces per acre) with Amplify™ (0.3 ounces per acre). This treatment must be used as a salvage treatment only for a horseweed infestation that was not controlled preplant. Application must be made between full emergency of the first trifoliolate leaf and 50 percent flowering stage or soybeans. At the time of treatment, horseweed must not exceed 6 inches in height.

Corn:

Preplant, At-Planting, Preemergence:

Apply a tank mixture of this product (17 ounces per acre) plus 2,4-D (0.5 pounds active ingredient of per acre) before horseweed exceeds 6 inches in height. See the 2,4-D product label for time intervals that are required between application and planting.

Atrazine (1 to 2 pounds active ingredient per acre) may be included in the tank mixture to provide residual control. Refer to the atrazine product label for specific use instructions.

In-crop (Roundup Ready Corn hybrids only):

In crop Roundup Ready corn, apply a tank mixture with this product (17 ounces per acre) plus Clarity (8 to 16 fluid ounces per acre) or 2,4-D (0.5 to 1.0 pound active ingredient per acre). Apply between corn emergence and the 5-leaf stage of growth (approximately 8 inches tall).

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

16.0 LIMIT OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, MEY Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESSED WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

To the extent consistent with applicable law buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop, or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

To the extent consistent with applicable law upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Amplify, Bullet, Degree, Harness, Lariat, Lasso, Micro-Tech, Monsanto and Vine symbol, Permit, Roundup UltraMAX, Roundup WeatherMAX and TransSorb are trademarks of Monsanto Technology LLC.

Assure, Authority, Canopy, Krovar, Leadoff and Lexone are trademarks of E.I. DuPont de Nemours & Co. Inc.

Direx, Karmex, Linex and Lorox are trademarks of Griffin LLC.

Bicep MAGNUM, Dual MAGNUM, Flexstar, Fusion, Princep Caliber, Reflex and Solicam are trademarks of Syngenta CropProtection Inc.

Firstrate, Fultime, Python, Surflan, Topnotch and Tordon are trademarks of Dow AgroSciences LLC.

Banvel, Clarity, Distinct, Frontier, Guardsman, Marksman, Outlook, Prowl, Pursuit Plus, Scepter, Squadron and Steel are trademarks of BASF Corp.

Balance, Folex, Ginstar, and Prep are trademarks of Aventis CropSciences.

Goal is a trademark of Rohm and Haas Company.

Deleted: 1
Deleted: March 19.

86/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Axiom, DEF, Domain, Epic, Sencor and Turbo are trademarks of Bayer Crop Protection.

Aim, Command and Gauntlet are trademarks of FMC Corporation.

Devrinol is a trademark of United Phosphorus Inc.

Sim-Trol is a trademark of Sipcam Agro USA, Inc.

Valor and Resource are trademarks of Valent USA Corporation

[[Revision] [Version Date] [MM/YYYY]]

Deleted: 1
Deleted: March 19,

87/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

(Note to Reviewer: The following information will be affixed to the unit package/container.)

GlySupreme 75-WDG

[Water Soluble Dry Herbicide] [Water Dispersible Granule [Herbicide]]

(Alternate Brand Names: GlySupreme 75-Water Dispersible Granule, GlySupreme 75-WDG Herbicide, GlySupreme 75-WDG Glyphosate Herbicide)

Formatted: Centered

This product is a post-emergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush, and trees. It is formulated as a water soluble granule. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions. Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees because severe injury or destruction may result.

ACTIVE INGREDIENT:

| | |
|--|--------|
| *Glyphosate, N-(phosphonomethy) glycine, in the form of its ammonium salt..... | 75.7% |
| Other Ingredients | 24.3% |
| Total | 100.0% |

*Equivalent to 68.7% of glyphosate acid

1 pound of [this product] [GlySupreme 75-WDG] contains 0.687 pounds of glyphosate acid.

EPA Reg. No.: 80967-13
EPA Est. No.: [73432-CHN-001]
Net Contents:

Manufactured [for][by]:
MEY Corporation
121 S. Estes Drive, Suite 101
Chapel Hill, NC 27514 USA

[Lot/Batch code/number]
(Note to reviewer: Lot or Batch number may appear on label or printed directly on packaging.)

KEEP OUT OF REACH OF CHILDREN CAUTION

See attached booklet for [additional precautionary statements] [first aid] [complete directions for use] [and warranty]

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE (800) 613-0427

IN CASE OF AN EMERGENCY INVOLVING THIS HERBICIDE PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL (800) 262-8200

| FIRST AID | |
|----------------------|---|
| If swallowed: | Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce |

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | |
|---|---|
| | vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If inhaled: | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. |
| If in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-262-8200. | |

Read the entire label before using this product. Use only according to label instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Caution. Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist or vapor.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE): Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

(Note to Reviewer: For products in containers >50 lbs): Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System

Deleted: 1
Deleted: March 19,

89/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

(NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use to prevent spills and contamination.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal Facility.

Container Handling: Non-refillable container. Do not reuse or refill this container.

(Note to reviewer: For non-refillable plastic containers (≤50 lbs) small enough to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable plastic containers (>50 lbs) too large to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable paper and plastic bags, large fiber sacks):

Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable fiber drums with liners or bag-in-box packaging):

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke. If [drum] [box] [carton] is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Deleted: 1
Deleted: March 19,

90/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

WARRANTY AND LIABILITY

To the extent consistent with applicable law, MEY Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESSED WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

[[Revision] [Version Date] [MM/YYYY]]

Deleted: 1
Deleted: March 19,

911/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

SUBLABEL B: Industrial, Turf, and Ornamental Use

GlySupreme 75-WDG

[Water Soluble Dry Herbicide] [Water Dispersible Granule [Herbicide]]

(Alternate Brand Names: GlySupreme 75-Water Dispersible Granule, GlySupreme 75-WDG Herbicide, GlySupreme 75-WDG Glyphosate Herbicide)

This product contains surfactants and ammonium sulphate in the granule resulting in superior weed control.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees because severe injury or destruction may result.

Read the entire label before using this product. Use only according to label instructions.

CONTENTS

- 1 1.0 INGREDIENTS AND FRONT PANEL STATEMENTS
- 2 2.0 IMPORTANT PHONE NUMBERS
- 3 3.0 PRECAUTIONARY STATEMENTS
 - 3.1 Hazards to Humans and Domestic Animals
 - 3.2 Environmental Hazards
 - 3.3 Physical or Chemical Hazards
 - 3.4 Directions for Use/WPS
- 4 4.0 STORAGE AND DISPOSAL
- 5 5.0 INFORMATION
 - (How This Product Works)
- 6 6.0 MIXING
 - 6.1 Mixing with Water
 - 6.2 Tank Mixing Procedure
 - 6.3 Mixing Percent Solutions
 - 6.4 Colorants or Dyes
- 7 7.0 APPLICATION EQUIPMENT AND TECHNIQUES
 - 7.1 Aerial Equipment
 - 7.2 Ground Broadcast Equipment
 - 7.3 Hand-Held Equipment
 - 7.4 Selective Equipment
 - 7.5 Injection Systems
 - 7.6 CDA Equipment
- 8 8.0 SITE AND USE INSTRUCTIONS
 - 8.1 Cut Stump
 - 8.2 Forestry Site Preparation

Deleted: 1

Deleted: March 19,

92/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- 8.3 Non-crop Areas and Industrial Sites
 - 8.4 Habitat Management
 - 8.5 Injection and Frill (Woody Brush and Trees)
 - 8.6 Ornamental, Plant Nurseries, and Christmas Trees
 - 8.7 Parks, Recreational and Residential Areas
 - 8.8 Railroads
 - 8.9 Roadsides
 - 8.10 Utility Sites
- 9 9.0 WEEDS CONTROLLED
- 9.1 Annual Weeds
 - 9.2 Perennial Weeds
 - 9.3 Woody Brush and Trees
- 10 LIMIT OF WARRANTY AND LIABILITY

1.0 INGREDIENTS AND FRONT PANEL STATEMENTS

ACTIVE INGREDIENT:

| | |
|--|--------|
| *Glyphosate, N-(phosphonomethy) glycine, in the form of its ammonium salt..... | 75.7% |
| Other Ingredients | 24.3% |
| Total | 100.0% |

*Equivalent to 68.7% of glyphosate acid

1 pound of [this product] [GlySupreme 75-WDG] contains 0.687 pounds of glyphosate acid.

EPA Reg. No.: 80967-13

EPA Est. No.: [73432-CHN-001]

Net Contents:

Manufactured [for][by]:
 MEY Corporation
 121 S. Estes Drive, Suite 101
 Chapel Hill, NC 27514 USA

[Lot/Batch code/number]

(Note to reviewer: Lot or Batch number may appear on label or printed directly on packaging.)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See attached booklet for [additional precautionary statements] [first aid] [complete directions for use] [and warranty]

2.0 IMPORTANT PHONE NUMBERS

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE (800) 613-0427

IN CASE OF AN EMERGENCY INVOLVING THIS HERBICIDE PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL (800) 262-8200

Deleted: 1

Deleted: March 19,

93/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

3.0 PRECAUTIONARY STATEMENTS

| FIRST AID | |
|---|---|
| If swallowed: | Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If inhaled: | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. |
| If in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on skin or clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-262-8200. | |

3.1 Hazards to Humans and Domestic Animals

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Caution. Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist or vapor.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE): Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

94/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

3.2 Environmental Hazards

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

(Note to Reviewer: For products in containers >50 lbs): Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

3.3 Physical or Chemical Hazards

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

3.4 Directions for Use/WPS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Deleted: 1
Deleted: March 19,

95/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Do not allow worker entry into treated areas during the restricted entry interval (REI) of four (4) hours or until solution has dried.

Exception: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter treated area if there is no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is: coveralls, chemical resistant gloves (made of any waterproof material) and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of the product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled.

Deleted: 1

Deleted: March 19,

96/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

4.0 STORAGE AND DISPOSAL

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use to prevent spills and contamination.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal Facility.

Container Handling: Non-refillable container. Do not reuse or refill this container.

(Note to reviewer: For non-refillable plastic containers (≤50 lbs) small enough to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable plastic containers (>50 lbs) too large to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable paper and plastic bags, large fiber sacks):

Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable fiber drums with liners or bag-in-box packaging):

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke. If [drum] [box] carton is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Deleted: 1

Deleted: March 19,

97/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

5.0 INFORMATION (HOW THIS PRODUCT WORKS)

Product Description: This product is a postemergent, systemic herbicide with no soil residual activity. It gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble granule. No additional surfactant is needed.

Time to Symptoms: This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Mode of Action in Plants: The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the listed stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash this product off of the foliage and a repeat application may be required for adequate control.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Tank Mixing: This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly listed in this label. Mixing this product with herbicides or other materials not listed on this label may result in reduced performance.

Grazing Restrictions: This product may be used to treat undesirable vegetation in rights-of-way that pass through pastures, rangeland and forestry sites that are being grazed. For tank mix applications, comply with all restrictions appearing on the tank mix product label.

There are no grazing restrictions for the following labeled applications of this product:

- Where the spray can be directed onto undesirable woody brush and trees, such as in handgun spray-to-wet or low volume directed spray treatments.
- For tree injection or frill applications and for cut stump treatments.

For broadcast applications, observe the following restrictions:

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- For application rates of greater than 5.6 but not to exceed 9.3 quarts per acre, no more than 15 percent of the available grazing area may be treated.
- For application rates that do not exceed 5.6 quarts per acre, no more than 25 percent of the available grazing area may be treated.
- All restrictions outlined above apply to lactating dairy animals. No other restrictions apply to lactating dairy animals.

These directions do not apply to rangeland outside of rights-of-way

Annual Maximum Use Rate: The combined total of all treatments must not exceed 11.6 pounds of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rates.

ATTENTION

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS (EXCEPT AS SPECIFIED FOR INDIVIDUAL ROUNDUP READY CROPS), DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combination of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

6.0 MIXING

Clean sprayer parts immediately after using this product by thoroughly flushing with water.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

6.1 Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows:

For hand held or backpack sprayers (less than or equal to 5 gal. capacity): Add the listed amount of this product to the spray tank. If adding Ammonium Sulfate, pre-dissolve in water before adding. Fill the spray tank with water and ensure thorough mixing. Alternatively, the

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

99/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

listed amount of this product can be mixed with water in a large container. Fill sprayer with the mixed solution.

For larger tank sprayers (greater than 5 gal. capacity): Fill the mixing or spray tank one-half full with water and start agitation. If adding Ammonium Sulfate, ensure that it is completely dissolved before proceeding. Add the listed amount of this product using a circular motion while pouring. Continue filling the spray tank with water and ensure thorough mixing.

Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

6.2 Tank Mixing Procedure

Use the following procedure to mix this product when preparing tank mixtures with other labeled products.

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If adding Ammonium Sulfate ensure that it is completely dissolved before proceeding.
4. Add [GlySupreme 75-WDG] [herbicide] using a circular motion while pouring.
5. If second product is a wettable powder, first make a slurry with the water carrier, then add the slurry SLOWLY through the screen into the tank. Continue agitation.
6. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
7. If an emulsifiable concentration formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
8. Continue filling the spray tank with water and add water soluble liquids near the end of the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift reduction additive and water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers must be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Refer to the "TANK MIXING" section of "INFORMATION" for additional precautions.

Deleted: 1
Deleted: March 19,

100/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

6.3 Mixing Percent Solutions

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table.

Percent Solution Amount of GlySupreme 75-WDG
% Weight to Weight *

| | 0.3% | 0.6% | 0.8% | 1.1% | 2.8% | 5.5% | 11.1% |
|-----------------------|--------|--------|--------|--------|---------|---------|---------|
| Desired Volume | | | | | | | |
| 1 Gal | 0.4 oz | 0.7 oz | 1 oz | 1.4 oz | 3.4 oz | 6.8 oz | 13.7 oz |
| 3 Gal | 1.1 oz | 2.2 oz | 3 oz | 4.1 oz | 10.4 oz | 1.3 lb | 2.6 lb |
| 25 Gal | 9.3 oz | 1.2 oz | 1.6 lb | 2.1 lb | 5.4 lb | 10.7 lb | 21.4 lb |
| 100 Gal | 2.3 lb | 4.6 lb | 6.2 lb | 8.5 lb | 21.7 lb | 42.6 lb | 86.0 lb |

*These amounts are calculated on a weight of this product per weight of final spray solution.

Above percentages are on a weight-to-weight basis with water as an 8.34 pound gallon. A 0.6% solution of GlySupreme 75-WDG herbicide contains approximately the same amount of glyphosate acid as a 1% volume-to-volume solution of Roundup Pro herbicide.

6.4 Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilution. Use colorants or dyes according to the manufacturer's recommendations. Certain blue dyes are not stable in the spray solution in the presence of this product. A jar test may be needed to determine if the desired blue dye is stable. If stability is a problem, consider switching to an alternate color dye.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all of these factors when making decisions.

Deleted: 1
Deleted: March 19,

101/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

Importance of droplet size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversions" sections of this label).

Controlling droplet size

- Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce large droplets.
- Pressure: Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles: Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation: Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- Boom Length: For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.
- Application Height: Application must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporate and wind.

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance must increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application must be avoided below 2 miles per hour due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sunsets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive areas

This product must only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

FOR AERIAL APPLICATION IN CALIFORNIA, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

This product plus, dicamba tank mixtures may not be applied by air in California.

Deleted: 1

Deleted: March 19,

103/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Use the listed rates of this herbicide in 3 to 25 gallons of water per acre.

Drift reduction additives may be used. When a drift reduction additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application – To avoid streaked, uneven or overlapped application, use appropriate marking devices.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear is most susceptible.

7.2 Ground Broadcast Equipment

For broadcast ground applications, unless otherwise specified use this product at the rate of 1.1 to 2.1 lbs. per acre for annual weeds, 2.1 to 5.5 lbs. per acre for perennial weeds and 4.3 to 10.7 lbs. per acre for woody brush and trees. When used according to label directions this product will give control or partial control of herbaceous weeds, woody brush and trees listed in the "WEEDS CONTROLLED" section of this label.

Apply the listed rate in 3 to 40 gallons per acre for best results. Broadcast applications with hand-held spray gun may require spray volumes higher than 40 gallons per acre for uniform coverage. As density of herbaceous weeds and woody brush increases, spray volume must be increased as needed to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. Check for even distribution of spray droplets.

7.3 Hand-Held Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For control of weeds listed in the "ANNUAL WEEDS" section of "WEEDS CONTROLLED", apply a 0.3 percent solution of this product to weeds less than 6 inches in height or runner length. For annual weeds over 6 inches tall, or unless otherwise specified, use a 0.6 percent solution. Make applications to annuals prior to seedhead emergence in grasses or bud formation in broadleaf weeds.

Use a 1 percent solution as a spray to wet application on harder-to-control perennials, woody vines, brush and trees. Make applications to perennials after seedhead emergence in grasses or bud formation in broadleaf weeds, woody brush and trees for best results.

For low volume directed spray applications, use a 2.8 to 5.5 percent solution of this product for control or partial control of annual weeds, perennial weeds, or woody brush and trees.

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Spray coverage must be uniform with at least 50 percent of the foliage contacted. Coverage of the top one half of the plant is important for best results. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

Unless otherwise specified, use the rates listed in the following "APPLICATION RATES" table for various methods of foliar application using high volume, backpack, knapsack and similar types of hand-held equipment. When used according to label directions this product will give control or partial control of herbaceous weeds, woody brush and trees listed in the "WEEDS CONTROLLED" section of this label.

APPLICATION RATES

| APPLICATION | GLYSUPREME 75-WDG | SPRAY VOLUME GALLONS/ACRE |
|---|----------------------|---------------------------|
| <u>SPRAY-TO-WET</u> | | |
| Handgun, or Backpack | 0.3 to 1% by weight | Spray-to-wet* |
| <u>LOW VOLUME DIRECTED SPRAY</u> | | |
| Backpack | 2. to 5.1% by weight | 15 to 25** |
| Modified High Volume | 1 to 2.1% by weight | 40 to 60** |

*For applications made on a spray-to-wet basis, spray coverage must be uniform and complete. Do not spray to the point of runoff.

**For low volume directed spray applications, coverage must be uniform with at least 50 to 75 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results. Low volume directed applications with backpacks work best when treating weeds and brush less than 10 feet tall. For taller weeds and brush, high volume handguns can be modified by reducing nozzle size and spray pressure to produce a low volume directed spray. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop.

7.4 Selective Equipment

This product may be applied through recirculating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, after dilution and thorough mixing with water, to listed weeds growing in any non-crop site specified on this label.

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION, AS SERIOUS INJURY OR DEATH IS LIKELY TO OCCUR.

Applicators used above desired vegetation must be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation is likely to result in discoloration, stunting or destruction.

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

105/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

Shielded and Hooded Applicators

A shielded or hooded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Wiper Applicators and Sponge Bar

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if two applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

For Rope or Sponge Wick Applicators – Prepare solutions by dissolving 1.4 to 3.2 pounds of this product per gallon of water.

For Panel Applicators and pressure-feed systems – Prepare solutions by dissolving 1.4 to 3.7 pounds of this product per gallon of water.

Note: In preparing these concentrated solutions always allow adequate time for product to dissolve. Use of warm water will shorten dissolution time.

7.5 Injection Systems

This product may be used in aerial or ground injection spray systems. This product may be injected into the spray stream after dilution and thorough mixing with water. Do not mix this product with the undiluted concentrate of other products when using injection systems unless specifically directed.

Deleted: 1

Deleted: March 19,

106/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

7.6 CDA Equipment

The rate of this product applied per acre by controlled droplet application (CDA) equipment must not be less than the amount listed in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units – Apply an 11 percent solution of this product (13.7 oz of product per gallon) at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 miles per hour (1 quart per acre). For the control of perennial weeds, apply an 11 to 22 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 miles per hour (2 to 4 quarts per acre).

CDA equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction is likely to result.

8.0 SITE AND USE INSTRUCTIONS

Detailed instructions follow alphabetically, by site.

Unless otherwise specified, applications may be made to control any weeds listed in the annual, perennial and woody brush tables. Refer also to the "SELECTIVE EQUIPMENT" section.

8.1 Cut Stump

Cut stump treatments may be made on any site listed on this label. This product will control many types of woody brush and tree species. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a solution prepared by dissolving 2.1 to 3.7 pounds of this product per gallon of water to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications must be made during periods of active growth and full leaf expansion.

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots.

Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

8.2. Forestry Site Preparation

This product can be used for the control or partial control of woody brush, trees and herbaceous weeds in forestry. This product can also be used in preparing or establishing wildlife openings within these sites and maintaining logging roads.

This product can be used in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

For applications using different types of equipment, see "APPLICATION RATES" table in "HAND-HELD EQUIPMENT" section of this label.

TANK MIXTURES

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements of all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any listed rate of this product may be used in a tank mix with the following products for forestry site preparation.

| PRODUCT | BROADCAST RATE |
|---------------------------------|--------------------------------|
| Arsenal Applicators Concentrate | 2. to 16 fluid ounces per acre |
| Escort™ | 0.5 to 3.5 ounces per acre |
| Chopper™ | 4 to 32 fluid ounce per acre |
| Garlon 4 | 1 to 4 quarts per acre |

| | |
|-------|------------------------|
| Oust™ | 1 to 4 ounces per acre |
|-------|------------------------|

| PRODUCT | SPRAY-TO-WET RATES |
|---------------------------------|-------------------------------|
| Arsenal Applicators Concentrate | 0.03 to 0.5 percent by volume |

| PRODUCT | LOW VOLUME DIRECTED SPRAY RATES |
|---------------------------------|---------------------------------|
| Arsenal Applicators Concentrate | 0.1 to 0.5 percent by volume |

For control of herbaceous weeds, use the lower tank mixture rate. For control of dense stands or tough-to-control woody brush and trees, use the higher rate.

Unless otherwise directed, do not apply this product as an over-the-top broadcast spray for forestry conifer or hardwood release.

8.3 Non-crop Areas and Industrial Sites

Use in areas including airports, apartment complexes, ditch banks, dry ditches, dry canals, fencerows, golf courses, industrial sites, lumber yards, manufacturing sites, office complexes, parks, parking areas. Petroleum tank farms and pumping installations, railroads, recreational areas, residential areas, roadsides, sod or turf seed farms, schools, storage areas, substations, warehouse areas, other public areas, and industrial and non-crop sites.

Weed control, Trim-and-edge and Bare ground

This product may be used in non-crop areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. This product

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product may be tank mixed with the following products. Refer to these products labels for approved non-crop sites and application rates.

- | | |
|-----------------|----------------|
| Arsenal™ | Plateau™ |
| Clarity | Princep™DF |
| Barricade™65 WG | Princep™Liquid |
| Duron | Ronstar™50WP |
| Endurance™ | Sahara™ |
| Escort™ | Simazine |
| Garlon™ 3A | Spike 80DF |
| Hyvar X | Surflan™ |
| Garlon 4 | Telar™ |
| Karmex™DF | Vanquish™ |
| Krovar™ 1DF | 2,4-D |
| Manage® | |
| Oust | |
| Pendulum™3.3 EC | |
| Pendulum WDG | |

This product plus dicamba tank mixtures may not be applied by air in California.

Brush Control Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any listed rate of this product may be used in a tank mix.

For control of herbaceous weeds, use the lower tank mixture rate. For control of dense stands or tough-to-control woody brush and trees, use the higher rate.

NOTE: For side trimming treatments, use this product alone or in tank mixture with Garlon 4.

| PRODUCT | BROADCAST RATE |
|----------------------|-------------------------------|
| Arsenal 2WSL | 6 to 32 fluid ounces per acre |
| Escort | 1 to 2 ounces per acre |
| Garlon 3A*, Garlon 4 | 1 to 4 quarts per acre |

| PRODUCT | SPRAY-TO-WET RATES |
|--------------|-------------------------|
| Arsenal 2WSL | 0.06 to 0.12% by volume |
| Escort | 1 to 2 ounces per acre |

PRODUCT LOW VOLUME DIRECTED SPRAY RATES

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | |
|--------------|------------------------|
| Arsenal 2WSL | 0.1 to 0.5% by volume |
| Escort | 1 to 2 ounces per acre |

*Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

Chemical Mowing – Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 3.2 ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 3.2 ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical Mowing – Annuals

For growth suppression of some annual grasses, including annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 2.2 to 2.6 ounces of this product in 10 to 40 gallons of spray solution per acre. Applications must be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

Bromus species and Medusahead in Pastures and Rangelands.

Bromus species. This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicas*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 4.3 to 8.5 ounces of this product per acre on a broadcast basis.

For best results, treatment must coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications must be made to the same site each year until used banks are depleted and the desirable perennial grasses can become reestablished on the site.

Medusahead. To treat medusahead, apply 8.5 ounces of this product per acre as soon as plants are actively growing, and prior to the 4-leaf stage. Applications may be made in the fall or spring.

Applications to brome and medusahead may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre. When applied as directed in this label, there are no grazing restrictions.

Dormant Turfgrass

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

110/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

This product may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 4.3 to 35 ounces of this product per acre. Apply the listed rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 8.5 ounces per acre may result in injury or delayed greenup in highly maintained areas, such as golf courses and lawns. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 8.5 ounces of this product per acre in highly maintained turfgrass areas. DO NOT apply tank mixtures of this product plus Oust in highly maintained turfgrass areas. For further uses, refer to the "ROADSIDES" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

Turfgrass Renovation, Seed, or Sod Production

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing must be delayed for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

8.4 Habitat Management

Habitat Restoration and Management

This product may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species,

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

Wildlife Food Plots

This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

8.5 Injection and Frill (Woody Bush and Trees)

This product may be used to control woody brush and trees by injection or frill applications. Apply this product using suitable equipment that must penetrate into the living tissue. Inject 0.03 ounces (0.7 grams) of this product per each 2 to 3 inches of trunk diameter at breast height (DBH). This is best achieved using 0.07 fluid ounces (2 mL) of a solution prepared by dissolving 2.2 to 3.7 pounds of this product per one gallon of water either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of this product. For best results, application must be made during periods of active growth and after full leaf expansion.

8.6 Ornamentals, Plant Nurseries, and Christmas Trees

Post-directed, Trim-and-edge

This product may be used as a post-directed spray around established woody ornamental species including arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew. This product may also be used to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a nursery setting.

Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. **THIS PRODUCT IS NOT FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES.** Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

Site Preparation

This product may be used prior to planting any ornamental, nursery or Christmas tree species.

Wipers

This product may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established eucalyptus or poplar trees. See the "SELECTIVE EQUIPMENT" section of this label for further information about the proper use of wiper applicators.

Greenhouse/Shadehouse

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

This product may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

8.7 Parks, Recreational, Residential Areas

This product may be used in parks, recreational areas. It may be applied with any application equipment described in this label. This product may be used to trim-and-edge around trees, fences, and paths, around buildings, sidewalks, and other objects in these areas. This product may be used for spot treatment of unwanted vegetation. This product may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plants. This product may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the "NON-CROP AREAS AND INDUSTRIAL SITES" section apply to park and recreational areas.

8.8 Railroads

All of the instructions in the "NON-CROP AREAS AND INDUSTRIAL SITES" section apply to railroads.

Bare Ground, Ballast and Shoulders, Crossings, Spot Treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of this product may be used, as weeds emerge, to maintain bare ground. This product may be used to control tall-growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used. This product may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments:

- | | |
|-----------|------------|
| Arsenal | Krovarl DF |
| Clarity | Oust |
| Diuron | Sahara |
| Escort | Spike™ |
| Garlon 3A | Telar |
| Garlon 4 | Vanquish |
| Hyvar™X | 2,4-D |

Brush Control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4.3 to 11.6 pounds of this product per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.6 to 1 percent solution of this product when using low volume directed sprays for spot treatment. This product may be mixed with the following products for enhanced control of woody brush and trees:

- | | |
|-----------|----------|
| Arsenal | Garlon 4 |
| Escort | Tordon™K |
| Garlon 3A | |

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Additional instructions are located in the "NON-CROP AREAS AND INDUSTRIAL SITES" section under "Brush Control Tank Mixtures".

Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 8.5 to 26 ounces of this product in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

- | | |
|------------------|-----------------|
| Bahiagrass | Johnsongrass |
| Bluestem, silver | Trumpet creeper |
| Fescue, tall | Vaseygrass |

This product may be tank-mixed with Oust. If tank-mixed, use no more than 8.5 to 26 ounces of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

- | | |
|------------------|-----------------|
| Bahiagrass | Fescue, tall |
| Blackberry | Johnsongrass |
| Bluestem, silver | Poorjoe |
| Broomsedge | Raspberry |
| Dallisgrass | Trumpet creeper |
| Dewberry | Vaseygrass |
| Dock, curly | Vervain, blue |
| Dogfennel | |

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not needed, since severe injury may occur.

8.9 Roadsides

All of the instructions in the "NON-CROP AREAS AND INDUSTRIAL SITES" section apply to roadsides.

Shoulder Treatments

This product may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

Guardrails and Other Obstacles to Mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

Spot Treatment

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

114/127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

Tank Mixtures

This product may be tank-mixed with the following products for shoulder, guardrail, spot and bare ground treatments:

| | |
|-----------------|----------------|
| Clarity | Princep DF |
| Diuron | Princep Liquid |
| Endurance | Ronstar 50WP |
| Escort | Sahara |
| Krovar I DF | Simazine |
| Oust | Surflan |
| Outrider | Telar |
| Pendulum 3.3 EC | Vanquish |
| Pendulum WDG | 2,4-D |

See the "NON-CROP AREAS AND INDUSTRIAL SITES" section of this label for instructions for tank mixing.

Release of Bermudagrass or Bahiagrass

Dormant Applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup. This product may also be tank-mixed with Oust for residual control. Tank mixtures of this product with Oust may delay greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 4.3 to 35 ounces of this product per acre alone or in a tank mixture with ¼ to 1 ounce per acre of Oust. Apply the listed rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in green up and minimize injury, add no more than 1 ounce of Oust per acre on Bermudagrass and no more than ½ ounce of Oust per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

Actively Growing Bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 8.5 to 26 ounces of this product in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

| | |
|------------------|-----------------|
| Bahiagrass | Johnsongrass |
| Bluestem, silver | Trumpet creeper |

Deleted: 1

Deleted: March 19,

115/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Fescue, tall

Vaseygrass

This product may be tank-mixed with Oust. If tank-mixed, use no more than 8.5 to 17 ounces of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

- | | |
|------------------|----------------|
| Bahiagrass | Fescue, tall |
| Bluestem, silver | Johnsongrass |
| Broomsedge | Poorjoe |
| Dallisgrass | Trumpetcreeper |
| Dock, curly | Vaseygrass |
| Dogfennel | Vervain, blue |

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not needed. Since severe injury may occur.

Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 3.2 ounces of this product in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full green up or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 2.1 ounces of this product per acre, followed by an application of 1 to 2.1 ounces per acre about 45 days later. Make no more than 2 applications per year.

A tank mixture of this product plus Oust may be used. Apply 3.2 ounces of this product plus 1/4 ounce of Oust per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

8.10 Utility Sites

In utilities, this product is for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, including substations, roadsides, railroads or rights-of-way that run in conjunction with utilities.

This product is also for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all product used. Use according to the most restrictive precautionary statements for each product in the mixture. Any listed rate of this product may be used in a tank mix.

For control of herbaceous weeds, use the lower tank mixture rate. For Control of dense stands or tough-to-control woody brush and trees, use the higher rate.

Deleted: 1
Deleted: March 19,

116/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

NOTE: For side trimming treatments, use this product alone or in tank mixture with Garlon 4.

PRODUCT

| PRODUCT | BROADCAST RATE | USE SITES |
|---------------------|-------------------------------|-----------------------------|
| Arsenal 2WSL | 6 to 32 fluid ounces per acre | Utility Sites |
| Escort | 1 to 2 ounces per acre | Utility Sites |
| Garlon 3A* Garlon 4 | 1 to 4 quarts per acre | Utility Sites/Side Trimming |
| Oust | 1 to 4 ounces per acre | Utility Sites |

PRODUCT SPRAY-TO-WET RATES USE SITES

| PRODUCT | SPRAY-TO-WET RATES | USE SITES |
|--------------|--------------------------------|---------------|
| Arsenal 2WSL | 0.06 to 0.12 percent by volume | Utility Sites |
| Escort | 1 to 2 ounces per acre | Utility Sites |

PRODUCT LOW VOLUME DIRECTED SPRAY RATES USE SITES

| PRODUCT | LOW VOLUME DIRECTED SPRAY RATES | USE SITES |
|--------------|---------------------------------|---------------|
| Arsenal 2WSL | 0.1 to 0.5 percent by volume | Utility Sites |
| Escort | 1 to 2 ounces per acre | Utility Sites |

**Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

Bare Ground and Trim-and-edge

This product may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plants. This product may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects. Repeated applications of this product may be used, as weeds emerge, to maintain bare ground.

This product may be tank mixed with the following products. Refer to these products' labels for approved non-crop sites and application rates.

| | |
|-----------------|-----------------|
| Arsenal | Plateau™ |
| Banvel | Princep™DF |
| Barricade™ 65WG | Princep™ Liquid |
| Diuron | Ronstar™50WP |
| Endurance™ | Sahara™ |
| Escort | Simazine |
| Garlon 3A | Surflan™ |

9.0 WEEDS CONTROLLED

Always use the higher rate of this product per acre within the listed range when weed growth is heavy or dense or weeds are growing in an undisturbed (noncultivated) area. Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Deleted: 1
Deleted: March 19,

117/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

Refer to the following label sections for application rates for the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, this product may be used at 5.4 to 11.6 pounds per acre for enhanced results.

9.1 Annual Weeds

Use 17 ounces per acre if weeds are less than 6 inches in height or runner length and 26 to 70 ounces per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions.

For spray-to-wet applications, apply a 0.3 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead emergence in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 0.6 to 1 percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall.

WEED SPECIES


- Annoda, spurred
- Barley&
- Barnyardgrass*
- Bittercress*
- Black nightshade*
- Bluegrass, annual*
- Bluegrass, bulbous*
- Bassia, fivehook
- Brome, downy*
- Brome, Japanese*
- Browntop panicum*
- Buttercup*
- Carolina foxtail*
- Carolina geranium
- Castor bean
- Cheatgrass*
- Cheeseweed (Malva parviflora)
- Chervil*
- Chickweed*
- Cocklebur*
- Copperleaf, hophornbeam
- Corn*
- Corn speedwell*
- Crabgrass*
- Dwarf dandelion*
- Eastern mannagrass*
- Eclipta*
- Fall panicum*
- Falsedandelion*
- Falseflax, smallseed*
- Fiddleneck
- Field pennycress*
- Filaree

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- Fleabane, annual*
- Fleabane, hairy (*Conyza bonariensis*)*
- Fleabane, rough*
- Florida pusley
- Foxtail*
- Goatgrass, jointed*
- Goosegrass
- Grain sorghum (milo)*
- Grousel, common*
- Hemp sesbania
- Henbit
- Horseweed/Marestail (*Conyza Canadensis*)
- Itchgrass*
- Johnsongrass, seedling
- Junglerice
- Knotweed
- Kochia
- Lamb's-quarters*
- Little barley*
- London rocket*
- Mayweed
- Medusahead*
- Morningglory (*Ipomoea* spp)
- Mustard, blue*
- Mustard, tansy*
- Mustard, tumble*
- Mustard, wild*
- Oats
- Pigweed*
- Plains/Tickseed coreopsis*
- Prickly lettuce*
- Puncturevine
- Purslane, common
- Ragweed, common*
- Ragweed, giant
- Red rice
- Russian thistle
- Rye*&
- Ryegrass*
- Sandbur, field*
- Shattercane*
- Shepherd's-purse*
- Sicklepod
- Signalgrass, broadleaf*
- Smartweed, ladysthumb*
- Smartweed, Pennsylvania*
- Sowthistle, annual
- Spanishneedles
- Speedwell, purslane*
- Sprangletop*
- Spurge, annual
- Spurge, prostrate*
- Spurge, spotted*
- Spurry, umbrella*



Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

- Starthistle, yellow
- Stinkgrass*
- Sunflower*
- Teaweed/Prickly sida
- Texas panicum*
- Velvetleaf
- Virginia copperleaf
- Virginia pepperweed*
- Wheat*
- Wild oats*
- Witchgrass*
- Woolly cupgrass*
- Yellow rocket

*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 8.7 ounces of this product per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

9.2 Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (boot stage in grasses and bud information in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the listed range.

Ensure thorough coverage when sing spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low volume directed spot treatments, apply a 2.8 to 5.5 percent solution of this product. Allow 7 or more days after application before tillage.

| WEED SPECIES | RATE (LB/A) | % SOLUTION BY WEIGHT |
|---|-------------|----------------------|
| Alfalfa* | 1.1-2.2 | 1 |
| Alligatorweed* | 4.4 | 0.7 |
| Anise (fennel) | 2.2-4.4 | 0.6-1 |
| Bahiagrass | 3.2-5.5 | 1 |
| Beachgrass, European (Ammophila arenaria) | - | 2.6 |
| Bentgrass* | 1.6 | 1 |
| Bermudagrass | 5.5 | 1 |
| Bermudagrass, water (knotgrass) | 1.6 | 1 |
| Bindweed, field | 0.6-5.5 | 1 |
| Bluegrass, Kentucky | 2.2 | 1 |
| Bluewed, Texas | 4.4-5.5 | 1 |
| Brackenfern | 3.2-4.4 | 0.6-0.7 |
| Bromegrass, smooth | 2.2 | 1 |
| Bursage, woolly-leaf | - | 1 |
| Canarygrass, reed | 2.2-3.4 | 1 |
| Cattail | 3.2-5.5 | 1 |
| Clover, red, white | 3.2-5.5 | 1 |
| Cogongrass | 3.2-5.5 | 1 |
| Dallisgrass | 3.2-5.5 | 1 |

Deleted: 1

Deleted: March 19,

120/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | |
|--------------------------|---------|---------|
| Dandelion | 3.2-5.5 | 1 |
| Dock, curly | 3.2-5.5 | 1 |
| Dogbane, hemp | 4.4 | 1 |
| Fescue (except tall) | 3.2-5.5 | 1 |
| Fescue tall | 1.1-3.2 | 1 |
| German ivy | 2.2-4.4 | 0.6 - 1 |
| Guineagrass | 3.2 | 0.6 |
| Horsenettle | 3.2-5.5 | 1 |
| Horseradish | 4.4 | 1 |
| Iceplant | 2.2 | 0.7-1 |
| Jerusalem artichoke | 3.2-5.5 | 1 |
| Johnsongrass | 0.6-3.2 | 0.6 |
| Kikuyugrass | 2.2-3.2 | 1 |
| Knapweed | 4.4 | 1 |
| Lantana | - | 0.6-0.7 |
| Lespedeza | 3.2-5.5 | 1 |
| Milkweed, common | 3.2 | 1 |
| Muhly, wirestem | 2.2 | 1 |
| Mullein, common | 3.2-5.5 | 1 |
| Napiergrass | 3.2-5.5 | 1 |
| Nightshade, silverleaf | 2.2 | 1 |
| Nutsedge, purple, yellow | 3.2 | 0.6 - 1 |
| Orchardgrass | 2.2 | 1 |
| Pampasgrass | 3.2-5.5 | 0.7-1 |
| Paragrass | 3.2-5.5 | 1 |
| Pepperweed, perennial | 4.4 | 1 |
| Phragmites* | 3.2-5.5 | 0.6-1 |
| Poison hemlock | 2.2-4.4 | 0.6-1 |
| Quackgrass | 1.2-3.2 | 1 |
| Redvine* | 2.2 | 1 |
| Reed, giant | 4.4-5.5 | 1 |
| Ryegrass, perennial | 1.2-3.2 | 0.6 |
| Smartweed, swamp | 3.2-5.5 | 1 |
| Spurge, leafy* | - | 1 |
| Sweet potato, wild* | - | 1 |
| Thistle, artichoke | 2.2-3.2 | 0.6-1 |
| Thistle, Canada | 2.2-3.2 | 1 |
| Timothy | 2.2-3.2 | 1 |
| Torpedograss* | 4-5.5 | 1 |
| Trumpetcreeper* | 2.2-3.2 | 1 |
| Vaseygrass | 3.2-5.5 | 1 |
| Velvetgrass | 3.2-5.5 | 1 |
| Wheatgrass, western | 2.2-3.2 | 1 |

*Partial control

9.3 Woody Brush and Trees

Apply this product after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

Deleted: 1
Deleted: March 19.

121/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low volume directed-spray spot treatments, apply a 2.6 to 5.1 percent solution of this product.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

| WEED SPECIES | BROADCAST RATE (LB/A) | HAND-HELD SPRAY-TO-WET % SOLUTION W/W |
|---------------------------|-----------------------|---------------------------------------|
| Alder | 3.2-4.4 | 0.6-1 |
| Ash* | 2.2-5.5 | 0.6-1 |
| Aspen, quaking | 2.2-3.2 | 0.6-1 |
| Bearclover (Bearmat)* | 2.2-5.5 | 0.6-1 |
| Beech* | 2.2-5.5 | 0.6-1 |
| Birch | 2.2 | 0.6 |
| Blackberry | 3.2-4.4 | 0.6-1 |
| Blackgum | 2.2-5.5 | 0.6-1 |
| Bracken | 2.2-5.5 | 0.6-1 |
| Broom; French, Scotch | 2.2-5.5 | 0.7-1 |
| Buckwheat, California* | 2.2-4.4 | 0.6-1 |
| Cascara* | 2.2-5.5 | 0.6-1 |
| Castor bean | 4.4 | 1 |
| Catsclaw* | - | 0.6-0.7 |
| Ceanothus* | 2.2-5.5 | 0.6-1 |
| Chamise* | 2.2-5.5 | 0.6 |
| Cherry; bitter, black pin | 2.2-3.2 | 0.6-0.7 |
| Cottonwood, eastern | 4.4 | 1 |
| Coyote brush | 3.2-4.4 | 0.7-1 |
| Cypress, swamp, bald | 4.4 | 1 |
| Deerweed | 2.2- | 0.6 |
| Dogwood* | 2.2-5.5 | 0.6-1 |
| Elderberry | 2.2 | 0.6 |
| Elm* | 2.2-5.5 | 0.6-1 |
| Eucalyptus | - | 1 |
| Gallberry | 4.4 | 1 |
| Hackberry, western | 4.4 | 1 |
| Gorse* | 2.2-5.5 | 0.6-1 |
| Hasardia* | 2.2-4.4 | 0.6-1 |
| Hawthorn | 2.2-3.2 | 0.6-1 |
| Hazel | 2.2 | 0.6 |
| Hickory* | 2.2-5.5 | 0.6-1 |
| Honeysuckle | 3.2-4.4 | 0.6-1 |
| Hornbeam, American* | 2.2-5.5 | 0.6-1 |

| |
|--------------------|
| Deleted: 1 |
| Deleted: March 19, |

122/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | | |
|---|---------|-------|
| Huckleberry | 4.4 | 1 |
| Kudzu | 4.4 | 1 |
| Locust, black* | 2.2-4.4 | 0.6-1 |
| Madrone resprouts* | - | 1 |
| Magnolia, sweetbay | 4.4 | 1 |
| Manzanita* | 2.2-5.5 | 0.6-1 |
| Maple, red | 2.2-4.4 | 0.6-1 |
| Maple, sugar | - | 0.6-1 |
| Monkey flower* | 2.2-4.4 | 0.6-1 |
| Oak; black, white* | 2.2-4.4 | 0.6-1 |
| Oak, post | 3.2-4.4 | 0.6-1 |
| Oak; northern, pin | 2.2-4.4 | 0.6-1 |
| Oak, Scrub* | 2.2-4.4 | 0.6-1 |
| Oak; southern red | 2.2-3.2 | 0.6-1 |
| Orange, Osage | 4 | 1 |
| Peppertree, Brazilian (Florida holly)* | 2.2-5.5 | 0.6-1 |
| Persimmon* | 2.2-5.5 | 0.6-1 |
| Pine | 2.2-5.5 | 0.6-1 |
| Poison Ivy | 4.4-5.5 | 1 |
| Poison oak | 4.4-5.5 | 1 |
| Poplar, yellow* | 2.2-5.5 | 0.6-1 |
| Redbud, eastern | 2.2-5.5 | 0.6-1 |
| Redcedar, eastern | 4.4 | 1 |
| Rose, multiflora | 2.2 | 0.6 |
| Russian olive* | 2.2-5.5 | 0.6-1 |
| Sage, black | 2.2-4.4 | 0.6 |
| Sage, white* | 2.2-5.4 | 0.6-1 |
| Sage brush, California | 2.2-4.4 | 0.6 |
| Salmonberry | 2.2 | 0.6 |
| Salt-cedar* | 2.2-5.5 | 0.6-1 |
| Sassafras* | 2.2-5.5 | 0.6-1 |
| Sourwood* | 2.2-5.5 | 0.6-1 |
| Sumac; laurel, poison, smooth, Sugarbrush, winged* | 2.2-4.4 | 0.6-1 |
| Sweetgum | 2.2-3.2 | 0.6-1 |
| Swordfern* | 2.2-5.5 | 0.6-1 |
| Tallowtree, Chinese | - | 0.6 |
| Tan oak resprouts* | - | 1 |
| Thimbleberry | 2.2 | 0.6 |
| Tobacco, tree* | 2.2-4.4 | 0.6-1 |
| Toyon* | - | 1 |
| Trumpet creeper | 2.2-3.2 | 0.6-1 |
| Vine maple* | 2.2-5.5 | 0.6-1 |
| Virginia creeper | 2.2-5.5 | 0.6-1 |
| Waxmyrtle, southern* | 2.2-5.5 | 0.6-1 |
| Willow | 3.2 | 0.6 |
| Yerbasenta, California* | - | 1 |

*Partial control

10.0 LIMIT OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, MEY Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set

Deleted: 1

Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESSED WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of the Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, applications to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

To the extent consistent with applicable law upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.

Manage, Monsanto and Vine symbol are trademarks of Monsanto Technology LLC.
Escort, Hyvar, Karmex, Krovar, Oust, and Telar are trademarks of E.I. duPont de Nemours & Co. Inc.
Garlon, Spike, Surflan and Tordon are trademarks of Dow AgroSciences
Barricade, Endurance, Princep and Vanquish are trademarks of Sygenta Group
Ronstar is a trademark of Aventis Group
Arsenal, Pendulum, Plateau, and Sahara are trademarks of BASF Corporation

This product is protected by U.S. Patent Nos. 6,184,12; 6,245,713; and 6,093,681. Other patents pending. No license is granted under any non-U.S. patent(s).

[[Revision] [Version Date] [MM/YYYY]]

Deleted: 1
Deleted: March 19,

124/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

(Note to Reviewer: The following information will be affixed to the unit package/container.)

GlySupreme 75-WDG

[Water Soluble Dry Herbicide] [Water Dispersible Granule [Herbicide]]

(Alternate Brand Names: GlySupreme 75-Water Dispersible Granule, GlySupreme 75-WDG Herbicide, GlySupreme 75-WDG Glyphosate Herbicide)

This product is a post-emergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush, and trees. It is formulated as a water soluble granule. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions. Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees because severe injury or destruction may result.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethy) glycine, in the form of its ammonium salt..... 75.7%

Other Ingredients..... 24.3%

Total..... 100.0%

*Equivalent to 68.7% of glyphosate acid

1-pound-of-[this product]-[GlySupreme 75-WDG] contains 0.687 pounds of glyphosate acid.

EPA Reg. No.: 80967-13

EPA Est. No.: [73432-CHN-001]

Net Contents:

Manufactured [for][by]:

MEY Corporation

121 S. Estes Drive, Suite 101

Chapel Hill, NC 27514 USA

[Lot/Batch code/number]

(Note to reviewer: Lot or Batch number may appear on label or printed directly on packaging.)

KEEP OUT OF REACH OF CHILDREN CAUTION

See attached booklet for [additional precautionary statements] [first aid] [complete directions for use] [and warranty]

FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE (800) 613-0427

IN CASE OF AN EMERGENCY INVOLVING THIS HERBICIDE PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL (800) 262-8200

FIRST AID

| | |
|----------------------|--|
| If swallowed: | Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not |
|----------------------|--|

Deleted: 1
Deleted: March 19,

125/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

| | |
|---|--|
| | give anything by mouth to an unconscious person. |
| If inhaled: | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. |
| If in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice |
| If on skin or clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-262-8200. | |

Read the entire label before using this product. Use only according to label instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Caution. Harmful if swallowed. Harmful if inhaled. Avoid breathing spray mist or vapor..

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Personal Protective Equipment (PPE): Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

(Note to Reviewer: For products in containers >50 lbs): Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System

Deleted: 1
Deleted: March 19,

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

(NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product must be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. Keep container closed when not in use to prevent spills and contamination.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal Facility.

Container Handling: Non-refillable container. Do not reuse or refill this container.

(Note to reviewer: For non-refillable plastic containers (≤50 lbs) small enough to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable plastic containers (>50 lbs) too large to shake):

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable paper and plastic bags, large fiber sacks):

Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Note to reviewer: For non-refillable fiber drums with liners or bag-in-box packaging):

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke. If [drum] [box] carton is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Deleted: 1

Deleted: March 19,

127/
127

(Note to reviewer: [bracketed text] is optional or alternate wording; (parenthetical text) is informational.)

WARRANTY AND LIABILITY

To the extent consistent with applicable law, MEY Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESSED WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

[[Revision] [Version Date] [MM/YYYY]]

Deleted: 1
Deleted: March 19,