

81198-1

8/8/2004

118



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
81198-1

Date of Issuance:
AUG 8 2004

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
FreeGrow Headstart SFM
(sulfometuron methyl)
75 Herbicide

Name and Address of Registrant (include ZIP Code):

Woodland Specialists, Inc
Hanger 209 Craig Field
Selma, AL 36701

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 81198-1."
 - b. On page 2 under Conifer and Conifer Release, West, revise "Vegetation Management" to read "Woodlands Specialists".
 - c. On page 3, in the Non-Agricultural Use Requirements Box, add the statement "Keep unprotected persons out of treated areas until sprays have dried."
 - d. On page 3, under Application information delete the phrase "other similar areas including" from the phrase "other similar areas including governmental and private lands." On page 4, delete the phrases "other noncrop areas" and "other industrial sites" and replace them a listing of specific sites. These types of phrases are no longer permitted to appear on Agency approved labels.

Signature of Approving Official:

Date:

8/6/04

3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release of this product for shipment constitutes acceptance of these conditions.

A stamped copy of labeling is enclosed for your records.

ACCEPTED
with COMMENTS
In EPA Letter Dated: FreeGrow
AUG - 6 2004

3/8

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

Headstart Herbicide

SFM 75 (SULFOMETURON METHYL)

SPECIMEN LABEL

ACTIVE INGREDIENT:

SULFOMETURON METHYL

{METHYL 2-[[[4,6-DIMETHYL-2-PYRIMIDINYL]AMINO]

-CARBONYL]AMINO]SULFONYL]BENZOATE}..... 75%

INERT INGREDIENTS:..... 25%

TOTAL:..... 100%

EPA REG. NO.

EPA EST. NO. 81198-AL-001

KEEP OUT OF REACH OF CHILDREN CAUTION

First Aid

If in Eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Causes moderate eye injury (irritation). Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

GENERAL INFORMATION

SFM 75 is a dispersible granule that is mixed in water and applied as a spray. SFM 75 controls many annual and perennial grasses and broadleaf weeds in forestry and noncrop sites.

SFM 75 may be used for general weed control on industrial noncrop sites and for selective weed control in certain types of unimproved turf grasses on industrial sites. It can also be used for selective weed control in forest site preparation and in the release of several types of pines and certain hardwoods.

SFM 75 may be applied by air (helicopter only) to control various weeds and grasses on railroad rights-of-way in the states of Colorado (except Saguache, Rio Grande, Alamosa, Costilla and Conejos counties), Kansas, Nebraska, New Mexico, North Dakota, South Dakota and Wyoming.

SFM 75 controls weeds by both preemergence and postemergence activity. Preemergence treatments control or suppress weeds through root uptake while postemergence control works through root and foliar uptake. The best results are obtained when the application is made before or during the early stages of weed growth before weeds develop an established root system. Moisture is required to move SFM 75 into the root zone of weeds for preemergence control. When rainfall is low, SFM 75 may not provide satisfactory control.

SFM 75 is noncorrosive, nonflammable, nonvolatile, and does not freeze.

For best postemergence results, apply SFM 75 to young, actively growing weeds. The use rate depends upon the weed species, weed size at application, and soil texture. The degree and duration of control may depend on the following:

- Weed spectrum and infestation intensity
- Weed size at application
- Environmental conditions at and following treatment.
- Soil pH, soil moisture, and soil organic matter

Use a high rate on established plants and on fine-textured soils and a lower rate on smaller weeds and coarse-textured soils.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

SFM 75 is absorbed by both the roots and foliage of plants, rapidly inhibiting the growth of susceptible weeds. Two to 3 weeks after application to weeds, leaf growth slows, and the growing points turn reddish-purple. Within 4 to 6 weeks of application, leaf veins and leaves become discolored, and the growing points subsequently die.

Warm, moist conditions following application accelerate the herbicidal activity of SFM 75; cold, dry conditions delay the herbicidal activity. In addition, weeds hardened-off by drought stress are less susceptible to SFM 75.

Rainfall is needed to move SFM 75 into the soil for preemergence weed control, but postemergence weed control may be reduced if rainfall occurs too soon after application.

RESISTANCE

When herbicides with the same mode of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant weed biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. These resistant weed biotypes may not be adequately controlled. Cultural practices such as tillage, preventing weed escapes from going to seed, and using herbicides with different modes of action within and between crop seasons can aid in delaying the proliferation and possible dominance of herbicide resistant weed biotypes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SFM 75 should be used only in accordance with recommendations on this label.

Woodlands Specialists will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by Woodlands Specialists. User assumes all risks associated with such nonrecommended use.

Do not use on food or feed crops.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult

SFM 75

Specimen Label

4/8

the agency in your State responsible for pesticide regulation.

AGRICULTURAL USES

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves, such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber \geq 14 mils.
- Shoes plus socks

FORESTRY

Application Information

SFM 75 is recommended to control many broadleaf weeds and grasses in forestry sites. Apply by ground equipment or by air (helicopter only).

Application Timing

Apply SFM 75 before herbaceous weeds emerge or shortly thereafter. Apply only during seasons when rainfall is sufficient to activate the herbicide in the soil.

Weeds Controlled

SFM 75 effectively controls the following weeds when applied at the use rates indicated for the respective crop species:

Chickweed	Nutsedge (yellow)
Crabgrass	Panicums (broadleaf, fall, narrow)
Dogfennel	Pokeweed
Fescue	Ragweed
Fireweed (willowweed)	Shepherd's purse
Goldenrod	White snakeroot
Horseweed	Yellow sweetclover
Kentucky bluegrass	

See also weeds controlled under Application Information-Noncrop (Industrial) Sites

Application Rates

Apply SFM 75 at the rates indicated by region. Use a low rate on coarse-textured soils (i.e. loamy sands, sandy loams) and a higher rate on fine-textured soils (i.e. sandy clay loams and silty clay loams).

CONIFERS

Conifer Site Preparation

-Application Before Transplanting

Make all applications before transplanting to control herbaceous weeds.

Southeast- Apply 2 to 8 oz per acre for loblolly, longleaf, slash, and Virginia pine. Transplant longleaf pine at least 60 days after treatment.

Northeast and Lake States- Apply 2 to 4 oz per acre for black spruce. Transplant at least 13 months after treatment.

Apply 2-1/2 to 4 oz SFM 75 plus Accord¹ (as registered) for larch and tamarack. Transplant the following spring or summer but not less than 8 months after

treatment.

West- Apply 2 to 4 oz per acre for coastal redwood, Douglas fir, grand fir, lodgepole pine, ponderosa pine, western larch, western white pine, and white fir. For ponderosa pine in California and other arid areas, apply in the fall and transplant the following spring. Other conifer species may be planted; however, Vegetation Management has not tested the response of unlisted conifer species, and, therefore, cannot assume responsibility for any injury that may occur to species not listed above.

Conifer Release

-Application After Transplanting

Apply SFM 75 after transplanting to control herbaceous weeds.

Southeast- Apply 2 to 8 oz per acre for loblolly, longleaf, slash or Virginia pine. Apply 1 to 1-1/2 oz per acre for eastern white pine.

Tank Mix Combinations (Southeast only)- To control a broader spectrum of weeds in stands of loblolly, longleaf, or slash pine, apply 2 to 4 oz of SFM 75 plus 2 to 3 pt of DuPont Velpar L² Herbicide or 2/3 to 1 lb of DuPont Velpar DF² Herbicide. Tank mix may injure or kill trees when applied during high humidity and temperature.

To enhance control of Bermudagrass and Johnsongrass in stands of loblolly pine, apply 2 oz of SFM 75 plus 4 to 6 fl oz of Arsenal³ Applicators Concentrate. For the best results make the application during late winter through spring when weeds first emerge. Arsenal may temporarily inhibit pine growth if it is applied when pine is actively growing.

For control of many annual weeds particularly on cropland conversion areas, apply 2 to 4 oz of SFM 75 plus 4 to 8 pt of Aatrex⁴ 4L per acre. Use the higher rates on medium to fine texture soils where organic matter exceeds 2%. Use only on tree species specifically listed on both the SFM 75 and Aatrex⁴ 4L labels.

Northeast and Lake States- Apply 2 to 8 oz per acre for jack or Virginia pine. Apply 1 to 1-1/2 oz per acre for eastern white pine. Apply 1 1/2 to 3 oz per acre for white spruce. Make applications when trees are dormant. Applications at budbreak and later stages of active growth may severely injure or kill trees.

West- Apply 2 to 4 oz per acre for coastal redwood, Douglas fir, grand fir, lodgepole pine, western larch, and western white pine. Application may be made for the release of other conifer species present on the site; however, Vegetation Management has not tested the response of unlisted conifer species, and, therefore, cannot assume responsibility for any injury that may occur to conifers not listed above. Applications made after dormancy break in the spring and before the final resting bud has hardened in the fall may severely injure or kill trees.

HARDWOODS

Hardwood Site Preparation

-Application Before Transplanting

Apply 3 to 5 oz on sites where northern red oak, white oak, chestnut oak, American sycamore, ash (white or green), red maple, sweetgum, or yellow poplar are to be planted. Make all applications before transplanting.

Hardwood Release

-Application After Transplanting

Apply 1 to 4 oz per acre in stands of American sycamore, ash (white or green), bald cypress, oaks (such as chestnut, northern red, southern red, overcup, pin, swamp chestnut, cherrybark, water, white, pin, etc.), red maple, sweetgum, or yellow poplar.

SFM 75 should be applied before the hardwood tree seedlings or transplants break dormancy (bud swell stage). Applications made over the top after the trees have broken dormancy may injure or kill the trees.

IMPORTANT PRECAUTIONS-FORESTRY ONLY

-Applications of SFM 75 made to trees, conifers, or hardwoods that are suffering from loss of vigor caused by insects, diseases, drought, winter damage, animal damage, excessive soil moisture, planting shock, or other stresses, may injure or kill the trees.

-Applications of SFM 75 made for release (trees present) should only be made after adequate rainfall has closed the planting slit and settled the soil around the roots following transplanting.

SFM 75

Specimen Label

-Do not apply SFM 75 to conifers or hardwoods grown for Christmas trees or ornamentals.

-If a surfactant is used with SFM 75, allowing the spray to contact tree foliage may injure or kill trees. The user assumes all responsibility for tree injury if a surfactant is used with SFM 75 treatments applied after planting.

-SFM 75 application may result in damage and mortality to other species of trees when they are present on sites with those listed in the preceding recommendations for forestry uses.

-Use on hardwood trees growing in soils having a pH of 7 or greater may injure or kill the trees.

-Careful consideration must be given by an experienced and knowledgeable forester to match the requirements of the hardwood tree species to the conditions of the site. Treatment of species mismatched to the site may injure or kill the trees.

-SFM 75 is not recommended for use on poorly drained or marshy sites, but it may be used where plantings are on raised beds.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Selective non-crop industrial weed control and weed control in turf (industrial, unimproved only) are not within the scope of the Worker Protection Standard.

NONCROP (INDUSTRIAL) SITES

Application Information

SFM 75 is recommended for use for general weed control on noncrop, industrial sites such as airports, military installations, fence rows, roadsides and associated rights-of-way, lumberyards, petroleum tank farms, pipeline and utility rights-of-way, pumping installations, railroads, storage areas, plant sites, and other similar areas including governmental and private lands. Apply by ground equipment only unless directed otherwise by this label or supplemental labeling.

Combination with other herbicides broadens the spectrum of weeds controlled. In addition, total vegetation control can be achieved with higher rates of SFM 75 plus residual-type companion herbicides. To improve the control of weeds, add surfactant at 0.25% by volume.

AERIAL APPLICATION ON RAILROAD RIGHT-OF-WAY IN THE STATES OF COLORADO, KANSAS, NEBRASKA, NEW MEXICO, NORTH DAKOTA, SOUTH DAKOTA AND WYOMING: SFM 75 may be applied by air (helicopter only) to control various weeds and grasses on railroad rights-of-way only. Helicopter applications may only be made in the states of Colorado (except Saguache, Rio Grande, Alamosa, Costil and Conejos counties), Kansas, Nebraska, New Mexico, North Dakota, South Dakota and Wyoming.

ARID AREAS (AREAS OF 20" OR LESS ANNUAL RAINFALL)

Application Timing

Apply SFM 75 as a preemergence or early postemergence spray during the rainy season when weeds are actively germinating or growing.

Weeds Controlled

SFM 75 effectively controls the following broadleaf weeds and grasses when applied at the rates shown.

Application Rates

Apply SFM 75 at the rates indicated by weed type. When applied at lower rates, SFM 75 provides short-term control of weeds listed; when applied at higher rates, weed control is extended.

Broadleaf Weeds -1 1/3 to 2 oz per acre

- | | |
|-------------------|--------------------|
| Annual sowthistle | Common yarrow |
| Black mustard | Curly dock |
| Buckhorn plantain | Prickly coontail |
| Burclover | Seaside heliotrope |
| Carolina geranium | Spreading orach |
| Chickweed | Sunflower |
| Common mallow | Western ragweed |
| Common speedwell | Whitestem filaree |

Grasses (up to 6 to 12" tall) -1 1/3 to 2 oz per acre

- | | |
|-------------------|--------------------|
| Annual bluegrass | Red brome |
| Barnyardgrass | Reed Canarygrass |
| Cheat | Ripgut brome |
| Foxtail barley | Seashore saltgrass |
| Foxtail fescue | Signalgrass |
| Italian ryegrass | Yellow foxtail |
| Jointed goatgrass | |

Grasses-2 to 3 oz per acre

Smooth brome

The weeds listed in **Areas of 20" or More Annual Rainfall** can also be controlled in Arid Areas; however, SFM 75 must be applied at 3 to 8 oz per acre to control those weeds. These higher rates also provide control of severe infestations and longer term control of weeds listed for arid areas.

AREAS OF 20" OR MORE ANNUAL RAINFALL

Application Timing

Apply SFM 75 as a preemergence or early postemergence spray during the rainy season when weeds are actively germinating or growing.

Weeds Controlled

SFM 75 effectively controls the following broadleaf weeds and grasses when applied at the rates shown.

Application Rates

Apply SFM 75 at the rates indicated by weed type. When applied at lower rates, SFM 75 provides short term control of weeds listed; when applied at higher rates, weed control is extended.

Broadleaf Weeds-3 to 5 oz per acre

- | | |
|-------------------------|---------------------|
| Bouncingbet | Pigweed |
| Burclover | Purple starthistle |
| Carolina geranium | Ragweed |
| Common chickweed | Sowthistle (annual) |
| Common dandelion | Sunflower |
| Common speedwell | Sweet clover |
| Common yarrow | Tansymustard |
| Crimson clover | Tansy ragwort |
| Dogfennel | Tumble mustard |
| Hoary cress (whitewort) | Vetch |
| Little mallow | Wild carrot |
| Mustard | Wild oats |
| Ox-eye daisy | Yellow rocket |
| Pepperweed | |

Broadleaf Weeds-6 to 8 oz per acre

- | | |
|-----------------|-----------------------|
| Bedstraw | Horsetail (Equisetum) |
| Canada thistle | Kudzu |
| Curly dock | Musk thistle |
| Redstem filaree | Turkey mullein |
| Goldernrod | Wild blackberry |

SFM 75

Specimen Label ^{6/8}

Grasses—3 to 5 oz per acre

Alta fescue	Kentucky bluegrass
Annual bluegrass	Little barley
Annual ryegrass	Red brome
Bahiagrass	Red fescue
Barnyardgrass	Reed canarygrass
Downy brome	Ripgut brome
Fescue	Ryegrass
Foxtails (except green)	Smooth brome
Foxtail barley	Sprangletop (annual)
Indiangrass	Wheat (volunteer)
Italian ryegrass	

Grasses—6 to 8 oz per acre

Johnsongrass

For short-term (up to 3 months) control of johnsongrass, apply early postemergence. Repeat treatment if additional control is desired or if regrowth occurs.

Note: Use the higher level of recommended dosage ranges under the following conditions:

- Heavy weed growth
- Soil containing more than 2-1/2% organic matter
- High soil moisture areas, such as along road edges or railroad shoulders

Specific Weed Problems

—Noncrop (Industrial) Sites

Kochia, Russian Thistle, and Prickly Lettuce

Since biotypes of kochia, Russian thistle, and prickly lettuce are known to be resistant to SFM 75, tank mixture combinations with herbicides having different modes of action, such as KARMEX DF, HYVAR X, or KROVAR I DF, must be used. In areas where resistance is known to exist, these weeds should be treated postemergence with other herbicides registered for their control, such as 2,4-D or dicamba. Do not allow kochia, Russian thistle, or prickly lettuce to form mature seed.

TANK MIX COMBINATIONS

To improve preemergence to early postemergence control of weeds and grasses, add 2 to 8 oz of SFM 75 per acre to the recommended rates of the following herbicides: DuPont HYVAR X herbicide; DuPont KARMEX DF Herbicide; DuPont KROVAR I DF herbicide; DuPont VELPAR L² Herbicide; DuPont VELPAR Herbicide; DuPont ESCORT² Herbicide (do not use in California); DuPont TELAR² Herbicide; glyphosate; dicamba; or 2,4-D; or registered generic equivalents of the trade names listed.

Apply SFM 75 plus a companion herbicide at the rates and timing as shown on package labels for target weeds. For application method and other use specifications, use the most restrictive directions for the intended combination.

Do not tank mix SFM 75 with DuPont HYVAR XL Herbicide.

UNDER ASPHALT AND CONCRETE PAVEMENT

Application Information

SFM 75 can be used to control weeds under asphalt and concrete pavement, such as that used in parking lots, highway shoulder, median strips, roadways, and other industrial sites.

SFM 75 will not control tubers, rhizomes, woody vegetation such as small trees, brush or woody vines.

SFM 75 should only be used in an area that has been prepared according to good construction practices. Use sufficient water to ensure uniform coverage, generally 100 gal per acre. Agitate the tank continuously to keep SFM 75 in suspension.

Application Timing

SFM 75 should be applied immediately before paving to avoid lateral movement of the herbicide as a result of soil movement due to rainfall or mechanical means.

Application Rate

Apply SFM 75 at 4 to 8 oz per acre. Use a higher rate on hard-to-control weeds and for long-term control.

Tank Mix Combinations

---Under Asphalt and Concrete Pavement

For broader spectrum control or for an extended period of control under asphalt or concrete pavement, SFM 75 may be applied as a tank mix with HYVAR X at 6 to 15 lb per acre or KROVAR I DF at 7 to 15 lb per acre.

IMPORTANT PRECAUTIONS—UNDER ASPHALT ONLY

-Do not use SFM 75 under pavement in residential properties such as driveways, or in recreational areas, including jogging or bike paths, tennis courts, or golf cart paths.

-Desirable plants may be injured if their roots extend into treated areas or if planted in treated areas.

TURF, INDUSTRIAL (UNIMPROVED ONLY)

Application Information

SFM 75 is recommended to control weeds on unimproved industrial turf, on roadsides, or on other noncrop sites where the turf is well established as a ground cover. Applications may temporarily suppress grass growth and inhibit seedhead formation (chemical mowing).

Bermudagrass Release

Application Timing

Apply SFM 75 after bermudagrass has broken dormancy and is well established, usually 30 days after initial spring flush. If additional applications are necessary, apply SFM 75 again during late spring to early summer. On established weeds, apply SFM 75 1 to 2 weeks after mowing for the best results.

SFM 75 may also be applied in late fall or early winter. Use the lower rates on small seedling weeds and a higher rate on larger weeds. Also, refer to the listing of **Weeds Controlled under Noncrop (Industrial) Weed Control**.

Weeds Controlled

SFM 75 may be used to control the following weeds when applied at the use rates shown.

Late Spring to Early Summer—1 to 2 oz per acre

Carolina Geranium	Goldenrod
Fescue	Spotted Spurge
Foxtail	Wild carrot

Spring to Fall—2 to 3 oz per acre

Johnsongrass

Late Fall to early Winter—1 to 4 oz per acre

Carolina geranium	Little barley
Common chickweed	Wild blackberry
Fescue	

Tank Mix Combinations—Bermudagrass (South Only)

Apply 1 to 2 oz SFM 75 per acre as a tank mix with 3 to 4 lb active ingredient of MSMA per acre on well-established bermudagrass during the summer. Refer to the MSMA package label for a list of additional weeds that may be controlled. Two or more sequential applications of MSMA alone may be necessary to maintain weed control.

Centipedegrass Release

Application Timing

Apply 1 to 2 ounces of SFM 75 in the fall or early winter, or in the early summer

SFM 75

Specimen Label

following greenup of the centipede. Refer to the listing of **Weeds Controlled under Bermudagrass Release.**

Bahiagrass Release and Seedhead Suppression

Application Timing

Apply 1/2 to 1 oz SFM 75 per acre to turf after green-up and before seedheads emerge (boot stage). Ensure that desirable grasses are well established at application, as premature treatment may result in top kill and stand reduction of desirable turf. Make only one application per year.

Smooth Brome and Crested Wheatgrass Release and Suppression

Application Timing

Apply 1 oz SFM 75 per acre to turf after green-up and before seedheads emerge (boot stage). Ensure that desirable grasses are well established at application, as premature treatment may result in top kill and stand reduction of desirable turf. Make only one application per year.

Weeds Controlled

SFM 75 may be used to control the following weeds when applied at the use rates shown.

Late Spring to Early Summer—1 oz. per acre

Downy Brome Goldenrod
Foxtail

IMPORTANT PRECAUTIONS – INDUSTRIAL, UNIMPROVED TURF

-Excessive injury to turf may result if a surfactant is used with SFM 75 application made to actively growing turf. The user assumes all responsibility for turf injury if a surfactant is used with SFM 75 treatments applied to actively growing turf.

-SFM 75 may temporarily discolor or cause top kill of turf grasses. Applications made while turf is dormant may delay green-up in the spring.

-Annual retreatments may reduce vigor, particularly at the higher recommended rates, where bahiagrass, crested wheatgrass and smooth brome are grown.

-SFM 75 application on turf that is under stress from drought, insects, disease, cold temperatures or late spring frost, may result in injury.

-Do not apply SFM 75 to turf within 1 year of planting as stand reduction may result.

SPRAY EQUIPMENT

Following a SFM 75 application, do not use sprayer for application to agricultural or ornamental crops. The mixing and application equipment must be used for forestry and noncrop application only. This is extremely important as low rates of SFM 75 can kill or severely injure most crops.

BROADCAST APPLICATION

Ground

Use 15 to 40 gal of water per acre when applying SFM 75 as a broadcast application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern. Be sure the sprayer is calibrated before use. Avoid overlapping and shut off spray booms while starting, turning, slowing, or stopping to avoid injury to desired species.

Air (Helicopter Only)

Use 5 to 15 gal of water per acre when applying SFM 75. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern. Do not use fixed-wing aircraft. Be sure the sprayer is calibrated. Avoid overlapping and shut off spray booms while starting, turning or slowing to avoid injury to desired species.

MIXING INSTRUCTIONS

1. Fill spray tank 1/2 full of water.
2. With the agitator running, add the recommended amount of SFM 75.

3. If using a companion product, add the recommended amount.
4. For postemergent applications, add the proper amount of spray adjuvant (i.e. surfactants, drift control agents, etc.).
5. Add the remaining water.
6. Agitate the spray tank thoroughly.

Use the spray preparation within 24 hours to avoid product degradation. If the spray preparation is left standing, agitate it thoroughly before using.

SPRAYER CLEANUP

Thoroughly clean all mixing and spray equipment following applications of SFM 75 as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water.
2. Fill the tank with clean water and 1 gal of household ammonia (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
Equivalent amounts of an alternate-strength ammonia solution or a commercial cleaner can be used in the cleanout procedure. If a commercial cleaner is used, carefully read and follow the individual cleaner instructions.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. Dispose of the rinsate on a labeled site or at an approved waste disposal facility. If a commercial cleaner is used follow the directions for rinsate disposal on the label.

Notes:

1. **Caution:** Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended before performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When SFM 75 is tank mixed with other pesticides, all required cleanout procedures should be examined and the most rigorous procedure should be followed.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 – 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind Temperature and Humidity, and Temperature Inversions sections of this label.

Controlling Droplet Size – General Techniques

-Volume- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

-Pressure- Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**

-Nozzle Type- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size – Aircraft

-Number of Nozzles- Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.

-Nozzle Orientation- Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.

8/8

SFM 75

Specimen Label

-Nozzle Type – Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

Boom Length and Height

-Boom Length (aircraft) – For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.

-Boom Height (aircraft)- Application more than 10 ft above the canopy increases the potential for spray drift.

-Boom Height (ground)-Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

IMPORTANT PRECAUTIONS

Injury to or loss of desirable trees or other plants may result from failure to observe the following:

-If equipment is drained or flushed on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

- Treatment of powdery, dry soil or light, sandy soil when there is little likelihood of rainfall soon after treatment may result in off target movement and possible damage to susceptible crops when soil particles are moved by wind or water. Do not apply SFM 75 when these conditions are identified and powdery, dry soil or light sandy soil are known to be prevalent in the area to be treated. Injury to crops may result if treated soil is washed, blown, or moved onto land used to produce crops. Exposure to SFM 75 may injure or kill most crops. Injury may be more severe when the crops are irrigated.

-Applications made where runoff water flows onto agricultural land may injure crops. Applications, made during periods of intense rainfall, to soils saturated with water, surfaces paved with materials such as asphalt or concrete, or soils through which rainfall will not readily penetrate may result in runoff and movement of SFM 75. Do not treat frozen soil. Treated soil should be left undisturbed to reduce the potential for SFM 75 movement by soil erosion due to wind or water.

Do not use on lawns, walks, driveways, tennis courts, or similar areas.

Keep from contact with fertilizers, insecticides, fungicides, and seeds.

Do not apply in or on irrigation ditches or canals including their outer banks.

Do not apply through any type of irrigation system.

Do not use the equipment (tanks, pumps, hoses, booms, etc.) used to mix or spray SFM 75 for applications on crops or ornamentals. The mixing and application equipment may be used for forestry and noncrop applications only. This is extremely important as low rates of SFM 75 can kill or severely injure most crops.

If noncrop or forested sites treated with SFM 75 are to be converted to a food, feed, or fiber agricultural crop, or to a horticultural crop, do not plant the treated sites for at least one year after the SFM 75 application. To avoid damage to crops planted in these areas, and to ensure complete SFM 75 dissipation in treated sites, soil samples should be quantitatively analyzed, and a bioassay should be conducted before planting.

Do not use this product in the following counties of Colorado: Saguache, Rio Grande, Alamosa, Costilla and Conejos.

STORAGE AND DISPOSAL

Storage: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

Product Disposal: Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: Read the information below before using this product. If the terms are not acceptable, you should return the unopened product container immediately for a complete refund.

LIMITED WARRANTY, TERMS OF SALE, AND LIMITATION OF LIABILITY

Upon purchase or use of this product, purchaser and user agree to the following terms:
Warranty: Woodlands Specialists (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. The Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. All such risks are assumed by the user.

Limitation of Liability: The exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. Under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

1. Accord is a registered trademark of Monsanto Company.
2. Velpar L, Velpar DF, KARMEK DF, HYVAR X, KROVAR I DF, TELAR, Escori Hyvar SL Velpar are registered trademarks of E. I. DuPont de Nemours and Company.
3. Arsenal is a registered trademark of American-Cyanamide Company.
4. Aalrex is a registered trademark of Ciba-Geigy Corporation.

For questions call 334-872-6001

Woodlands Specialists Inc.
Hangar 209
Craig Field
Selma Alabama 36701