

81803-6

11/18



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511C)
1200 Pennsylvania Avenue NW
Washington, DC 20460

EPA Reg. Number:
81803-6

Date of Issuance:
APR 16 2008

NOTICE OF PESTICIDE:

Registration Re-registration
(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:
Jet-Ag

Name and Address of Registrant (include ZIP Code):

Jet Harvest Solutions
P.O. Box 915139
Longwood, FL 32791

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA Sec. 3(c) (5) provided you:

1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the following label change before you release the product for shipment: Revise the EPA Registration Number to read, "EPA Reg. No. 81803-6."
3. Submit three (3) copies of you final printed labeling before you release the product for shipment.

Signature of Approving Official:

W. Michael McDavit

Date:

4-16-08

W. Michael McDavit, Associate Director
Biopesticides and Pollution
Prevention Division (7511P)

EPA Form 8570-6

CONCURRENCES

SYMBOL	▶ 7511P						
SURNAME	▶ COLE	Foley					
DATE	▶ 4/11/08	4-16					

2/8

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Sincerely,

W. Michael McDavit, Associate Director
Biopesticides and Pollution
Prevention Division (7511P)

Enclosure

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

JET-AG

Jet-Ag is a liquid horticulture, algacide and fungicide used in horticulture on the following, such as greenhouse surfaces and equipment, for reasonably clean, non-porous surfaces, evaporative coolers, irrigation systems.

Use **Jet-Ag** as a fungicide, as a pre-plant dip treatment, as a soil drench on potting soil and growing mediums.

Use **Jet-Ag** as a foliar spray treatment in greenhouse and in the field for the reduction of plant pathogens.

Use **Jet-Ag** for the control of algae, fungi and bacteria on turf and seed.

Use **Jet-Ag** to treat soil prior to the inoculation with beneficial microorganisms.

Do not apply this product through any irrigation system unless directed by the label. Refer to Chemigation Directions for Use.

KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panel for additional precautionary statements.

For Commercial Horticultural Use Only

Active Ingredients:

Hydrogen Peroxide	26.5%
Peroxyacetic Acid	4.9%
Other Ingredients:.....	68.6%
Total:	100.0%



Jet Harvest Solutions
P.O. Box 915139
Longwood, FL 32791

EPA Registration No. (pending as File Symbol 81803-A)
 EPA Est. No. 54289-0H-OOI

Net Contents: _____ gal. (_____ fl. oz.)

ACCEPTED

APR 16 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 81803-6

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

Precautionary Statements
Hazards to Humans and Domestic Animals

DANGER - Corrosive. Concentrate causes irreversible eye damage. Concentrate may be fatal if swallowed or absorbed through skin. Concentrate causes skin burns or temporary discoloration on exposed skin. Do not breathe vapor of concentrate. Do not get concentrate in eyes, on skin or on clothing. Wear protective eyewear such as goggles or face shield. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE) – Applicators and handlers must wear coveralls over long-sleeved shirt, long pants, and chemical resistant footwear plus socks. When mixing and loading wear a chemical resistant apron. For overhead exposure wear chemical-resistant headgear. Wear protective eyewear (goggles, face shield, or safety glasses), and chemical resistant gloves. When cleaning equipment wear a chemical resistant apron. Follow manufacturer’s instructions for cleaning / maintaining PPE. If no such instruction exists for washables, use detergent and hot water.

User Safety Recommendations

User should wash hands thoroughly with soap and water before eating, drinking or using tobacco or using the toilet. Users should remove clothing immediately if contaminated by pesticide. Wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards - This pesticide is toxic to birds and fish. Do not contaminate water when disposing of equipment washwaters or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

This product is highly toxic to bees and other beneficial insects exposed to direct contact on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Do not apply this product or allow it to drift to crops where beneficials are part of an Integrated Pest Management strategy.

Physical or Chemical Hazards - Corrosive. Strong oxidizing agent. Do not use in concentrated form. Mix only with water in accordance with label instructions. Never bring concentrate in contact with other pesticides, cleaners or oxidative agents.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains the requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted-Entry Interval (REI). The requirements in this box apply to the uses of this product that are covered by the Worker Protection Standard.

For enclosed environments:

There is a restricted entry of one (1) hour for this product when applied via fogging or spraying to growing plants, surfaces, equipment, structures and non-porous surfaces in enclosed environments such as glasshouses and greenhouses. PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls worn over

long-sleeved shirt and pants, waterproof gloves and shoes plus socks.

There is a restricted entry of zero (0) hours for pre-plant dip, seed treatment, soil drench, mop, sponge, dip, soak, rinse or other non-spraying or fogging application methods when used in enclosed environments such as glasshouses and greenhouses.

For field applications:

Keep unprotected persons out of treated areas until sprays have dried.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are **not** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for the pesticide regulation.

To prevent, suppress, or eliminate Algae – *Alternaria* – Anthracnose – *Aphanomyces* – Black Spot – *Botrytis* (grey mold) – Downy Mildew – *Erwinia*- *Fusarium* (root rot) – Leaf Spot – *Phytophthora* (blights, rots) – *Plasmopara* – Powdery Mildew – *Pseudomonas* – *Pythium* – *Rhizoctonia* – Rust – Scab – Smut – *Thielaviopsis* – *Uncinula* (powdery mildew) – Wilts and Blights.

Compatibility

Jet-Ag™ has been formulated to provide a balanced source of the active ingredient directly to the plant surface and has been shown not to cause adverse cosmetic effects on most plants. Since not all plant species have been tested, it is always advisable to test on a few plants before treating large numbers.

Solution Preparation

Jet-Ag™ works best when diluted with water containing low levels of organic or inorganic materials and having a neutral pH. Thoroughly rinse out mixing tank with water before mixing concentrate. **Jet-Ag™** will readily mix with clean, neutral water and does not require agitation. **Jet-Ag™** must **not** be combined



5/8

or mixed with any other pesticide or fertilizer. **Jet-Ag™** can be used with surfactants for plants with waxy or hairy surfaces.

Use Rates and Directions

For greenhouse surfaces and equipment, use **Jet-Ag™** to suppress/control fungi and slime forming algae on greenhouse structures such as glazing, plastic, benches, walkways, floors, walls, fan blades, ventilation ducts, watering systems, coolers, storage rooms and equipment.

1. Sweep and remove all plant debris. Use power sprayer to wash all surfaces to remove loose dirt.
2. Use 7.8 ounce per 5 gallons of clean water. Use additional surfactant if needed.
3. Apply solution with mop, sponge, power sprayer or fogger to thoroughly wet all surfaces.
4. Scrub off heavy growths of algae and fungi following application. Use a solution of **Jet-Ag™** to wash away dead growth.
5. Reapply as often as needed for control.

For reasonably clean, non-porous surfaces, use 1.3 to 1.8 fluid ounces per 5 gallons of clean water. Use additional surfactant, if needed.

For evaporative coolers, treat cooler water every week with 0.7 to 1.3 fluid ounces per 5 gallons of cooler water.

For irrigation systems (flooded floors, flooded benches, recycled water systems, humidification and misting systems), treat already contaminated water with 0.7 to 2.6 fluid ounces per 5 gallons of water. Treat clean water with 0.5 to 0.7 gallon of **Jet-Ag™** per 10,000 gallons of water.

As a pre-plant dip treatment, use **Jet-Ag™** for the control/suppression of damping-off, root and stem rot diseases caused by *Pythium*, *Phytophthora*, *Rhizoctonia*, *Fusarium* or *Thielaviopsis* on ornamental and nursery plants, seed beds, seeds, seedlings, bulbs or cuttings.

1. Use 39.5 to 78.0 fluid ounces per 50 gallons of water.
2. Immerse plants or cuttings, remove and allow draining. Do not rinse.

As a soil drench, **Jet-Ag™** is effective for the control/suppression of soil borne plant diseases such as *Pythium*, *Phytophthora*, *Rhizoctonia*, *Fusarium* and *Thielaviopsis*. Use as a soil drench at the time of seeding or transplanting, as well as periodic drench. Use **Jet-Ag™** on potting soil and growing mediums prior to planting.

1. Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water.

2. Apply to soil or growing media to the point of saturation.
3. Wait fifteen minutes before planting or watering.

As a foliar spray treatment in greenhouses, **Jet-Ag™** works immediately on contact with any plant surface for control/suppression of plant pathogens. Apply **Jet-Ag™** to ornamentals, bedding plants, flowering plants, shrubs and trees.

Initial Curative Application:

1. Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water. Do not reuse already mixed solution; make fresh daily.
2. Spray, mist or fog plants in the early morning or late evening.
3. Thoroughly wet all surfaces of plant, upper and lower foliage, including stems, branches and stalks to ensure full contact with plant and flower tissue.
4. Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application.

Weekly Preventative Treatment:

1. Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water.
2. Spray, mist or fog plants.
3. Thoroughly wet all surfaces of plant, upper and lower foliage, including stems, branches and stalks.
4. Spray every five to seven days as a preventative treatment.
5. At the first sign of disease, spray daily with 3.6 to 7.8 fluid ounces per 5 gallons of water for three consecutive days and then resume weekly preventative treatment.

As a foliar spray treatment in the field, **Jet-Ag™** works immediately on contact with any plant surface for control / suppression of disease. Apply **Jet-Ag™** to nursery stock such as woody ornamentals, bedding plants, flowering plants, roses, container plants, azaleas, rhododendrons, conifers and shade trees. Complete coverage and wetting of the foliage is necessary.

Initial Curative Application:

1. Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water. Do not reuse already mixed solution; make fresh daily.
2. Spray, mist or fog plants and trees including application through irrigation or chemigation systems. If application is to be made through irrigation or chemigation systems, refer to the Chemigation Directions for Use section of this label for further requirements and instructions.
3. Thoroughly wet all surfaces of plant, upper and lower foliage, including stems, branches and stalks to ensure full contact with plant and flower tissue.
4. Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application.

Weekly Preventative Treatment:

1. Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water.
2. Spray, mist or fog plants and trees, including application through irrigation or chemigation systems.
3. Thoroughly wet all surfaces of plant, upper and lower foliage, including stems, branches and stalks.
4. Spray every five to seven days as a preventative treatment.
5. At the first sign of disease, spray daily with 3.6 to 7.8 fluid ounces per 5 gallons of water for three consecutive days and then resume weekly preventative treatment.

For cut flowers, use Jet-Ag™ to prevent *Botrytis*, Downy Mildew and Powdery Mildew on flowers in cold storage or in transit. Apply as a post harvest treatment. Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water. Spray flowers after grading and prior to storage or shipment. Repeat weekly for flowers in storage.

For bareroot nursery stock, use Jet-Ag™ to prevent *Botrytis* on bud wood and nursery stock in storage. Use 7.8 fluid ounces per 5 gallons of water. Dip plants or spray until dripping wet. Repeat weekly if necessary.

Turf Applications

Broad spectrum treatment for the control of algae, fungi and bacteria on turf. For use on all turf types such as commercial turf, lawns, athletic fields and golf course fairways, greens and tees.

Use Jet-Ag™ to control *Anthrachnose*, Brown Spot, Dollar Spot, Copper Spot, Fairy Ring, Pink Snow Mold, *Pythium*, *Phytophthora*, Summer Patch, *Rhizoctonia*, Scum, *Fusarium* Blight, Stripe Smut, Leaf Spot, Algae, Slime Molds and their spores.

Jet-Ag™ controls on contact.

For algacide/bactericide treatment, use Jet-Ag™ to control algae and bacterial diseases and the odors and the conditions these organisms may cause.

1. Typical treatment rates involve treating approximately 1,000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until run-off, saturation of the entire area will help ensure the solution penetrates algal crusts and deposits. Add a spreader surfactant for best results.
2. For initial (curative) treatment of heavy infestations of algae or bacterial disease, dilute 3.9 to 15.5 fluid ounces in 5 gallons of clean water. Apply 5-10 gallons of

diluted solution per 1000 square feet.

3. For preventative treatment of algae and bacterial disease, dilute 1.56 to 39.0 fluid ounces in 5 gallons of clean water. Apply 1 to 5 gallons of diluted solution per 1000 square feet.
4. Repeat application every 5 to 7 days or as needed to control new or established disease conditions. For best results, apply immediately after grass has been cut.

Optimum treatment time is early morning or late afternoon. Applications can be made during wet or rainy weather. Use the spray solution the same day it is prepared; do not store and reuse mixed spray solution. Jet-Ag™ can be injected through automatic irrigation systems in turf areas. Refer to Chemigation Directions for Use for specific instructions on using this product through irrigation systems.


For severe conditions of crusted algae, use Jet-Ag™ diluted at 14 to 25 fluid ounces per 5 gallons of clean water, and apply to 1000 square feet of affected area. Severe conditions require increased rates of active ingredient and increases in water volume to help penetrate layers of algae. Under severe conditions, double applications either by increasing the amount of active ingredient per 1000 square feet of turf or by applying twice over the same area.

For fungicide treatment of turf, use on golf course fairways, greens and tees of Bent grass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass and their mixtures.

1. Typical treatment rates involve treating approximately 1,000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until run-off. Add a spreader surfactant for best results.
2. Start applications at the first sign of disease and repeat every 5 to 7 days or as needed to control new or established disease condition. For best results, apply uniformly over the area immediately after grass has been cut.
3. For initial (curative) treatment of heavy infestation of fungal disease, dilute 1.80 to 7.80 fluid ounces in 5 gallons of clean water. Apply 5 to 10 gallons of dilute solution per 1000 square feet.
4. For preventative treatment of fungal disease, dilute 1.8 to 8.10 fluid ounces per 5 gallons of clean water. Apply 1-5 gallons of dilute solution per 1,000 square feet.

Optimum treatment time is early morning or late afternoon. Applications can be made during wet or rainy weather. Use spray solution the same day it is prepared; do not store mixed spray solution for later use.

For seedbed treatment, prior to sowing seed, use 2.5 fluid ounces per 5 gallons of clean water. Thoroughly wet or drench the seed bed, to the point of saturation, with 60 to 100 gallons of dilute solution per 1,000 square feet. Let sit for one hour then



immediately seed soil.

After seeds have germinated, use 0.8 to 1.2 fluid ounces per 5 gallons of clean water. Lightly spray or irrigate the soil and seedlings until thoroughly wetted. Retreat once per week until seed is well established.

For soil treatment prior to inoculation with beneficial microorganisms, use Jet-Ag™ to reduce the number of plant pathogenic microorganisms in the soil. Use 1.3 to 2.5 fluid ounces per 5 gallons of clean water. Thoroughly wet or drench the area to be inoculated. Wait one day before inoculating the soil.

To treat turf following inoculation of soil with beneficial microorganisms, use Jet-Ag™ to control plant pathogens on the foliar portion of turf. Do not drench the root system, or a temporary reduction in beneficial soil microorganisms can occur.

Use 0.8 to 1.9 fluid ounces per 5 gallons of clean water. Apply to the turf by lightly spraying leaf surfaces. Do not allow solution to be drenched into the soil and root systems. Drenching of Jet-Ag™ into the soil can result in temporary reduction of beneficial microorganisms.

CHEMIGATION DIRECTIONS FOR USE

General Requirements

1. Apply this product through one of the following types of irrigation systems: center pivot, lateral move, end tow, side wheel roll, traveler, solid set, and hand move, flood basin or drip trickle irrigation system. Do not apply this product through any other type of irrigation system.
2. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
3. Ensure that the irrigation system is properly calibrated. If you have questions about calibration, contact the State Extension Service specialist, the equipment manufacturer, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless proper safety devices for public water systems are in place. Read specific requirements provided below.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

Specific Requirements for all Irrigation Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or

- regularly serves an average of at least 25 individuals daily at least 60 days throughout the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back towards the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump or equivalent, effectively designed and constructed of materials that are compatible with pesticide and capable of being filled with a system interlock.
7. Do not apply outdoors when wind speed favors drift beyond the area intended for treatment.

Application Instructions

1. Remove scale, pesticide residues and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residue may cause product to lose effectiveness of strength.
2. Determine the treatment rates as indicated in the directions for use and make proper dilutions.
3. Prepare a solution by filling the tank with the required volume of water and then adding product as required. The product will immediately go into solution without any required agitation.
4. Jet-Ag™ Liquid Agriculture and Horticulture must not be applied in conjunction with any other pesticides or fertilizers; this may cause reduced performance of the product and should be avoided.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers,

hospitals, in-patient clinics, nursing homes or any public area such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.


Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words "PESTICIDES IN IRRIGATION WATER."

forth, ensuring at least one complete revolution, for 30 seconds. Stand container on its end and tip back and for the several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat the procedure two more times. Then offer for recycling or dispose in a sanitary landfill, or incineration, if allowed by state and local authorities by burning.

WARRANTY

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing, method of application, weather, watering practices, and nature of soil, potting medium, disease problem, and condition of crop, incompatibility with other chemicals, pre-existing conditions and other conditions influencing the use of this product are beyond the control of the seller. Buyer assumes all risks associated with the use, storage, or handling of this material not in strict accordance with directions given herewith. TO THE EXTENT ALLOWABLE BY STATE LAW, NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.



Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original containers in a cool, well-vented area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip the container on its side and roll it back and