

81803-6

9/29/2011

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

September 29, 2011

OFFICE OF
PREVENTION PESTICIDES AND
TOXIC SUBSTANCES

Austin Grant, Inc
c/o William F Stoneman
W F Stoneman Company, LLC
P O Box 465
McFarland, WI 53558-0465

Subject Application for Label Notification to change the net contents from 4 35 and 53 gallons to 5, 53, and 265 gallons
Jet-Ag
EPA Reg No 81803-6
Your submission dated September 9, 2011
Decision Number 455100

Dear Mr Stoneman

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under PR Notice 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PR Notice 98-10 and it has been determined that the action(s) requested falls within the scope of PR Notice 98-10. Our records have been duly noted and the application has been stamped "Notification Accepted"

If you have any questions concerning this action, please feel free to contact Mr Colin Walsh at (703) 308-0298 or via email at walsh.colin@epa.gov

Sincerely,

Linda A Hollis

Linda A Hollis, Chief
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention
Division (7511P)



United States Environmental Protection Agency Washington DC 20460

Registration Amendment Other

OPP Identifier Number

Application for Pesticide - Section I

1 Company/Product Number 81803-6 2 EPA Product Manager Linda Hollis Branch Chief/ BPPD 3 Proposed Classification None Restricted 4 Company/Product (Name) JET-AG PM# Biochemical Pesticides Branch 5 Name and Address of Applicant (Include ZIP Code) Austin Grant Inc DBA Jet Harvest Solutions PO Box 9151139 Longwood FL 32791-5139 6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No Product Name

Section - II

Amendment Explain below Resubmission in response to Agency letter dated Notification Explain below Final printed labels in response to Agency letter dated "Me Too" Application Other Explain below Notification Accepted Date SEP 29 2011

Explanation Use additional page(s) if necessary (For section I and Section II) Reviewer C Walsh Please note that the container sizes for JET AG (EPA Reg No 81803 6) are changing from the previous 53 gallon or 4 35 gallon containers to these three container sizes 53 gallons 5 gallons 265 gallons All containers will be appropriately labeled as required

Section - III

1 Material This Product Will Be Packaged In Child Resistant Packaging Unit Packaging Water Soluble Packaging 2 Type of Container Metal Plastic Glass Paper Other (Specify) 3 Location of Net Contents Information Label Container 4 Size(s) Retail Container 53 gallons 5 gallons 265 gallons 5 Location of Label Directions On each container 6 Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Other

Section - IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application) Name William F Stoneman W F Stoneman Company LLC Title Regulatory Consultant Telephone No (Include Area Code) 606 268 7040 or 608 712 4916 Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law 2 Signature William F. Stoneman 3 Title Regulatory Consultant 4 Typed Name William F Stoneman 5 Date September 9th, 2011 6 Date Application Received (Stamped)



Jet Ag is a liquid horticulture algacide and fungicide used in horticulture on the following such as greenhouse surfaces and equipment for reasonably clean non porous surfaces, evaporative coolers irrigation systems

Use **Jet Ag** as a fungicide as a pre plant dip treatment as a soil drench on potting soil and growing mediums

Use **Jet Ag** as a foliar spray treatment in greenhouse and in the field for the reduction of plant pathogens

Use **Jet Ag** for the control of algae fungi and bacteria on turf and seed

Use **Jet Ag** to treat soil prior to the inoculation with beneficial microorganisms

Do not apply this product through any irrigation system unless directed by the label Refer to Chemigation Directions for Use

KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

See side panel for additional precautionary statements

For Commercial Horticultural Use Only

| | |
|--------------------|--------|
| Active Ingredients | |
| Hydrogen Peroxide | 26.5% |
| Peroxyacetic Acid | 4.9% |
| Other Ingredients | 68.6% |
| Total | 100.0% |



Jet Harvest Solutions
P O Box 915139
Longwood, FL 32791

EPA Registration No 81803 6 EPA Est No XXXXXX

Net Contents 53 gallons or 4 35 gallons or 30 gallons

| FIRST AID | |
|--|--|
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 – 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 – 20 minutes • Call a poison control center or doctor for treatment advice |
| If swallowed | <ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible • Call a poison control center or doctor for treatment advice |
| HOTLINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 800 222 1222 for emergency medical treatment information | |
| NOTE TO PHYSICIAN | |
| Probable mucosal damage may contraindicate the use of gastric lavage | |

Precautionary Statements
Hazards to Humans and Domestic Animals

DANGER Corrosive Concentrate causes irreversible eye damage Concentrate may be fatal if swallowed or absorbed through skin Concentrate causes skin burns or temporary discoloration on exposed skin Do not breathe vapor of concentrate Do not get concentrate in eyes on skin or on clothing Wear protective eyewear such as goggles or face shield Wash thoroughly with soap and water after handling Remove and wash contaminated clothing before reuse

Personal Protective Equipment (PPE) – Applicators and handlers must wear coveralls over long sleeved shirt long pants and chemical resistant footwear plus socks When mixing and loading wear a chemical resistant apron For overhead exposure wear chemical resistant headgear Wear protective eyewear (goggles face shield or safety glasses) and chemical resistant gloves When cleaning equipment wear a chemical resistant apron Follow manufacturer s instructions for cleaning / maintaining PPE If no such instruction exists for washables use detergent and hot water



User Safety Recommendations

User should wash hands thoroughly with soap and water before eating drinking or using tobacco or using the toilet Users should remove clothing immediately if contaminated by pesticide Wash thoroughly and put on clean clothing Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

Environmental Hazards This pesticide is toxic to birds and fish Do not contaminate water when disposing of equipment washwaters or rinsate Exposed treated seed may be hazardous to birds and other wildlife Dispose of all excess treated seed and seed packaging by burial away from bodies of water

This product is highly toxic to bees and other beneficial insects exposed to direct contact on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area Do not apply this product or allow it to drift to crops where beneficials are part of an Integrated Pest Management strategy

Physical or Chemical Hazards Corrosive Strong oxidizing agent Do not use in concentrated form Mix only with water in accordance with label instructions Never bring concentrate in contact with other pesticides cleaners or oxidative agents

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170 This standard contains the requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and Restricted Entry Interval (REI) The requirements in this box apply to the uses of this product that are covered by the Worker Protection Standard

For enclosed environments

There is a restricted entry of one (1) hour for this product when applied via fogging or spraying to growing plants surfaces equipment structures and non porous surfaces in enclosed environments such as glasshouses and greenhouses PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is coveralls worn over

long sleeved shirt and pants waterproof gloves and shoes plus socks

There is a restricted entry of zero (0) hours for pre plant dip seed treatment soil drench mop sponge dip soak rinse or other non spraying or fogging application methods when used in enclosed environments such as glasshouses and greenhouses

For field applications

Keep unprotected persons out of treated areas until sprays have dried

Non Agricultural Use Requirements

The requirements in this box apply to uses of this product that are **not** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Keep unprotected persons out of treated areas until sprays have dried

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for the pesticide regulation

To prevent suppress or eliminate Algae – *Alternaria* – *Anthraco*se – *Aphanomyces* – Black Spot – *Botrytis*(grey mold) – Downy Mildew – *Erwinia Fusarium* (root rot) – Leaf Spot – *Phytophthora* (blights rots) – *Plasmopara* – Powdery Mildew – *Pseudomonas* – *Pythium* – *Rhizoctonia* – Rust – Scab – Smut – *Thielaviopsis* – *Uncinula* (powdery mildew) – Wilts and Blights

Compatibility

Jet Ag™ has been formulated to provide a balanced source of the active ingredient directly to the plant surface and has been shown not to cause adverse cosmetic effects on most plants Since not all plant species have been tested it is always advisable to test on a few plants before treating large numbers

Solution Preparation

Jet Ag™ works best when diluted with water containing low levels of organic or inorganic materials and having a neutral pH Thoroughly rinse out mixing tank with water before mixing concentrate **Jet Ag™** will readily mix with clean neutral water and does not require agitation **Jet Ag™** must **not** be combined or mixed with any other pesticide or fertilizer **Jet Ag™** can be used with surfactants for plants with waxy or hairy surfaces

JET-AG

Use Rates and Directions

For greenhouse surfaces and equipment use Jet Ag™ to suppress/control fungi and slime forming algae on greenhouse structures such as glazing plastic benches walkways floors walls fan blades ventilation ducts watering systems coolers storage rooms and equipment

- 1 Sweep and remove all plant debris Use power sprayer to wash all surfaces to remove loose dirt
- 2 Use 7.8 ounce per 5 gallons of clean water Use additional surfactant if needed
- 3 Apply solution with mop sponge power sprayer or fogger to thoroughly wet all surfaces
- 4 Scrub off heavy growths of algae and fungi following application Use a solution of Jet Ag™ to wash away dead growth
- 5 Reapply as often as needed for control

For reasonably clean, non porous surfaces use 1.3 to 1.8 fluid ounces per 5 gallons of clean water Use additional surfactant if needed

For evaporative coolers treat cooler water every week with 0.7 to 1.3 fluid ounces per 5 gallons of cooler water

For irrigation systems (flooded floors flooded benches recycled water systems humidification and misting systems) treat already contaminated water with 0.7 to 2.6 fluid ounces per 5 gallons of water Treat clean water with 0.5 to 0.7 gallon of Jet Ag™ per 10,000 gallons of water

As a pre plant dip treatment, use Jet Ag™ for the control/suppression of damping off root and stem rot diseases caused by *Pythium Phytophthora Rhizoctonia Fusarium or Thielaviopsis* on ornamental and nursery plants seed beds seeds seedlings bulbs or cuttings

- 1 Use 39.5 to 78.0 fluid ounces per 50 gallons of water
- 2 Immerse plants or cuttings remove and allow draining Do not rinse

As a soil drench Jet Ag™ is effective for the control/suppression of soil borne plant diseases such as *Pythium Phytophthora Rhizoctonia Fusarium* and *Thielaviopsis* Use as a soil drench at the time of seeding or transplanting as well as periodic drench Use Jet Ag™ on potting soil and growing mediums prior to planting

- 1 Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water
- 2 Apply to soil or growing media to the point of saturation
- 3 Wait fifteen minutes before planting or watering

As a foliar spray treatment in greenhouses Jet Ag™ works immediately on contact with any plant surface for control/suppression of plant pathogens Apply Jet Ag™ to ornamentals bedding plants flowering plants shrubs and trees

Initial Curative Application

- 1 Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water Do not reuse already mixed solution make fresh daily
- 2 Spray mist or fog plants in the early morning or late evening
- 3 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks to ensure full contact with plant and flower tissue
- 4 Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application

Weekly Preventative Treatment

- 1 Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water
- 2 Spray mist or fog plants
- 3 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks
- 4 Spray every five to seven days as a preventative treatment
- 5 At the first sign of disease spray daily with 3.6 to 7.8 fluid ounces per 5 gallons of water for three consecutive days and then resume weekly preventative treatment

As a foliar spray treatment in the field, Jet Ag™ works immediately on contact with any plant surface for control / suppression of disease Apply Jet Ag™ to nursery stock such as woody ornamentals bedding plants flowering plants roses container plants azaleas rhododendrons conifers and shade trees Complete coverage and wetting of the foliage is necessary

Initial Curative Application

- 1 Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water Do not reuse already mixed solution make fresh daily
- 2 Spray mist or fog plants and trees including application through irrigation or chemigation systems If application is to be made through irrigation or chemigation systems refer to the Chemigation Directions for Use section of this label for further requirements and instructions
- 3 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks to ensure full contact with plant and flower tissue
- 4 Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application



Weekly Preventative Treatment

- 1 Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water
- 2 Spray mist or fog plants and trees including application through irrigation or chemigation systems
- 3 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks
- 4 Spray every five to seven days as a preventative treatment
- 5 At the first sign of disease spray daily with 3.6 to 7.8 fluid ounces per 5 gallons of water for three consecutive days and then resume weekly preventative treatment

For cut flowers, use **Jet Ag™** to prevent *Botrytis* Downy Mildew and Powdery Mildew on flowers in cold storage or in transit Apply as a post harvest treatment Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water Spray flowers after grading and prior to storage or shipment Repeat weekly for flowers in storage

For bareroot nursery stock use **Jet Ag™** to prevent *Botrytis* on bud wood and nursery stock in storage Use 7.8 fluid ounces per 5 gallons of water Dip plants or spray until dripping wet Repeat weekly if necessary

Turf Applications

Broad spectrum treatment for the control of algae fungi and bacteria on turf For use on all turf types such as commercial turf lawns athletic fields and golf course fairways greens and tees

Use **Jet Ag™** to control *Anthraco*se Brown Spot Dollar Spot Copper Spot Fairy Ring Pink Snow Mold *Pythium* *Phytophthora* Summer Patch *Rhizoctonia* Scum *Fusarium* Blight Stripe Smut Leaf Spot Algae Slime Molds and their spores

Jet Ag™ controls on contact

For algaecide/bactericide treatment use **Jet Ag™** to control algae and bacterial diseases and the odors and the conditions these organisms may cause

- 1 Typical treatment rates involve treating approximately 1 000 square feet of lawn area with 1 to 10 gallons of diluted solution Spray entire area until run off saturation of the entire area will help ensure the solution penetrates algal crusts and deposits Add a spreader surfactant for best results
- 2 For initial (curative) treatment of heavy infestations of algae or bacterial disease dilute 3.9 to 15.5 fluid ounces in 5 gallons of clean water Apply 5 10 gallons of diluted solution per 1000 square feet

- 3 For preventative treatment of algae and bacterial disease dilute 1.56 to 39.0 fluid ounces in 5 gallons of clean water Apply 1 to 5 gallons of diluted solution per 1000 square feet
- 4 Repeat application every 5 to 7 days or as needed to control new or established disease conditions For best results apply immediately after grass has been cut

Optimum treatment time is early morning or late afternoon Applications can be made during wet or rainy weather Use the spray solution the same day it is prepared do not store and reuse mixed spray solution **Jet Ag™** can be injected through automatic irrigation systems in turf areas Refer to Chemigation Directions for Use for specific instructions on using this product through irrigation systems

For severe conditions of crusted algae use **Jet Ag™** diluted at 14 to 25 fluid ounces per 5 gallons of clean water and apply to 1000 square feet of affected area Severe conditions require increased rates of active ingredient and increases in water volume to help penetrate layers of algae Under severe conditions double applications either by increasing the amount of active ingredient per 1000 square feet of turf or by applying twice over the same area

For fruit and vegetable storage **Jet Ag™** can be applied by fogging to control the growth of non public health microorganisms that may cause decay and/or spoilage on raw post harvest fruits and vegetables during the post harvest process

- 1 Ensure room is properly ventilated Vacate all personnel from room during fogging and for a minimum of 2 hours after fogging Ensure there is no strong odor characteristic of acetic acid before having personnel return to work area Do not enter room until hydrogen peroxide concentrations are correctly tested and are below 1 ppm on a time weighted average
- 2 Fog areas using 3.5 fluid ounces into humidified air 38 pounds per square feet of fruit and vegetables per 1 000 cu ft of room volume Inject concentrate into water used in humidification of postharvest fruit and vegetables in storage using any type of fogging equipment including but not limited to cold foggers thermal foggers low pressure air assisted and high pressure fog systems

For the fungicide treatment control or suppression **Jet Ag™** can be applied to growing crops such as but not limited to berries citrus fruit pome fruit herbs spices peppers tomatoes onions

Initial Curative Application

- 5 Use 3.9 to 7.8 fluid ounces per 5 gallons of clean water Do not reuse already mixed solution make fresh daily

JET-AG

- 6 Spray mist or fog plants and trees including application through irrigation or chemigation systems. If application is to be made through irrigation or chemigation systems refer to the Chemigation Directions for Use section of this label for further requirements and instructions
- 7 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks to ensure full contact with plant tissue
- 8 Apply for one to three consecutive days and then follow directions for preventative treatment after the initial application

Weekly Preventative Treatment

- 6 Use 0.75 to 1.3 fluid ounces per 5 gallons of clean water
- 7 Spray mist or fog plants and trees including application through irrigation or chemigation systems
- 8 Thoroughly wet all surfaces of plant upper and lower foliage including stems branches and stalks
- 9 Spray every five to seven days as a preventative treatment
- 10 At the first sign of disease spray daily with 3.6 to 7.8 fluid ounces per 5 gallons of water for three consecutive days and then resume weekly preventative treatment

For fungicide treatment of turf, use on golf course fairways, greens and tees of Bent grass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St Augustinegrass and their mixtures

- 1 Typical treatment rates involve treating approximately 1,000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until run off. Add a spreader surfactant for best results
- 2 Start applications at the first sign of disease and repeat every 5 to 7 days or as needed to control new or established disease condition. For best results apply uniformly over the area immediately after grass has been cut
- 3 For initial (curative) treatment of heavy infestation of fungal disease dilute 1.8 to 7.8 fluid ounces in 5 gallons of clean water. Apply 5 to 10 gallons of dilute solution per 1,000 square feet
- 4 For preventative treatment of fungal disease dilute 1.8 to 8.1 fluid ounces per 5 gallons of clean water. Apply 1.5 gallons of dilute solution per 1,000 square feet

Optimum treatment time is early morning or late afternoon. Applications can be made during wet or rainy weather. Use spray solution the same day it is prepared. Do not store mixed spray solution for later use.

For seedbed treatment prior to sowing seed use 2.5 fluid ounces per 5 gallons of clean water. Thoroughly wet or drench the seed bed to the point of saturation with 60 to 100 gallons of dilute solution per 1,000 square feet. Let sit for one hour then immediately seed soil.

After seeds have germinated use 0.8 to 1.2 fluid ounces per 5 gallons of clean water. Lightly spray or irrigate the soil and seedlings until thoroughly wetted. Retreat once per week until seed is well established.

For soil treatment prior to inoculation with beneficial microorganisms use Jet Ag™ to reduce the number of plant pathogenic microorganisms in the soil. Use 1.3 to 2.5 fluid ounces per 5 gallons of clean water. Thoroughly wet or drench the area to be inoculated. Wait one day before inoculating the soil.

To treat turf following inoculation of soil with beneficial microorganisms use Jet Ag™ to control plant pathogens on the foliar portion of turf. Do not drench the root system or a temporary reduction in beneficial soil microorganisms can occur.

Use 0.8 to 1.9 fluid ounces per 5 gallons of clean water. Apply to the turf by lightly spraying leaf surfaces. Do not allow solution to be drenched into the soil and root systems. Drenching of Jet Ag™ into the soil can result in temporary reduction of beneficial microorganisms.

CHEMIGATION DIRECTIONS FOR USE

General Requirements

- 1 Apply this product through one of the following types of irrigation systems: center pivot, lateral move, end tow, side wheel roll, traveler, solid set, and hand move, flood basin, or drip/trickle irrigation system. Do not apply this product through any other type of irrigation system.
- 2 Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- 3 Ensure that the irrigation system is properly calibrated. If you have questions about calibration, contact the State Extension Service specialist, the equipment manufacturer, or other experts.
- 4 Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless proper safety devices for public water systems are in place. Read specific requirements provided below.
- 5 A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.



9/10

Specific Requirements for all Irrigation Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days throughout the year
- 2 Chemigation systems connected to public water systems must contain a functional reduced pressure zone (RPZ) backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least the inside diameter of the fill pipe
- 3 The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of liquid back towards the injection pump
- 4 The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down
- 5 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6 Systems must use a metering pump such as a positive displacement injection pump or equivalent effectively designed and constructed of materials that are compatible with pesticide and capable of being filled with a system interlock
- 7 Do not apply outdoors when wind speed favors drift beyond the area intended for treatment

fertilizers this may cause reduced performance of the product and should be avoided

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas labor camps businesses day care centers hospitals in patient clinics nursing homes or any public area such as schools parks playgrounds or other public facilities not including public roads or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses. Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

Application Instructions

- 1 Remove scale pesticide residues and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank void of scale or residue may cause product to lose effectiveness of strength
- 2 Determine the treatment rates as indicated in the directions for use and make proper dilutions
- 3 Prepare a solution by filling the tank with the required volume of water and then adding product as required. The product will immediately go into solution without any required agitation
- 4 Jet Ag™ Liquid Agriculture and Horticulture must not be applied in conjunction with any other pesticides or



Storage and Disposal

Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Store in original containers in a cool well vented area away from direct sunlight Do not allow product to become overheated in storage This may cause increased degradation of the product which will decrease product effectiveness In case of spill flood area with large quantities of water Do not store in a manner where cross contamination with other pesticides or fertilizers could occur

PESTICIDE DISPOSAL Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility Open dumping is prohibited If wastes cannot be disposed of according to label directions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL Nonrefillable container Do not reuse or refill this container Triple rinse (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or mix tank Fill the container ¼ full with water Replace and tighten closures Tip the container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand container on its end and tip back and for the several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use of disposal Repeat the procedure two more times Then offer for recycling or dispose in a sanitary landfill or incineration if allowed by state and local authorities by burning

WARRANTY

This material conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use Timing method of application weather watering practices and nature of soil potting medium disease problem and condition of crop incompatibility with other chemicals pre existing conditions and other conditions influencing the use of this product are beyond the control of the seller Buyer assumes all risks associated with the use storage or handling of this material not in strict accordance with directions given herewith TO THE EXTENT ALLOWABLE BY STATE LAW NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE

