	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

**Application for Pesticide - Section I**

1. Company/Product Number 81880-2	2. EPA Product Manager Jim Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) GWN-3061	PM# Team# 25, Phone: 703-305-5697	
5. Name and Address of Applicant (Include ZIP Code) Canyon Group c/o Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	--

**NOTIFICATION**  
SEP 23 2005

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)


-Notification of Minor Label revisions - correction of typographical errors, rewording for clarification purposes per PR-Notice 98-10 II.N.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

<b>1. Material This Product Will Be Packaged In:</b>				<b>2. Type of Container</b>	
<b>Child-Resistant Packaging</b> <input type="checkbox"/> Yes <input type="checkbox"/> No	<b>Unit Packaging</b> <input type="checkbox"/> Yes <input type="checkbox"/> No	<b>Water Soluble Packaging</b> <input type="checkbox"/> Yes <input type="checkbox"/> No			<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
<b>3. Location of Net Contents Information</b>		<b>4. Size(s) Retail Container</b>		<b>5. Location of Label Directions</b>	
<input type="checkbox"/> Label <input type="checkbox"/> Container				<input type="checkbox"/> _____	
<b>6. Manner in Which Label is Affixed to Product</b>			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		
			<input type="checkbox"/> Other _____		

**Section - IV**

<b>1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)</b>		
Name Rebecca A. Hargadine	Title Agent for Canyon Group	Telephone No. (Include Area Code) 928-819-1531
<b>Certification</b>		<b>6. Date Application Received (Stamped)</b>
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		_____ (Stamped)
<b>2. Signature</b> 	<b>3. Title</b> Agent for Canyon Group	
<b>4. Typed Name</b> Rebecca A. Hargadine	<b>5. Date</b> 08-30-05	

2/23

# GWN-3061

## Herbicide

NOTIFICATION

SEP 23 2005

GWN-3061 is a selective herbicide for the control of listed annual broadleaf weeds and nutsedge in field corn, field corn grown for seed, grain sorghum (milo), rice, sugarcane, fallow ground, cotton, and dry beans

	<b>% By Wt.</b>
ACTIVE INGREDIENT: *Halosulfuron-methyl.....	75.0%
OTHER INGREDIENTS.....	25.0%
	<b>TOTAL 100.0%</b>

### KEEP OUT OF REACH OF CHILDREN

## CAUTION!

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUSES MODERATE EYE IRRITATION. HARMFUL IF SWALLOWED. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after 5 minutes, then continue rinsing eye.</li> <li>• Call poison control center or physician for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call poison control center or physician immediately for treatment advice.</li> <li>• Remove visible particles from mouth.</li> <li>• Have person rinse mouth thoroughly with water, spit out rinse water.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or physician, or going for treatment.</p> <p><b>FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL TOLL FREE 1-888-478-0798</b></p> <p>This product is identified as GWN-3061, EPA Reg. No. 81880-xxx.</p>	

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

**Applicators and other handlers must wear:**

- long-sleeved shirt and long pants
- and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

EPA Reg. No. 81880-2  
EPA Est. No.

Canyon Group LLC c/o Gowan Company  
P.O. Box 5569  
Yuma, Arizona 85366-5569

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions For Use on this label or in separately published Canyon Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- shoes plus socks
- chemical-resistant gloves, such as nitrile rubber, neoprene rubber or polyethylene. For more options, follow instructions for category A (dry and water-based formulations) on an EPA chemical resistant category selection chart.

For more product information, call toll-free 1-800-888-1844.

### GENERAL INFORMATION

#### Biological Information

The level of weed control following GWN-3061 application is dependent upon application rate, weed species and size at application time, and growing conditions. For best results, applications should be made to actively growing weeds at the heights defined in the "USE RATE GUIDE" sections of this label. Heavy infestations should be treated early before the weeds become too competitive with the crop. When early post-emergence treatments are used (in corn), sequential applications may be required to control later weed flushes.

Soon after GWN-3061 is applied, growth of susceptible weeds is inhibited, and susceptible weeds are no longer competitive with the crop. Following growth inhibition, the leaves and growing point begin to discolor. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions.

### MIXING INSTRUCTIONS

Fill the spray tank to about three-fourths of the desired volume with water or carrier. Add the recommended amount of this product as listed in the "WEEDS CONTROLLED" sections. Complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Add nonionic surfactant and other adjuvants as the last ingredients in the tank.

Spray solutions should be applied within 24 hours after mixing.

**Adjuvants:** A nonionic surfactant (NIS) is the only adjuvant required in the spray solution. Use only nonionic surfactants which are approved by EPA for use on food crops and which contain at least 80 percent active ingredient. Use 0.25 to 0.5 percent nonionic surfactant concentration (1 to 2 quarts per 100 gallons of spray solution).

**Crop oil concentrate (COC)** may be used with GWN-3061 instead of nonionic surfactants. Do not use both NIS and COC in the spray mixture. Add COC to the spray mixture at 1% vol./vol. (1 gallon per 100 gallons of spray mixture). Use only good quality petroleum or vegetable-based crop oil concentrates which contain at least 14 percent emulsifiers.

4/23

Nonionic surfactant or COC are the only additives necessary for GWN-3061 applications. Liquid nitrogen fertilizer solution (e.g., 28-0-0) may be added to the spray solution to improve the control of certain species, particularly if GWN-3061 is being tank mixed with a companion herbicide which requires use of a liquid nitrogen additive. However, a nonionic surfactant or COC will still be necessary. Refer to the companion product label for specific additive requirements. Otherwise, add liquid nitrogen fertilizer at a rate of 2 to 4 quarts per acre. Do not use liquid nitrogen fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur. A high quality, spray grade ammonium sulfate (e.g., 21-0-0) may be applied at a rate of 2 to 4 pounds per acre in place of the liquid nitrogen fertilizer.

### APPLICATION EQUIPMENT AND INSTRUCTIONS

Applications may be made by ground [(or aerial) if corn, sorghum, or rice] equipment to healthy, actively growing weeds. For best results, avoid applications when weeds are under drought, stress, disease, or insect damage. Rainfall or irrigation occurring within 4 hours after application may also reduce effectiveness.

#### Ground Applications

Apply GWN-3061 uniformly with properly calibrated ground equipment in 10 or more gallons of water per acre. Other water based spray carriers may be used for directed applications, avoiding contact with crop foliage. Select spray volumes that ensure thorough and uniform weed coverage. Choose nozzles which provide optimum spray distribution and coverage at the appropriate pressure (psi). Use only ground application equipment. Thoroughly clean equipment prior to mixing spray solution. Avoid streaking, skips, overlaps, and spray drift during applications.

Do not apply this product through any type of irrigation system.

Avoid disturbing (e.g., cultivation) treated areas for at least 7 days following application.

Thoroughly clean application equipment immediately after GWN-3061 use and prior to spraying a crop other than corn or grain sorghum. Prepare a tank cleaning solution which consists of a 1 percent solution of household ammonia (one quart of ammonia for every 25 gallons of water). Use sufficient cleaning solution to thoroughly rinse all surface and to flush all hoses. Repeat the procedure with the ammonia solution. Complete the cleaning process by rinsing with clean water.

#### Aerial Applications [For Corn, Sorghum, & Rice]

**Aerial applications may only be made to Rice, Corn, and Sorghum.** Apply this product or approved tank mixtures with properly calibrated equipment in 3 to 15 gallons of water per acre.

Thoroughly clean equipment prior to mixing spray solution. Avoid streaking, skips, overlaps, and spray drift during applications.

#### Spray Drift Management

**AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.**

The interaction of many equipment – and weather – related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

#### The importance of spray droplet size:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but may not prevent drift if applications are made improperly or under unfavorable environmental conditions (see the following "Wind", "Temperature and Humidity", and "Temperature Inversion" sections of this advisory).

#### Controlling initial droplet size:

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher flow rates produce larger droplets.
- **Pressure** – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation** – Orienting nozzles so the spray stream is released backwards, parallel to the air stream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

- **Nozzle type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Controlling placement of spray droplets:

- **Boom length** – For some use patterns, reducing the effective boom length to less than ¾ of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application height** – Applications should not be greater than 10 feet above the top of the tallest plants unless a greater height is required for aircraft safety. Greater application heights result in greater droplet size reduction through evaporation and greater movement in air currents. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
- **Application speed**- Slower aircraft speeds within a safe range will produce less air turbulence and fewer small droplets.
- **Swath adjustment** – When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distances should increase with increasing drift potential (wind speed, droplet size, etc.).

Key environmental factors:

- **Wind** – Drift potential is the lowest between wind speeds of 2 to 10 mph. However, many factors including droplet size and equipment type determine drift potential at any given speed. Application should be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Applicators should be familiar with local wind patterns and how they affect drift.
- **Temperature and humidity** – When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.
- **Temperature inversions** – Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable air currents that are common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke detector. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive areas:**

Pesticides should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid disturbing (e.g., cultivation) treated areas for at least 7 days following application.

Thoroughly clean application equipment immediately after the use of GWN-3061. Prepare a tank cleaning solution that consists of a 1 percent solution of household ammonia (one quart of ammonia for every 25 gallons of water). Use sufficient cleaning solution to thoroughly rinse all surfaces and to flush all hoses. Repeat the procedure with the ammonia solution. Complete the cleaning process by rinsing with clean water.

**CALIFORNIA ONLY**

**Sensitive Crops:**

Cotton Prunes

**Buffer Zones:**

1. Aerial applications shall not be made closer than four miles from sensitive crops.
2. Ground applications shall not be made closer than 1 mile from sensitive crops unless wind direction during the application is away from sensitive crops. When wind direction during the ground application is away from sensitive crops ground applications shall not be made closer than 0.5 miles from sensitive crops.

**TANK MIXTURES**

This product may be applied in combination with other products that are registered for the same crop and application.

6/23

**Refer to the companion product label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated, and application restrictions.**

Before mixing in the spray tank, it is recommended that compatibility be tested by mixing all components in a small container in proportionate quantities. For tank mixtures, add individual formulations to a spray tank in the following sequence: water soluble bags, dry flowables, emulsifiable concentrates, drift control additive, water soluble liquids followed by nonionic surfactant or crop oil concentrate.

Tank mixtures should not be applied if the crop is under severe stress due to drought, water-saturated soils, poor fertility (especially low nitrogen levels, hail, frost, insects or when the maximum daytime temperature is above 92° F. Tank-mix applications under these conditions may cause temporary crop injury.

**ROTATIONAL CROP INFORMATION**

Labeled crops may be planted at specified time intervals following application of approved rates of GWN-3061. Use the time intervals listed below to determine the required time interval before planting.

**TIME INTERVAL BEFORE PLANTING  
(Months after treatment with GWN-3061)**

<b>Crop</b>	<b>Months</b>
IR/LMR Field corn	0
Sugarcane	0
IT Field corn	1
Normal Field corn	1
Barley (winter)	2
Forage Grasses	2
Oats	2
Proso Millet	2
Rice	2
Rye (winter)	2
Seed corn	2
Sorghums	2
Spring cereal crops	2
Wheat (winter)	2
Popcorn, Sweetcorn*	3
Cotton	4
Peanuts	6
Tomato (transplant)	8
Alfalfa	9
Clovers	9
Dry Beans	9
Field Peas	9
Peas	9
Potatoes	9
Cucumbers, Pumpkins, Squash	9
Snap Beans	9
Soybeans	9
Peppers	10
Eggplant	12
Radish	12
Cabbage	15
Canola	15
Carrot	15
Mint	15
Broccoli, Cauliflower, Collards	18
Leeks, Onions	18
Lettuce crops	18
Sunflowers	18
Sugarbeet (Michigan only)	21
Sugarbeet and Red Beet	24
Spinach	24
Sugarbeet (ND, MN, Red River Valley)**	36

\* In-crop and preplant applications of GWN-3061 to sweetcorn and popcorn are based on application rates and timings specific for use in those crops. Rotational interval must be adhered to for planting subsequent sweet

corn or popcorn crops after GWN-3061 applications in sweetcorn or popcorn crops that are lost, terminated, or harvested.

\*\* Also includes other regions where rainfall is sparse or irrigation is required.

**Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.**

#### LIST OF COMMON AND SCIENTIFIC NAMES OF BROADLEAF WEEDS

Bindweed, hedge <i>Calystegia sepium</i>	Morningglory, tall <i>Ipomoea purpurea</i>
Burcucumber <i>Sicyas angulatus</i>	Mustard, wild <i>Sinapis arvensis</i>
Cocklebur, common <i>Xanthium strumarium</i>	Nutsedge, yellow <i>Cyperus exculentus</i>
Dogbane, hemp <i>Apocynum cannabinum</i>	Nutsedge, purple <i>Cyperus rotundus</i>
Fleabane, Philadelphia <i>Erigeron philadelphicus</i>	Passionflower, maypop <i>Passiflora incarnata</i>
Horsenettle <i>Solanum carolinense</i>	Pigweed, redroot <i>Amaranthus retrofractus</i>
Kyllinga, green <i>Kyllinga brevifolia</i>	Pokeweed, common <i>Phytolacca Americana</i>
Kochia <i>Kochia scoparia</i>	Radish, wild <i>Raphanus raphanistrum</i>
Jimsonweed <i>Datura stramonium</i>	Ragweed, common <i>Ambrosia artemisiifolia</i>
Lambsquarters, common <i>Chenopodium album</i>	Ragweed, giant <i>Ambrosia trifida</i>
Mallow, Venice <i>Hibiscus trionum</i>	Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>
Milkweed, common <i>Asclepias syriaca</i>	Sunflower, common <i>Helianthus annuus</i>
Milkweed, honeyvine <i>Ampelamus albidus</i>	Velvetleaf <i>Abutilon theophrasti</i>
Morningglory <i>Ipomoea spp.</i>	Waterhemp, common <i>Amaranthus rudis</i>
Morningglory, ivyleaf <i>Ipomoea hederacea</i>	Waterhemp, tall <i>Amaranthus tuberculatus</i>

#### FIELD CORN AND FIELD CORN GROWN FOR SEED

Corn Growth Stage: When used alone, GWN-3061 can be applied over-the-top or with drop nozzles from the spike through layby stage of field corn.

GWN-3061 may be applied up to 2 applications with a total application not to exceed 2 2/3 ounces of product by weight (0.125 pound active ingredient) per acre per use season.

Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

8/23

**WEEDS CONTROLLED**

**GWN-3061**

**CORN USE RATE GUIDE**

Use Rate – 2/3 ounce of product by weight per acre  
(0.031 pound active ingredient per acre)

Weed Species	Size Range
	Height (inches)
Cocklebur, common	1 to 9
Fleabane, Philadelphia	1 to 3
Kochia	1 to 3
Mallow, Venice	1 to 3
Passionflower, maypop	1 to 3
Pigweed, redroot	1 to 3
Pokeweed, common	1 to 6
Ragweed, common	1 to 9
giant	1 to 3
Smartweed, Pennsylvania	1 to 2
Sunflower, common	1 to 12
Velvetleaf	1 to 9

Use Rate – 1 to 1 1/3 ounces of product by weight per acre  
(0.047 to 0.062 pound active ingredient per acre)

Weed Species	Size Range
	Height (inches)
Cocklebur, common	9 to 14
Mallow, Venice	4 to 12
Milkweed, honeyvine	1 to 6
Mustard, wild	4 to 6
Nutsedge, yellow <sup>1</sup>	4 to 12
purple	4 to 12
Pigweed, redroot <sup>2</sup>	4 to 6
Radish, wild	4 to 6
Ragweed: common	9 to 12
giant	4 to 6
Sunflower, common	12 to 15
Velvetleaf <sup>2</sup>	9 to 12

<sup>1</sup> Heavy infestations of nutsedge may require sequential applications. An earlier treatment may be required to prevent nutsedge from competing with the crop.

<sup>2</sup> For large velvetleaf and pigweed, the addition of liquid nitrogen fertilizer (2 to 4 quarts per acre) plus crop oil concentrate or nonionic surfactant is recommended.

**WEEDS SUPPRESSED**

Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

Weed Species	ounces by weight per acre	
	2/3 ounce Height (in.)	1 to 1 1/3 ounce Height (in.)
Burcucumber	1 to 3	4 to 12
Kochia	*	3 to 6
Lambsquarters, common	1 to 2	-----
Milkweed, common	3 to 5	6 to 12
Milkweed, honeyvine	1 to 3	-----
Morningglory	-----	1 to 3
Nutsedge: yellow	4 to 12	*
purple	4 to 12	*

\* Refer to "WEEDS CONTROLLED" section of this label.

Refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.



9/23

**Tank Mixtures for Corn Only**

Ensure that spray equipment is set up to avoid applying an excessive rate directly over the rows and into the whorl of the cornstalk. To insure good spray coverage of weeds and to reduce the risk of spraying directly into the whorl, tank-mix applications made after corn is 24 inches tall should be directed or semi-directed using drop nozzles.

**GWN-3061 Tank-Mixture Options in Field Corn & Seed Corn**

Tank Mix Partners	Rate per Acre	Additives	Application Method	Comments
Banvel or Clarity™	2 to 8 fl. ounces	NIS	<ul style="list-style-type: none"> <li>Broadcast up to 36" tall corn.</li> <li>Use lower Banvel rates or directed sprays on corn taller than 8".</li> </ul>	<ul style="list-style-type: none"> <li>COC may cause crop injury, especially with higher Banvel/Clarity rates.</li> <li>For large corn, avoid direct spraying into whorl of cornstalk.</li> </ul>
Marksman™	1/2 to 2 pints	NIS	<ul style="list-style-type: none"> <li>Broadcast up to 8" tall corn.</li> </ul>	<ul style="list-style-type: none"> <li>COC may cause crop injury.</li> </ul>
2,4-D (4 pounds/gal)	4 to 8 fl. ounces	NIS	<ul style="list-style-type: none"> <li>Broadcast up to 8" tall corn.</li> </ul>	<ul style="list-style-type: none"> <li>If corn exceeds 8" directed sprays with drop nozzles are required.</li> </ul>
Buctril™	1/2 to 1 pint	NIS	<ul style="list-style-type: none"> <li>Broadcast to corn up to tassel emergence.</li> </ul>	<ul style="list-style-type: none"> <li>Leaf burn may occur.</li> <li>COC or 28 percent may cause additional leaf burn</li> </ul>
BUCTRIL+ atrazine	1 to 2 pints	NIS	<ul style="list-style-type: none"> <li>Broadcast to corn up to 12" tall.</li> </ul>	<ul style="list-style-type: none"> <li>Leaf burn may occur.</li> <li>COC or 28 percent may cause additional leaf burn</li> </ul>
Atrazine 4L	1 1/2 to 3 pints	COC	<ul style="list-style-type: none"> <li>Broadcast to corn up to 12" tall.</li> </ul>	<ul style="list-style-type: none"> <li>Control is best when weeds are small.</li> <li>Effective for burndown of grass weed escapes.</li> <li>Antagonism may occur on larger broadleaf weeds.</li> </ul>
Accent	2/3 ounce	COC or NIS	<ul style="list-style-type: none"> <li>Broadcast or apply with drop nozzles to corn up to 24" tall.</li> <li>For corn 24" to 36" tall, apply with drop nozzles only.</li> </ul>	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer (e.g., 28 percent) is also recommended as an additive.</li> <li>Avoid spraying directly into whorls of larger cornstalks.</li> <li>Refer to Accent label for soil insecticide interaction information.</li> </ul>
Beacon	0.76 ounce (1/2 packet)	COC or NIS	<ul style="list-style-type: none"> <li>Broadcast or apply with drop nozzles to corn up to 20" tall.</li> <li>For corn 20" to pre-tassel, apply with drop nozzles only.</li> </ul>	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer (e.g., 28 percent) is also recommended as an additive.</li> <li>Avoid spraying directly into whorls of larger corn.</li> <li>Refer to Beacon label for soil insecticide interaction restrictions.</li> <li>Consult your dealer, seed supplier, or Syngenta representative for a list of susceptible hybrids.</li> </ul>
Accent Gold	2.9 ounces	COC	<ul style="list-style-type: none"> <li>Broadcast to corn up to 12" tall.</li> </ul>	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer (e.g. 28 percent) is also recommended as an additive.</li> <li>Do not apply to seed corn.</li> <li>Refer to Accent Gold label for soil insecticide interactions.</li> </ul>
Basis Gold	14 ounces	COC or NIS	<ul style="list-style-type: none"> <li>Broadcast to corn up to 12" tall.</li> </ul>	<ul style="list-style-type: none"> <li>Ammonium nitrogen fertilizer (e.g. 28 percent) is also recommended as an additive.</li> <li>Do not apply to seed corn.</li> <li>Refer to Accent Gold label for soil insecticide.</li> </ul>

NIS = Nonionic surfactant. COC = Crop oil concentrate.

Refer to "MIXING INSTRUCTIONS," "TANK MIXTURES" and "USE RATE GUIDES" sections of this label for detailed information.

**Refer to the specific product labels and observe all precautions, mixing and application instructions for all products used in tank mixtures.**

10/23

**GWN-3061 plus ACCENT™**

A tank mixture of GWN-3061 plus Accent may be used for the post-emergence control of annual broadleaf weeds and annual grasses in corn only. GWN-3061 plus Accent may be applied over-the-top or with drop nozzles to field corn up to 24 inches tall (free standing). For corn 24 to 36 inches tall, refer to the Accent label for application restrictions. Banvel, Marksman, Clarity, Buctril or BUCTRIL+atrazine may also be added to the tank mixtures for improved control of certain weed species.

**Refer to the Accent label for use instructions and restrictions on corn varieties and insecticides.**

**GWN-3061 plus BEACON™**

A tank mixture of GWN-3061 plus Beacon may be used for the post-emergence control of annual broadleaf weeds and annual grasses in corn only. GWN-3061 plus Beacon may be applied over-the-top or directed to field corn when corn height is between 4 and 20 inches tall. Drop nozzles are required with the Beacon mixture when corn is between 20 inches tall and tassel emergence. Banvel, Marksman, Clarity, Buctril or BUCTRIL+atrazine may also be added to the tank mixtures for improved control of certain weed species.

**Refer to the Beacon label for use instructions and restrictions on corn varieties and insecticides.**

**Additional grass species controlled by tank mixing with Accent or Beacon.**

Grasses	Size Range Height (inches)	
	GWN-3061 + Accent (2/3-1oz)+(2/3 oz)	GWN-3061 + Beacon (2/3-1oz)+(1/2 packet)
Barnyardgrass	2 to 4	
Cupgrass, wooly	2 to 4	
Foxtails: giant, yellow	2 to 4	
green, bristly	2 to 4	
Itchgrass	2 to 6	
Johnsongrass, rhizome	8 to 18	8 to 16
Seedling	4 to 12	4 to 12
Millet, wild proso	1 to 4	
Oats, wild	2 to 4	
Panicum, browntop	1 to 3	
Panicum, fall	2 to 4	Less than 2
Panicum, Texas	1 to 3	
Quackgrass	4 to 10	4 to 8
Ryegrass, Italian	2 to 6	
Sandbur	1 to 3	
Shattercane	4 to 12	4 to 12
Signalgrass, broadleaf	1 to 2	
Sorghum-almum	4 to 12	4 to 12

**GWN-3061 plus SOIL RESIDUALS**

Micro-Tech® or Bullet or Harness® Xtra or Harness® Xtra 5.6L or Degree™ or Degree Xtra™ plus GWN-3061 may be applied early post-emergence for control of additional broadleaf weeds and nutsedge in field corn (including seed corn).

These tank mixtures will provide post-emergence control of small emerged grasses and broadleaf weeds as well as residual pre-emergence control or reduced competition of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" SECTION OF THE Micro-Tech, Bullet, Harness, Harness Xtra, Harness Xtra 5.6L, Degree, Degree Xtra herbicide labels.

Apply these tank-mixtures to emerged grasses at the 2-leaf stage or less and to corn less than 11 inches tall (5 inch corn for Micro-Tech and Bullet). Include 28 percent nitrogen fertilizer at a rate of 4 gallons per 100 gallons of spray solution plus NIS at 1 quart per 100 gallons of spray solution in 15 to 30 gallons of water per acre. The addition of Banvel or Clarity at 2 ounces of product per acre is recommended to these mixtures to control emerged lambsquarters less than 4 inches tall.

**RECOMMENDED RATES/ACRE:**

Soil Residual	Plus	GWN-3061
(quarts)		(ounces)
Labeled rate		2/3

11/23

### GWN-3061 plus ACCENT plus SOIL RESIDUALS

Micro-Tech® or Bullet or Harness® or Harness® Xtra or Harness® Xtra 5.6L or Degree™ or Degree Xtra™ plus GWN-3061 plus Accent may be applied early post-emergence for control of additional broadleaf weeds and nutsedge in field corn (including seed corn).

These tank mixtures will provide post-emergence control of emerged foxtails as well as residual pre-emergence control or reduced competition of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of the Micro-Tech, Bullet, Harness, Harness Xtra, Harness Xtra 5.6L, Degree, Degree Xtra herbicide labels.

Apply these tank-mixtures to emerged foxtails less than 2 inches tall and to corn less than 11 inches tall (5 inch corn for Micro-Tech and Bullet). Include 28 percent nitrogen fertilizer at a rate of 4 gallons per 100 gallons of spray solution plus NIS at 1 quart per 100 gallons of spray solution in 15 to 30 gallons of water per acre. The addition of Banvel or Clarity at 2 ounces of product per acre is recommended to these mixtures to control emerged lambsquarters less than 4 inches tall.

#### RECOMMENDED RATES/ACRE:

Soil Residual	<u>Plus</u>	GWN-3061	<u>Plus</u>	Accent
<u>(Quarts)</u>		<u>(Ounces)</u>		<u>(Ounces)</u>
Labeled rate		2/3		1/3 to 1/2

### GWN-3061 plus GLYPHOSATE AGRICULTURAL HERBICIDES plus NONIONIC SURFACTANT

GWN-3061 may be applied at 2/3 ounce by weight per acre in combination with Glyphosate herbicides labeled for agricultural uses for pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge with Pioneer IR corn hybrids only. Pioneer IR hybrids are required to ensure crop safety due to the pre-plant application. Banvel or 2,4-D may also be applied in this tank mixture for enhanced pre-plant burndown of broadleaf weeds.

### GWN-3061 SOIL APPLICATIONS

When used exclusively with Pioneer IR field corn hybrids, GWN-3061 may be soil applied at the rate of 1 1/3 to 2 ounces by weight per acre (0.062 to 0.094 pound of active ingredient per acre) for residual control of velvetleaf, common cocklebur, common lambsquarters, common ragweed, pigweed, smartweed, sunflower and other difficult to control weeds.

This product is recommended as an early pre-plant surface-applied, pre-plant incorporated, or pre-emergence treatment. GWN-3061 offers effective broadleaf control across all tillage systems and is intended for use in tank mixtures with pre-emergence grass herbicides, including but not limited to: Harness, Harness Xtra, Harness Xtra 5.6L, Degree, Degree Xtra, Micro-Tech, Bullet, Lariat and Lasso.

Refer to the labels for these products, or any other grass pre-emergence herbicide used for use instructions, weeds controlled, and application restrictions.

### GWN-3061 plus BANVEL plus NONIONIC SURFACTANT

For the control of additional broadleaf weeds, GWN-3061 may be applied in tank mixtures with Banvel. A GWN-3061 tank mixture with low rates of Banvel may be applied during the period beginning at corn emergence and continuing until corn is 36 inches in height. Applications should not be made after corn exceeds 36 inches or 15 days before tassel emergence, whichever comes first. Clarity or Marksman may be substituted in this tank mixture.

Refer to the labels for Banvel, Clarity, and Marksman products for label restrictions.

### GWN-3061 plus 2,4-D plus NONIONIC SURFACTANT

For the control of additional broadleaf weeds, GWN-3061 may be applied in tank mixtures with 2,4-D. Avoid spraying just after corn leaves unfold, as injury may occur. A GWN-3061 tank mixture with 2,4-D may be applied during the period from corn emergence through the 5 leaf stage or 8 inches tall, whichever comes first. If corn exceeds 8 inches, directed spray applications with drop nozzles must be used for tank mixtures with 2,4-D.

Refer to the labels for 2,4-D products for label restrictions.

12/23

**GWN-3061  
USE RATE GUIDE**

**GWN-3061 plus BANVEL plus NONIONIC SURFACTANT  
GWN-3061 plus 2,4-D plus NONIONIC SURFACTANT**

GWN-3061 Use Rate – 2/3 ounce of product by weight per acre  
 Banvel Use Rate – 1/4 to 1/2 pint per acre  
 2,4-D Use Rate – 1/4 to 1/2 pint per acre  
 (0.125 to 0.25 pound active ingredient per acre)

<b>Weed Species</b>	<b>Size Range Height (inches)</b>
Bindweed	1 to 6
Burcucumber <sup>1</sup>	4 to 2
Cocklebur, common	1 to 12
Dogbane, hemp <sup>3</sup>	1 to 6
Horsenettle	1 to 8
Jimsonweed	1 to 4
Kochia <sup>1</sup>	1 to 6
Lambsquarters, common <sup>2</sup>	1 to 6
Mallow, Venice	1 to 3
Milkweed, common	1 to 6
Morningglory, ivyleaf	1 to 6
Morningglory, tall	1 to 6
Nightshade, black <sup>1</sup>	1 to 6
Pigweed, redroot	1 to 12
Pokeweed, common <sup>1</sup>	1 to 18
Ragweed: common	1 to 12
giant <sup>1</sup>	1 to 6
Smartweed, Pennsylvania	1 to 3
Sunflower, common	1 to 12
Thistle, Canada <sup>1</sup>	1 to 6
Velvetleaf	1 to 12

<sup>1</sup> Banvel tank mixture only.  
<sup>2</sup> A tank mixture of GWN-3061 (2/3 ounce by weight per acre) plus Banvel at 2 fluid ounces per acre (1/8 pint) is recommended for the control of common lambsquarters less than 4 inches in height.  
<sup>3</sup> Provides suppression of this weed.

**GWN-3061 plus BUCTRIL plus NONIONIC SURFACTANT**

GWN-3061 may be applied in combination with Buctril or BUCTRIL + atrazine herbicides for post-emergence control of many annual broadleaf weeds in corn. Use 2/3 ounce of GWN-3061 by weight plus surfactant in combination with 1/2 to 1 pint of Buctril and 1 to 2 1/2 pints of BUCTRIL + atrazine herbicide.

**Refer to Buctril and Buctril + atrazine labels for use instructions, weeds controlled and application restrictions.**

**GWN-3061 plus ATRAZINE**

GWN-3061 may be applied in combination with atrazine for post-emergence control of labeled broadleaf weeds. The addition of atrazine will also aid in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Applications should be made when broadleaf weeds are small (3 inches or less).

Mixtures with atrazine may result in reduced control (antagonism) of larger broadleaf weeds. Use the labeled rate for GWN-3061 plus atrazine 4L at 1 1/2 to 3 pints per acre (0.75 to 1 1/2 pounds active ingredient per acre). The addition of crop oil concentrate (COC) is recommended for this mixture.

**Refer to the atrazine 4L label for use instructions, additive requirements, weeds controlled and application restrictions.**

13/23

**SWEET CORN AND POPCORN**

Corn Growth Stage: When used alone, this product may be applied over-the-top or with drop nozzles from the spike through layby stage of the corn.

Apply 2/3 ounce by weight (0.031 pound active ingredient) of this product per acre broadcast over the top or with drop nozzles in sweetcorn and popcorn. Mechanical cultivation may be required to control weeds species not on the label. Avoid cultivation for at least 7 days following application. If necessary, a sequential treatment of this product at 2/3 ounce by weight per acre may be applied only with drop nozzles semi-directed or directed to avoid application into the corn plant whorl. No more than 2 applications of this product may be made per year in sweetcorn and popcorn. (Any single application must not exceed 2/3 ounce by weight per acre).

Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

This product may be applied to sweetcorn and popcorn, however, the user assumes responsibility for such use. All hybrids/varieties have not been tested for sensitivity to GWN-3061 nor does Canyon have access to all seed company or processor data. Consequently, any injury arising from the use of this product on sweetcorn and popcorn is the responsibility of the user. Do not apply this product to sweetcorn or popcorn unless the seed company, processor or State Agricultural Extension service has tested this product on the particular hybrid/variety and specifically approves and recommends the use. Do not apply this product to sweetcorn or popcorn if the crop is under severe stress due to drought, water-saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions. Refer to the following "WEEDS CONTROLLED" section for use rate recommendations. Also refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.

This product is not recommended for use on 'Jubilee' sweetcorn.

Canyon does not recommend application of this product to sweetcorn or popcorn previously treated with soil applied organophosphate insecticides. Do not apply an organophosphate insecticide within 7 days before or 3 days after any application of this product.

**WEEDS CONTROLLED  
SWEET CORN AND POPCORN  
USE RATE GUIDE**

Use Rate – 2/3 ounce of product by weight per acre  
(0.031 pound active ingredient per acre)

<b>Weed Species</b>	<b>Size Range Height (inches)</b>
Cocklebur, common	1 to 9
Fleabane, Philadelphia	1 to 3
Kochia	1 to 3
Mallow, Venice	1 to 3
Passionflower, maypop	1 to 3
Pigweed, redroot	1 to 3
Pokeweed, common	1 to 6
Ragweed: common	1 to 9
giant	1 to 3
Smartweed, Pennsylvania	1 to 2
Sunflower, common	1 to 12
Velvetleaf	1 to 9

**WEEDS SUPPRESSED**

Use Rate – 2/3 to ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

<b>Weed Species</b>	<b>ounces by weight per acre</b>
	<b>2/3 ounce Height (in.)</b>
Burcumber	1 to 3
Kochia	*
Lambsquarters, common	1 to 2
Milkweed, common	3 to 5
Milkweed, honeyvine	1 to 3
Morningglory	----
Nutsedge, yellow	4 to 12
purple	4 to 12

4/23

**GRAIN SORGHUM (MILO)**

Grain Sorghum Growth Stage: GWN-3061 alone, can be applied from the 2-leaf through layby stage (before grain head emergence).

Only apply GWN-3061 in a single application with the total application rate not to exceed 1.0 ounce of product by weight (0.047 pound active ingredient) per acre per use season.

Temporary stature reduction may occur to the crop following application of GWN-3061 if the grain sorghum is under stress. This effect will be most evident 7 to 10 days after application. The crop will quickly recover under normal growing conditions.

Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

**WEEDS CONTROLLED**

**GWN-3061**

**SORGHUM USE RATE GUIDE**

Use rate – 2/3 ounce of product by weight per acre  
(0.031 pound active ingredient per acre)

<b>Weed Species</b>	<b>Size Range Height (inches)</b>
Cocklebur, common	1 to 9
Fleabane, Philadelphia	1 to 3
Kochia	1 to 3
Mallow, Venice	1 to 3
Passionflower, maypop	1 to 3
Pigweed, redroot	1 to 3
Pokeweed, common	1 to 6
Ragweed: common	1 to 9
giant	1 to 3
Smartweed, Pennsylvania	1 to 2
Sunflower, common	1 to 12
Velvetleaf	1 to 9

Use Rate – 1.0 ounce of product by weight per acre  
(0.047 pound active ingredient per acre)

<b>Weed Species</b>	<b>Size Range Height (inches)</b>
Nutsedge: yellow	4 to 12
purple	4 to 12

**WEEDS SUPPRESSED**

Use rate – 2/3 ounce of product by weight per acre  
(0.031 pound active ingredient per acre)

<b>Weed Species</b>	<b>Size Range Height (inches)</b>
Burcucumber	1 to 3
Lambsquarters, common	1 to 2
Milkweed, common	3 to 5
Milkweed, honeyvine	1 to 3
Nutsedge: yellow	4 to 12
purple	4 to 12

Refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.

**Tank Mixtures for Grain Sorghum**

**GWN-3061 plus 2,4-D plus NONIONIC SURFACTANT**

A GWN-3061 tank mixture with 2,4-D may be applied to grain sorghum when the crop is 6 to 15 inches tall. If sorghum exceeds 8 inches, use drop nozzles and keep the spray off foliage. Do not treat during the boot, flowering or dough stage.

Applications should not be made when grain sorghum exceeds 15 inches. Do not treat grain sorghum during the boot, flowering, or dough stage. Clarity or Marksman may be substituted in this tank mixture.

**Refer to the labels for 2,4-D, Clarity and Marksman products for label restrictions.**

**GWN-3061  
USE RATE GUIDE**

**GWN-3061 plus 2,4-D plus NONIONIC SURFACTANT**

GWN-3061 Use Rate – 2/3 ounce of product by weight per acre  
2,4-D Use Rate – 1/4 to 1/2 pint per acre  
– (0.125 to 0.25 pound active ingredient per acre)

Weed Species	Size Range Height (inches)
Bindweed	1 to 6
Burcucumber <sup>1</sup>	4 to 12
Cocklebur, common	1 to 12
Dogbane, hemp <sup>3</sup>	1 to 6
Horsenettle	1 to 8
Jimsonweed	1 to 4
Kochia <sup>1</sup>	1 to 6
Lambsquarters, common <sup>2</sup>	1 to 6
Mallow, Venice	1 to 3
Milkweed, common	1 to 6
Morningglory, ivyleaf	1 to 6
Morningglory, tall	1 to 6
Nightshade, black <sup>1</sup>	1 to 6
Pigweed, redroot	1 to 12
Pokeweed, common <sup>1</sup>	1 to 18
Ragweed: common	1 to 12
giant <sup>1</sup>	1 to 6
Smartweed, Pennsylvania	1 to 3
Sunflower, common	1 to 12
Thistle, Canada <sup>1</sup>	1 to 6
Velvetleaf	1 to 12

<sup>1</sup> Banvel tank mixture only.

<sup>2</sup> A tank mixture of GWN-3061 (2/3 ounce by weight per acre) plus Banvel at 2 fluid ounces per acre (1/8 pint per acre) is recommended for the control of common lambsquarters less than 4 inches in height.

<sup>3</sup> Provides suppression of this weed.

**GWN-3061 plus BUCTRIL plus NONIONIC SURFACTANT**

GWN-3061 may be applied in combination with Buctril or BUCTRIL + atrazine herbicides for post-emergence control of many annual broadleaf weeds in grain sorghum. Use 2/3 ounce of GWN-3061 by weight plus surfactant in combination with 1/2 to 1 pint of Buctril and 1 to 2 1/2 pints of BUCTRIL + atrazine herbicide.

**Refer to Buctril and BUCTRIL + atrazine labels for use instructions, weeds controlled and application restrictions.**

**GWN-3061 plus ATRAZINE**

GWN-3061 may be applied in combination with atrazine for post-emergence control of labeled broadleaf weeds. The addition of atrazine will also aid in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Applications should be made when broadleaf weeds are small (3 inches or less).

Mixtures with atrazine may result in reduced control (antagonism) of larger broadleaf weeds. Use the labeled rate for GWN-3061 plus atrazine 4L at 1 1/2 to 3 pints per acre (0.75 to 1 1/2 pounds active ingredient per acre). The addition of crop oil concentrate (COC) is recommended for this mixture.

**Refer to the atrazine 4L label for use instructions, additive requirements, weeds controlled and application restrictions.**





17/23

## RICE PRE-EMERGENCE AND POST-EMERGENCE APPLICATIONS TO RICE

GWN-3061, when applied alone, may be applied for post-emergent weed control from prior to the emergence of rice until after permanent flood is established. GWN-3061 may be applied at 2/3 to 1 1/3 ounce by weight per acre, with the total application rate not to exceed 1 1/3 ounce of product by weight (0.062 lb. active ingredient) per acre per use season.

GWN-3061 can be applied as a foliar spray or dry broadcast.

GWN-3061 may be applied at 2/3 ounce by weight per acre in combination with Glyphosate agricultural herbicides for pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge. If this product is applied pre-plant burndown, refer to "TIME INTERVAL BEFORE PLANTING" table in complete Directions for Use.

This product may be tank-mixed with propanil containing rice herbicides (e.g. Stam M4 and Propanil 4E) at 2/3 to 1 1/3 ounce per acre of this herbicide and labeled rates of the tankmix products.

Foliar applications of GWN-3061 may be made at the 3-5 leaf stage of rice when weeds have 2-4 leaves. Dry broadcast applications may be made at the 1-2 leaf stage of rice when weeds have two leaves or less.

This product may also be applied post flood with dry broadcast applications of GWN-3061 herbicide at 1 to 1 1/3 ounce by weight per acre, with the total application rate not to exceed 1 1/3 ounce product by weight per acre per use season.

It is best to use 0.25 to 0.5 percent nonionic surfactant which contains at least 80% active ingredient with foliar applications of GWN-3061.

With all foliar applications of GWN-3061 use a minimum 3-15 gallons of water per acre for aerial equipment and a minimum of 10 gallons of water per acre for ground equipment. It is best to apply spray solutions the day they are mixed. **Note:** See "APPLICATION EQUIPMENT AND INSTRUCTION" section for spray drift management techniques.

Water levels in rice fields and checks should remain static (3 inch to 6 inch depth) following dry broadcast applications of GWN-3061. Do not reintroduce water into rice fields or checks for at least five days following dry broadcast applications of GWN-3061. Rice fields and checks may be irrigated to maintain water level, but this may reduce weed control.

Control of emerged weeds with foliar applications is best when 70% - 80% of the weed foliage is exposed. Control of submerged weeds is best when weeds have 2 leaves or less. Do not reintroduce water into rice fields or checks for at least 24 hours following foliar applications of GWN-3061.

Do not apply within 48 days of harvest. Do not apply within 69 days of harvest in California.

**CAUTION:** To ensure product effectiveness avoid using GWN-3061 on rice fields which have a history of weed biotypes resistant to Londax.

### SEQUENTIAL APPLICATIONS

GWN-3061 herbicide may be applied sequentially with Ordram, Bolero, Clincher, Regiment and Shark. Read the Ordram, Bolero, Clincher, Regiment and Shark labels for application information, restrictions and precautions.

18/23

**WEEDS CONTROLLED BY  
GWN-3061**

**RICE USE RATE**

Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

Weed Species	ounces by weight per acre	
	2/3 ounce Height (inches)	1 to 1 1/3 oz. Height (inches)
Cocklebur, common	1 to 9	9 to 14
Dayflower	1 to 2	3 to 4
Eclipta	1 to 4	4 to 8
Flatsedge rice	1 to 9	9 to 12
Fleabane, Philadelphia	1 to 3	-----
Jointvetch	1 to 2	3 to 4
Kochia	1 to 3	-----
Mallow, Venice	1 to 3	4 to 12
Milkweed, honeyvine	-----	1 to 6
Mustard, wild	-----	4 to 6
Nutsedge <sup>1</sup> :		
yellow	1 to 6	6 to 12
purple	1 to 6	6 to 12
Passionflower, maypop	1 to 3	-----
Pigweed, redroot <sup>2</sup>	1 to 3	4 to 6
Pokeweed, common	1 to 6	-----
Radish, Wild	-----	4 to 6
Ragweed:		
common	1 to 9	9 to 12
giant	1 to 3	4 to 6
Sesbania, Hemp	1 to 3	3 to 6
Sida, Prickly	1 to 2	3 to 4
Smartweed, Pennsylvania	1 to 2	-----
Sunflower, common	1 to 12	12 to 15
Velvetleaf <sup>2</sup>	1 to 9	9 to 12

<sup>1</sup> Heavy infestations of nutsedge may require sequential applications. An earlier treatment may be required to prevent nutsedge from competing with the crop.

<sup>2</sup> For large velvetleaf and pigweed, the addition of liquid nitrogen fertilizer (2 to 4 quarts per acre) plus crop oil concentrate or nonionic surfactant is recommended.

**WEEDS CONTROLLED**

Use Rate – 1 to 1 1/3 ounces of product by weight per acre  
(0.047 to 0.062 pound active ingredient per acre)

- |                   |                           |
|-------------------|---------------------------|
| Ricefield Bulrush | Smallflower Umbrellaplant |
| Redstem           | California Arrowhead      |

**WEEDS SUPPRESSED**

Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

Weed Species	ounces by weight per acre	
	2/3 ounce Height (inches)	1 to 1 1/3 oz. Height (inches)
Burcucumber	1 to 3	4 to 12
Kochia	*	3 to 6
Lambsquarters, common	1 to 2	-----
Milkweed, common	3 to 5	6 to 12
Milkweed, honeyvine	1 to 3	-----
Morningglory	-----	1 to 3
Nutsedge:		
Yellow	4 to 12	*
Purple	4 to 12	*

\* Refer to "WEEDS CONTROLLED" section of the label booklet.

19/23

**SUGARCANE**

When used alone, this product may be applied prior to planting, prior to emergence or after the emergence of the sugarcane, and until row closure. Apply 2/3 to 1 1/3 ounces by weight (0.031 to 0.062 pound active ingredient) of this product per acre. Mechanical cultivation may be required to control weed species not on the label. If so, a **sequential treatment** may be required to control weeds in areas disturbed soil. No more than 3 applications (including pre-plant applications) may be made with the total use rate not to exceed 2 2/3 ounces of product by weight (0.125 pound active ingredient) per acre per year.

Following application to foliage allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

This product may be applied at 2/3 to 1 1/3 ounces by weight per acre (0.031 to 0.062 pound active ingredient per acre) in combination with Glyphosate agricultural herbicides for pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge in sugarcane.

**WEEDS CONTROLLED  
SUGARCANE**

Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

Weed Species	ounces by weight per acre	
	2/3 ounce Height (inches)	1 to 1 1/3 oz. Height (inches)
Cocklebur, common	1 to 9	9 to 14
Fleabane, Philadelphia	1 to 3	-----
Kochia	1 to 3	-----
Mallow, Venice	1 to 3	4 to 12
Milkweed, honeyvine	-----	1 to 6
Mustard, wild	-----	4 to 6
Nutsedge <sup>1</sup> :		
yellow	-----	4 to 12
purple	-----	4 to 12
Passionflower, maypop	1 to 3	-----
Pigweed, redroot <sup>2</sup>	1 to 3	4 to 6
Pokeweed, common	1 to 6	-----
Radish, Wild	-----	4 to 6
Ragweed:		
common	1 to 9	9 to 12
giant	1 to 3	4 to 6
Smartweed, Pennsylvania	1 to 2	-----
Sunflower, common	1 to 12	12 to 15
Velvetleaf <sup>2</sup>	1 to 9	9 to 12

<sup>1</sup> Heavy infestations of nutsedge may require sequential applications. An earlier treatment may be required to prevent nutsedge from competing with the crop.

<sup>2</sup> For large velvetleaf and pigweed, the addition of liquid nitrogen fertilizer (2 to 4 quarts per acre) plus crop oil concentrate or nonionic surfactant is recommended.

**WEEDS SUPPRESSED**

Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre  
(0.031 to 0.062 pound active ingredient per acre)

Weed Species	ounces by weight per acre	
	2/3 ounce Height (inches)	1 to 1 1/3 oz. Height (inches)
Burcucumber	1 to 3	4 to 12
Kochia	*	3 to 6
Lambsquarters, common	1 to 2	-----
Milkweed, common	3 to 5	6 to 12
Milkweed, honeyvine	1 to 3	-----
Morningglory	-----	1 to 3
Nutsedge, yellow	4 to 12	*
Purple	4 to 12	*

\* Refer to "WEEDS CONTROLLED" section of this label.

20/23

**Tank Mixtures for Sugarcane**

GWN-3061 may be tank mixed with Asulox™, atrazine 4L, Evik™ or 2,4-D for application in sugarcane.

**GWN-3061 plus GLYPHOSATE AGRICULTURAL HERBICIDES plus NONIONIC SURFACTANT**

GWN-3061 may be applied at 2/3 to 1 1/3 ounces by weight per acre (0.031 to 0.062 pound ai/acre) in combination with recommended rates of Glyphosate agricultural herbicides for pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge in sugarcane.

Refer to the Glyphosate agricultural herbicide label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated, and application restrictions.

**GWN-3061 plus ASULOX plus NONIONIC SURFACTANT or CROP OIL CONCENTRATE**

GWN-3061 may be applied in tank mixtures with Asulox for the control of labeled grasses. A GWN-3061 tank mixture with Asulox may be applied to sugarcane before crop emergence or post-emergence until 90 days before harvest. Up to 2 applications per year may be made in accordance with label recommendations.

GWN-3061 Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre. Up to 3 treatments per year may be applied, not exceeding 2 2/3 ounces of product by weight per acre per year.

Asulox Use Rate – 6 to 8 pints of product per acre. Up to 2 treatments per year may be applied.

Refer to the Asulox label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated, and application restrictions.

**GWN-3061 plus ATRAZINE 4L plus NONIONIC SURFACTANT or CROP OIL CONCENTRATE**

GWN-3061 may be applied in combination with atrazine 4L for post-emergence control of labeled broadleaf weeds in sugarcane. The addition of atrazine will also aid in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Applications should be made when broadleaf weeds are small (3 inches or less). Mixtures with atrazine may result in reduced control (antagonism) of larger broadleaf weeds.

GWN-3061 Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre. Up to 3 treatments per year may be applied not exceeding 2 2/3 ounces of product by weight per acre per year.

Atrazine Use Rate – 4 to 8 pints per acre (1 to 2 pounds active ingredient). Follow the specific recommendations on the atrazine label for number and timing of applications and for maximum number of applications per year.

Refer to the atrazine 4L label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated and application restrictions.

**GWN-3061 plus EVIK plus NONIONIC SURFACTANT**

GWN-3061 may be applied in tank mixtures with Evik for the control of additional broadleaf weeds and grasses. A GWN-3061 tank mixture with Evik may be applied to sugarcane before crop emergence or post-emergence until row closure.

GWN-3061 Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre. Up to 3 treatments per year may be applied, not exceeding 2 2/3 ounces of product by weight.

Evik Use Rate – 1/2 to 1 1/2 pounds of product per acre. Follow the specific recommendations on the Evik label for number and timing of applications and for maximum number of applications per year.

Refer to the Evik label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated, and application restrictions.

**GWN-3061 plus 2,4-D AMINE plus NONIONIC SURFACTANT**

GWN-3061 may be applied in tank mixtures with 2,4-D amine for the control of additional broadleaf weeds. A GWN-3061 tank mixture with 2,4-D may be applied to sugarcane before crop emergence or post-emergence until 6 weeks before harvest.

GWN-3061 Use Rate – 2/3 to 1 1/3 ounces of product by weight per acre. Up to 3 treatments per year may be applied, not exceeding 2 2/3 ounces of product by weight.

2,4-D Use Rate – 2 to 4 pints per acre (1 to 2 pounds active ingredient per acre). Up to 4 treatments per year may be applied.

Refer to the 2,4-D amine label for use instructions, additive requirements, weeds controlled, the size range of weeds that should be treated, and application restrictions.

Refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.

### COTTON

GWN-3061 may be applied as a directed spray in hooded equipment for post-emergent weed control in emerged cotton. Applications may be made anytime after cotton emergence until row closure inhibits use of hooded spray equipment. The applicator is responsible for maintaining proper spray speed and equipment position so spray mist does not contact cotton plants.

GWN-3061 alone, may be applied at 2/3 to 1 1/3 ounce by weight per acre, with the total application rate not to exceed 1 1/3 ounce of product by weight (0.062 lb. active ingredient) per acre per use season. Contact of the herbicide solution with desirable vegetation may result in damage or destruction.

Do not apply within 28 days of harvest.

Also refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.

### FALLOW GROUND

Applications of GWN-3061 may be made to fallow ground at use rates ranging between 2/3 to 1 1/3 ounces of product by weight per acre. GWN-3061 may be applied up to 2 applications with a total application not to exceed 2 2/3 ounces of product by weight (0.125 pound active ingredient) per acre per use season.

Refer to the "FIELD CORN" section of this label for weed control recommendations. Also refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions.

### DRY BEANS

**GWN-3061 can be applied to direct seeded dry beans either pre-emergence, or as a row middle/furrow application.**

For pre-emergence applications, apply after planting but prior to soil cracking. Apply a rate of 1/2 - 2/3 ounces uniformly with ground equipment in a minimum of 15 gallons of water per acre. Use the lower rate on lighter textured soils with low organic matter.

GWN-3061 may be applied at a rate of 1/2 - 1 ounce between rows of crop for the control of nutsedge and listed broadleaf weeds. Avoid contact of the herbicide with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Reduce rate and spray volume in proportion to area actually sprayed.

Do not apply more than 1 ounce GWN-3061 per acre per crop-cycle, not to exceed 2 ounces per acre per 12-month period (includes applications to the crop and to Row Middles/Furrows).

### Tank Mixtures for Dry Beans

#### GWN-3061 and EPTAM 7E

A tank-mix combination of GWN-3061 plus EPTAM 7-E will give a broader spectrum of weed control than either product used separately. Read both labels carefully before using. Observe all cautions and limitations on labeling of both products.

Apply and incorporate 1/2 to 2/3 ounce GWN-3061 and 3-1/2 to 4-1/2 pints EPTAM 7-E per acre to a depth of approximately 2 inches just before planting. Use lower rate on lighter textured soils with low organic matter.

Refer to EPTAM 7-E label for specific incorporation directions. Rotary hoe lightly during or shortly after emergence of the beans to break any crust which occurs.

Do not apply more than 2/3 ounce GWN-3061 per acre per crop-cycle, not to exceed 2 ounces per acre per 12-month period (includes applications to the crop and to Row Middles/Furrows).

Do not use EPTAM 7-E on Adzuki beans, cowpeas (black-eyed peas, black-eyed beans), soybeans, lima beans, Mung beans, Garbanzo beans or other flat-podded beans except Romano. Under abnormal weather conditions, stunting may occur on Gratiot, Michilite, Sanilac, Seafarer, and Seaway varieties. Do not exceed 9 pints EPTAM 7-E per acre per crop.

Do not exceed 3-1/2 pints EPTAM 7-E per acre on small white beans or green beans grown on coarse textured soils.

Do not exceed 7 pints per acre per crop of Eptam in the Southwestern and Southeastern regions. Do not exceed 8 pints per acre per crop of Eptam in the Western Region. Do not exceed 9 pints per acre per crop of Eptam in the Pacific Northwestern Region. Do not exceed 9 3/4 pints of Eptam in the Northern Region.

*Rebecca A. Hargadine*  
Rebecca A. Hargadine

22/23

### STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store under cool, dry conditions (below 120° F). Do not store under moist conditions.

**Water Soluble Packaging [GWN-3061]**

Keep outer pouch **TIGHTLY** sealed to prevent moisture from damaging any unused water soluble bags.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal in accordance with applicable Federal, state or local procedures, or in such other method as is approved under those procedures.

**Water Soluble Packaging [GWN-3061]:** Emptied foil pouch and cardboard box retain vapor and product residue. Observe all labeled safeguards until containers are destroyed. Do not reuse foil pouch or box.

Completely use water soluble bags in application equipment. Then dispose of empty foil pouch and cardboard box in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. If burned stay out of smoke.

**Plastic Bottle Packaging [GWN-3061]:** Do not reuse container. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. If burned, stay out of smoke.

**DISPOSAL AUTHORITIES:** If none of the foregoing procedures is permitted by state and local authorities, then contact your State Pesticide or Environmental Control Agency, or your local Hazardous Waste Disposal office, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.**

### NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

**Important:** Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

**Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Canyon Group. All such risks shall be assumed by the Buyer and User.**

Canyon Group warrants that his product conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label when used in strict conformance with Directions for Use, subject to the above stated risk limitations. CANYON GROUP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**BUYER'S OR USER'S EXCLUSIVE REMEDY AND CANYON GROUP'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT CANYON GROUP'S SOLE DISCRETION.**

**THIS IS AN END-USE PRODUCT. AND CANYON DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKING.**

Bullet, Harness, Degree, Degree Xtra, Lasso, Micro-Tech, and Partner are registered trademarks of Monsanto Technology LLC.

Pioneer is a registered trademark of Pioneer Hi-Bred International, Inc.

Accent is a trademark of E.I. DuPont de Nemours & Co.

Banvel, Clarity, and Marksman are trademarks of BASF Corporation.

Beacon, Dual Magnum and Evik are trademarks of Novartis Finance Corporation.

Asulox and Buctil are registered trademarks of Aventis CropScience USA, Inc.

Eradicane is a registered trademark of Gowan Company.

Eptam is a registered trademark of Gowan Company

Sutan is a registered trademark of Syngenta Corporation.

Surpass is a registered trademark of Dow AgroSciences LLC.

Formulated in the United States using Active Ingredient made in Japan.

\*Manufactured by Nissan Chemical Industries, Ltd.

This product is protected by U.S. Patent No. 4,668,277

23/23

# Gowan®

*The Go To Company*

P.O. Box 5569 ▲ Yuma, AZ 85366-5569 ▲ Phone (928) 783-8844 ▲ FAX (928) 343-9255

**FEDERAL EXPRESS Acct. 3055-4440 3 Day**

August 30, 2005

Document Processing Desk (FPL)  
U.S.E.P.A. OPP (7504C)  
1801 South Bell Street  
Arlington, Virginia 22202-4501

**ATTN: Jim Tompkins, PM Team 25 (703) 305-5697  
Herbicide Branch, Registration Division (7505C)**

**RE: GWN-3061, EPA Reg. No. 81880-2  
Notification of Minor Label Revision**

Dear Mr. Tompkins:


In accordance with PR-Notice 98-10, we have made minor label revisions to the referenced label including correction of typographical errors and rewording for clarification purposes.

- Application for Pesticide (EPA Form 8570-1)
- Revised label (one copy)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Thank you very much for your help. If you have any questions or concerns, please do not hesitate to contact me either by e-mail at: RHargadine@gowanco.com, by Fax at: (928) 373-1828 or by phone at: (928) 819-1531.

Sincerely,



Rebecca A. Hargadine,  
Agent for Canyon

Enclosures

