



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 9, 2026

Ana Sanchez  
Agent  
Canyon Group C/O Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366

Subject: Label Amendment - Registration Review Mitigation for Pyridaben  
Product Name: Pyridaben T&O SC  
EPA Registration Number: 81880-29  
Case Number: 473003  
Application Date: 7/10/2020

Dear Ana Sanchez:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Pyridaben Interim Decision and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Carolyn Smith by phone at (202) 566-2273, or via email at [smith.carolyn@epa.gov](mailto:smith.carolyn@epa.gov).

Sincerely,



Jaclyn Pyne, Team Leader  
Risk Management and Implementation Branch 3  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

**ACCEPTED**

02/09/2026

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under

EPA Reg. No. 81880-29

PYRIDABEN GROUP 21A ACARICIDE

# PYRIDABEN T&O SC

## Miticide/Insecticide

**ACTIVE INGREDIENT:** Pyridaben[2-*tert*-butyl-5-(4-*tert*-butylbenzylthio)-4-chloropyridazin-3(2*H*)-one] .....**OTHER INGREDIENTS:** .....

% By Wt.

42.47%

57.53%

**Total** 100.00%

Contains 3.75 pounds active ingredient per gallon.

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
If swallowed	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have the person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. <b>For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-478-0798.</b>	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)****Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves
- Protective eye wear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**NET CONTENTS:**

**Gowan**  
The Go To Company

EPA Reg. No. 81880-29  
EPA Est. No.

Distributed By:  
Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366-5569

**Engineering Controls Statement:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from target area. Drift or runoff from treated areas may be hazardous to fish in adjacent sites. This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively foraging the treatment area. Application early in the morning or at dusk is suggested.

#### PHYSICAL AND CHEMICAL HAZARDS

Do not allow contact with oxidizing agents. Hazardous chemical reaction may occur.

#### ENDANGERED SPECIES CONCERNS

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Notice of Conditions of Sale and Warranty and Liability Limitations** are to be followed.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

#### USE INFORMATION

**Pyridaben T&O SC** is a selective contact Miticide/Insecticide that controls mites and whiteflies on ornamental plants, flowers, and foliage crops. **Pyridaben T&O SC** provides knockdown and residual control. A good performance evaluation can be made 4-7 days after treatment. For optimum results, **Pyridaben T&O SC** should be applied as pest populations build prior to reaching economic thresholds.

#### Mite Resistance Management

Naturally occurring strains of mites and insects listed on this label may not be effectively controlled due to reduced sensitivity. If insensitive strains are present, use a product with a different mode of action to ensure control.

For resistance management, **Pyridaben T&O SC** contains a Group (21A) insecticide/acaricide. Any insect/mite population may contain individuals naturally resistant to **Pyridaben T&O SC** and other Group (21A) insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of **Pyridaben T&O SC** or other Group 21A insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.

- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Gowan Company representative

#### Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and by triple rinsing the equipment before and after applying this product.

#### MANDATORY SPRAY DRIFT MANAGEMENT

##### **Ground Boom Applications:**

- Apply with the nozzle height recommended by the manufacturer, but no more than 4 ft. above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

##### **Boomless Ground Applications:**

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

#### SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

##### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

##### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

##### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

##### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

##### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

##### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

##### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

##### **Boom-less Ground Applications:**

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

##### **Handheld Technology Applications:**

- Take precautions to minimize spray drift

#### APPLICATION INSTRUCTIONS

**Pyridaben T&O SC** may be applied by ground equipment using either diluted or concentrated sprays. Apply listed rates of **Pyridaben T&O SC** as instructed by **Application Directions Section**. Spray the last 3 rows windward of surface water using nozzles on only one side with the spray directed away from surface water. Spraying over the tops of trees can be prevented by adjusting or turning off the top nozzles. Shut the nozzles on the side away from the grove off when spraying the outside row. Shut the nozzles off when turning at the ends of the rows and when passing tree/vine gaps in rows.

## Coverage

Apply **Pyridaben T&O SC** in sufficient water to ensure thorough coverage of foliage and fruit. Thorough coverage is required for optimum control. Spraying alternate rows may reduce **Pyridaben T&O SC** performance. **Pyridaben T&O SC** must be applied to each row for optimum control. To achieve thorough coverage, use proper spray pressure, nozzles, nozzle spacing, volume per acre, and tractor speed. Consult spray nozzle and accessory guide for information pertaining to proper equipment calibration.

## GROUND APPLICATION (BROADCAST)

**Water Volume:** Use 100-400 gallons of spray solution per broadcast acre for optimal performance.

## ADDITIVES

In general, no additives or adjuvants are necessary for effective use of **Pyridaben T&O SC**. However, the use of additives may be considered for certain conditions such as obtaining better spray distribution, adhesion or penetration of product onto leaf or plant surfaces. Consult a Canyon representative or local agricultural authorities for more information concerning additives.

## TANK MIXING INFORMATION

The phytotoxic potential of **Pyridaben T&O SC** has been assessed on a wide variety of plants with no phytotoxicity observed. However, all varieties and cultivars have not been tested with possible tank mix combinations, sequential pesticide treatments, and adjuvants and surfactants. Local conditions can also influence crop tolerance and may not match the information under which testing had been conducted. Therefore, before using **Pyridaben T&O SC**, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

## Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the Mixing Order (see below) using 2 teaspoons for each pound or 1 teaspoon for each pint of listed label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

## Mixing Order

1. Water. Begin by agitating a thoroughly clean sprayer tank half full of clean water.
2. Products in PVA bags. Place any product contained in water soluble PVA bags. Allow time for thorough mixing.
3. Water-dispersible products (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
4. Water soluble products
5. Emulsifiable concentrates
6. Water soluble additives
7. Remaining quantity of water. Maintain constant agitation during application.

## CROPS

This product can be used on the following crops:

- **Ornamental plants**
- **Foliage plants, such as poinsettias**
- **Greenhouse tomatoes**
- **Greenhouse cucumbers**

## PESTS

This product can be used on the following pests:

COMMON NAME	SCIENTIFIC NAME
Broad mite	Family: Tarsonemidae <i>Polyphagotarsonemus latus</i>
False spider mite	Family: Tenuipalpidae <i>Brevipalpus phoenicis</i>
Citrus flat mite	<i>Brevipalpus lewisi</i>
Apple rust mite	Family: Eriophyidae <i>Aculus schlechtendali</i>
Citrus bud mite	<i>Aceria sheldoni</i>
Citrus rust mite	<i>Phyllocoptrus oleivora</i>
Peach silver mite	<i>Aculus fockeui</i>
Pear rust mite	<i>Epitrimerus pyri</i>
Pink citrus rust mite	<i>Aculops pelekassi</i>
Citrus red mite	Family: Tetranychidae <i>Panonychus citri</i>
European red mite	<i>Panonychus ulmi</i>
McDaniel spider mite	<i>Tetranychus mcdanieli</i>
Pacific spider mite	<i>Tetranychus pacificus</i>

Pecan leaf scorch mite	<i>Eotetranychus hicoriae</i>
Sixspotted mite	<i>Eotetranychus sexmaculatus</i>
Southern red mite	<i>Oligonychus ilicis</i>
Texas citrus mite	<i>Eutetranychus banksi</i>
Twospotted spider mite	<i>Tetranychus urticae</i>
Willamette spider mite	<i>Eotetranychus willamettei</i>
Citrus root weevil	Family: Curculionidae <i>Pachnaeus litus</i>
Apple aphid	Family: Aphididae <i>Aphis pomi</i>
Black pecan aphid	<i>Melanocallis caryaefoliae</i>
Black margined aphid	<i>Monellia caryella</i>
Brown citrus aphid	<i>Toxoptera citricida</i>
Yellow pecan aphid	<i>Monelliopsis pecanis</i>
Sweet potato whitefly	Family: Aleyrodidae <i>Bemisia tabaci</i>
Silverleaf whitefly	<i>Bemisia argentifolii</i>
Pear Psylla	Family: Psyllidae <i>Cacopsylla pyricola</i>
Eastern grape leafhopper	Family: Cicadellidae <i>Erythroneura comes</i>
Grape leafhopper	<i>Erythroneura elegantula</i>
Variegated leafhopper	<i>Erythroneura variabilis</i>
Virginia creeper leafhopper	<i>Erythroneura ziczac</i>
White apple leafhopper	<i>Typhlocyba pomaria</i>

#### APPLICATION DIRECTIONS

CROP	RATE	COMMENTS
Ornamental plants and Foliage plants grown in greenhouses and outdoors	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.  • Do not apply more than 34.14 oz of product per acre, per year. • Avoid drift to all other crops and non-target areas.
Greenhouse tomatoes	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.  • PHI = 2 days • Do not make more than 2 applications per crop per year. • Do not apply more than 12.8 oz of product per crop per season. • Allow a minimum of 30 days between sequential applications
Greenhouse cucumbers	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.  • PHI = 1 day • Do not make more than 2 applications per crop per year. • Do not apply more than 19.2 oz of product per crop per season. • Allow a minimum of 30 days between sequential applications. • Not for use on cucumbers grown in the field.

#### USE RESTRICTIONS

- Do not enter a treated greenhouse or a treated indoor area without protective equipment for 12 hours unless one of the following items is completed:
  - 10 air exchanges
  - 2 hours of system ventilation
  - 4 hours of ventilation using vents, windows or other passive ventilation
  - All required PPE is worn.
- Sequential Treatment: Do not use in successive miticide applications. Use in rotation with other treatments having different modes of action.
- Do not use less than 6.4 ozs of **Pyridaben T&O SC** per acre.
- Do not use with nutritional sprays that contain boron.
- Do not apply by air.
- Do not apply through any type of irrigation equipment.

- Drift: Do not apply product when weather conditions favor drift to surface water. Do not apply within 110 feet upwind of surface water or when wind speed is above 8 mph. Do not apply during a temperature inversion.
- **Pyridaben T&O SC** is not for sale, distribution, or use in Nassau and Suffolk counties in New York State. In the remainder of the state, read and follow all applicable directions, restrictions and precautions on this label.

#### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place away from heat or open flame.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**FOR 24 HOUR EMERGENCY ASSISTANCE (SPILL, LEAK, OR FIRE) CALL CHEMREC® (800) 424-9300  
For other information, contact Gowan Company or see Safety Data Sheet.**

#### NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

**Important:** Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

All other brands are registered trademarks of their respective owners.

EPA TEXT PENDING: Pyridaben T&O SC (to EPA 1-28-21 Reg Review ONLY)