

81927-18

4-4-2008

1/47



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
HUMAN SAFETY
NOTIFICATION

Michael Kellog
Agent for Alligare, LLC
Pyris Regulatory Consulting, Inc.
4110 136th St. N.W.
Gig Harbor, WA 98332

APR 4 2008

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Mr. Kellog:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated March 20, 2008 for:

EPA Registration 81927-~~12~~¹⁸ Alligare Picloram 22K

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-8893.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

PYXIS REGULATORY CONSULTING, INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
www.PyxisRC.com

2/47

March 20, 2008

DR

COURIER DELIVERY

Jim Tompkins (PM 25)
Document Processing Desk (**NOTIF**)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

RE: Alligare, LLC – Alligare Picloram 22K (EPA Reg. No. 81927-18)
Revision to Container Disposal Instructions per PRN 2007-4

Dear Mr. Tompkins,

On behalf of Alligare, LLC please find the enclosed label notification revising the container disposal instructions per PRN 2007-4.

In support of this notification submission, we submit the following documents:

1. Completed Application for Registration (EPA Form 8570-1)
2. One (1) copy of the Alligare Picloram 22K labeling with changes tracked
3. One (1) copy of the Alligare Picloram 22K labeling with changes incorporated
4. Letter of Authorization

Please feel free to contact me by phone (253) 853-7369 or by email at Mike@PyxisRC.com if you have any questions or need any additional information.

Sincerely,



Michael Kellogg

Enclosures

cc: Jay Golz; Alligare, LLC

3147



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

| | | |
|---|---|--|
| 1. Company/Product Number 81927-12 | 2. EPA Product Manager J. Tompkins | 3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted |
| 4. Company/Product (Name) Alligare, LLC / Alligare Picloram 22K | PM# 25 | |
| 5. Name and Address of Applicant (Include ZIP Code) Alligare, LLC c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Gig Harbor, WA 98332 <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ | |

Section - II

Amendment - Explain below. Final printed labels in response to Agency letter dated _____

Resubmission in response to Agency letter dated _____ "Me Too" Application.

Notification - Explain below. Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

| | | | |
|---|--|---|---|
| Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | 2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| * Certification must be submitted | | If "Yes" Unit Packaging wgt. No. per container | If "Yes" Package wgt No. per container |

3. Location of Net Contents Information: Label Container

4. Size(s) Retail Container: 1 qt., 1 gal., 2.5 gal.

5. Location of Label Directions: On Label On Labeling accompanying product

6. Manner in Which Label is Affixed to Product: Lithograph Paper glued Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

| | | |
|-------------------------|----------------|---|
| Name Michael Kellogg | Title Agent | Telephone No. (Include Area Code) (253) 853-7369 |
|-------------------------|----------------|---|

Certification
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received:
(Stamped)

| | |
|----------------------------------|--------------------|
| 2. Signature | 3. Title Agent |
| 4. Typed Name Michael Kellogg | 5. Date 3/20/08 |

4/47

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible, non-target plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

Alligare Picloram 22K

For use in areas west of the Mississippi River for the control of susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, spring seeded wheat, barley and oats not underseeded with a legume, non-cropland, and on Conservation Reserve Program (CRP) acres.

ACTIVE INGREDIENT:

Picloram: 4-amino-3,5,6-trichloropicolinic acid, potassium salt.....24.4%

OTHER INGREDIENTS:.....75.6%

TOTAL:.....100.0%

Alligare Picloram 22K contains the following acid equivalent:

Picloram: 4-amino-3,5,6-trichloropicolinic acid – 21.1% (2 lbs./gal.)

**KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

| FIRST AID | |
|---|---|
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information. | |

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals.

EPA Reg. No. 81927-18

EPA Est. No.

Manufactured For:
Alligare, LLC
13 N. 8th Street
Opelika, AL 36801

Net Contents:

5/47

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves Category A, such as barrier laminate \geq 14 mils, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to some plants at very low concentrations. Non-target plants may be adversely affected if pesticide is allowed to drift from areas of application. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of wastes or rinsate. Do not allow runoff or spray to contaminate wells, irrigation ditches or any body of water used for irrigation or domestic purposes. Do not make application when circumstances favor movement from treatment site.

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, picloram may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetative filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not intended for manufacturing or formulating.

6/47

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves Category A, such as barrier laminate \geq 14 mils, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications on rangeland, permanent grass pastures, and non-cropland, do not enter or allow worker entry into treated areas until sprays have dried, unless applicator and other handler PPE is worn.

GENERAL INFORMATION

In areas west of the Mississippi River use Alligare Picloram 22K to control susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, spring seeded wheat, barley and oats not underseeded with a legume, non-cropland, and on Conservation Reserve Program (CRP) acres. This product is NOT for sale or use in the San Luis Valley of Colorado.

PRECAUTIONS AND RESTRICTIONS

Use this product only as specified on this label.

Observe any special use and application restrictions and limitations, including method of application and permissible areas of use as promulgated by state or local authorities.

Do not mix with dry fertilizer.

Chemigation: Do not apply this product through any type of irrigation system.

Grazing Poisonous Plants: Application of this herbicide may increase the palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.

7/97

Maximum Use Rates:

Non-cropland Areas: Total use of picloram, including retreatments or spot treatments, must not exceed 1 lb ae per acre picloram (2 quarts per acre Alligare Picloram 22K) per annual growing season.

Rangeland and Permanent Grass Pastures: For control of noxious weeds as defined by federal, state, or local authorities, picloram may be applied at up to 1 lb ae per acre (2 quarts Alligare Picloram 22K per acre) per annual growing season as a broadcast treatment. Spot treatments may be applied at the equivalent broadcast rate of up to 1 lb ae picloram (2 quarts Alligare Picloram 22K) per acre.

For control of other broadleaf weeds and woody plants, picloram may be applied broadcast at up to 0.5 lb ae per acre (1 quart Alligare Picloram 22K per acre) per annual growing season. Apply spot treatments at an equivalent broadcast rate of up to 1 lb ae per acre (2 quarts Alligare Picloram 22K per acre) per annual growing season, but not more than 50% of an acre may be treated. Repeat treatments may be applied as necessary, but total use must not exceed the maximum amount specified.

Cropland (Spring-Seeded Wheat, Barley and Oats): Do not apply more than 0.09 lb ae picloram (1 1/2 fluid ounces of Alligare Picloram 22K) per acre during the small grain growing season.

Fallow Cropland (Not Rotated to Broadleaf Crops): Do not apply more than 1 lb ae picloram per acre (1 pint per acre of Alligare Picloram 22K) as a broadcast treatment per annual growing season.

Conservation Reserve Program (CRP) for Seeding to Permanent Grasses Only: Do not broadcast apply more than 0.5 lb ae per acre of picloram (1 quart per acre of Alligare Picloram 22K) per annual growing season or apply more than 1 lb ae per acre picloram (2 quarts per acre of Alligare Picloram 22K) per annual growing season as a spot application. To reduce potential damage to subsequent small grain crops, use the lower rate or discontinue the use of Alligare Picloram 22K at least 2 years prior to the seeding of small grain crops. After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay shows that no detectable picloram is present in the soil.

Precautions for Avoiding Injury to Non-Target Plants

- Do not move treated soil to other areas or use it to grow plants if they are not registered for use with picloram until an adequate sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.
- Do not spray if the loss of forage legumes, including clover cannot be tolerated. Alligare Picloram 22K may injure or kill legumes. New legume seedlings may not grow for several years following application of this product.
- Do not make application when circumstances favor movement from treatment site.
- Do not rotate to food or feed crops on treated land if they are not registered for use with picloram until an adequately sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.
- Do not contaminate water intended for irrigation, or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or runoff to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not apply Alligare Picloram 22K on residential or commercial lawns or near ornamental trees and shrubs. Untreated trees can occasionally be affected by root uptake of herbicide through movement into the topsoil or by excretion of the product from the roots of nearby treated trees. Do not apply Alligare Picloram 22K within the root zone of desirable trees unless such injury can be tolerated.
- Allow 7 days of grazing on an untreated grass pasture (or feeding of untreated hay) before transferring livestock from treated grazing areas (or feeding of treated hay) onto sensitive broadleaf crop areas. Otherwise, urine and manure may contain enough picloram to cause injury to sensitive broadleaf plants.
- Do not use manure from animals grazing treated areas or feeding on treated hay on land used for growing broadleaf crops, ornamentals, orchards or other susceptible, desirable plants. Manure may contain enough picloram to cause injury to susceptible plants.

- Do not use grass or hay from treated areas for composting or mulching of susceptible broadleaf plants.
- Do not apply to snow or frozen ground. Application during very cold (near freezing) weather is not advisable.
- Do not use on flood or sub-irrigated land.

Sprayer Clean-Out: To avoid injury to desirable plants, equipment used to apply Alligare Picloram 22K should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
2. Rinse a second time, adding 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15-20 min.). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed and cleaned separately.

Precautions for Avoiding Injurious Spray Drift

For aerial applications on rights-of-way or other areas near susceptible crops, use drift control additive as recommended by the manufacturer.

Do not apply or otherwise permit Alligare Picloram 22K or spray containing Alligare Picloram 22K to contact crops or other desirable broadleaf plants, including but not limited to alfalfa, beans, grapes, melons, peas, potatoes, safflower, soybeans, sugar beets, sunflower, tomatoes, and other vegetable crops, flowers, fruit plants, ornamentals or shade trees or the soil containing roots of nearby valuable plants.

Applications should be made to avoid spray drift since very small quantities of spray, which may not be visible, may seriously injure susceptible crops during both growing and dormant periods. To minimize spray drift, use low nozzle pressure; apply as a coarse spray; and use nozzles designed for herbicide application that do not produce a fine droplet spray. To aid in further reducing spray drift, a drift control and deposition aid may be used with this product, especially when water alone is used as the carrier. If a drift control aid is used, follow all use recommendations and precautions on the product label. Do not use a thickening agent with Microfoil or Thru-Valve booms, or other systems that cannot accommodate thick sprays.

Ground Equipment: With ground equipment, spray drift can be reduced by keeping the spray boom as low as possible; by applying 10 gallons or more of spray per acre; by keeping the operating spray pressures at manufacturer's recommended minimum pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); by spraying when wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to air inversions. Direct sprays no higher than the tops of target vegetation and keep spray pressures low enough to provide coarse spray droplets to minimize drift. A spray thickening agent may be used to further reduce the potential for drift.

Aerial Application:

AERIAL SPRAY DRIFT MANAGEMENT

Spray Drift Management

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

9/47

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{1}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Aerial Drift Reduction Advisory

[This section is advisory in nature and does not supersede the mandatory label requirements.]

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than $\frac{1}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2

U.S. GOVERNMENT PRINTING OFFICE: 1981 O-101-101

10/47

mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

MIXING INSTRUCTIONS

Mix the required amount of Alligare Picloram 22K in water and apply as a coarse, low pressure spray using ground equipment or aircraft. Use enough spray volume to provide uniform coverage of the weeds.

Use with Surfactants: Addition of wetting or penetration agents is not usually necessary when using Alligare Picloram 22K. However, under certain conditions, such as drought, addition of a surfactant may improve efficacy. However, if foliar burn occurs too rapidly, translocation of Alligare Picloram 22K will not occur and control of perennial weeds, such as field bindweed, may be reduced.

Mixing with Water

To prepare the spray, add about half the desired amount of water in the spray tank. Then with agitation, add the recommended amount of Alligare Picloram 22K and other registered tank mix herbicides. Finally, with continued agitation, add the rest of the water and additives such as surfactants or drift control and deposition aids.

Mixing Oil-Water Emulsions (Ground and Aerial Applications)

For aerial application, add oil to the total spray mix at the ratio of 1 part oil to 5 parts water (1:5 ratio). For ground application, add oil to the spray mix at a rate of 5 to 10% of the total mix. Do not use more than 1 gallon of oil per acre for aerial or ground application. Use agricultural spray emulsifiers such as Sponto 712 or Triton X-100 according to mixing instructions given below.

Batch Mixing Instructions

With continuous, vigorous agitation:

1. Add half the amount of water to be used to the spray tank.
2. Add the required amount of water-soluble herbicides such as Alligare Picloram 22K, Reclaim™ herbicide or 2,4-D Amine.
3. With continued, vigorous agitation, slowly add a premix of oil, emulsifier and oil soluble herbicides such as Remedy™ herbicide or a 2,4-D ester as required. Note: Do not add water or mixtures containing water to the premix or oil soluble herbicide since a thick "invert" (water in oil) emulsion may

11/47

be formed that will be difficult to break. An invert emulsion will also form if the premix is added to the mixing tank before the addition of water.

4. Finish filling the spray tank and maintain sufficient agitation to ensure uniformity of the spray mixture during application.

Mixing with Sprayable Liquid Fertilizer Solutions

Alligare Picloram 22K is compatible with most non-pressurized liquid fertilizer solutions; however, a compatibility test (jar test) should be made prior to mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank.

Note: The lower the temperature of the liquid fertilizer, the greater the likelihood of mixing problems.

Use of a compatibility aid such as Unite or Compex may help obtain and maintain a uniform spray solution during mixing and application. Compatibility is best with straight liquid nitrogen fertilizer solutions. Mixing with N-P-K fertilizer solutions or suspensions is more difficult and should not be attempted without first conducting a successful jar test. Agitation in the spray tank must be vigorous to be comparable with jar test agitation. For best results, liquid fertilizer rates should not exceed 50% of the total spray volume. Premix Alligare Picloram 22K with water and add to the liquid fertilizer/water mixture while agitating contents of the spray tank. Apply the spray the same day it is prepared while maintaining continuous agitation. Rinse spray tank thoroughly after use.

Note: Foliar applied liquid fertilizers used as carrier for Alligare Picloram 22K can cause yellowing or leaf burn of crop foliage.

Do not use spray equipment used to apply Alligare Picloram 22K for other applications to land planted to, or to be planted to susceptible crops or desirable sensitive plants, unless it has been determined that all phytotoxic residue of this herbicide has been removed by thorough cleaning of equipment.

Local conditions may affect the use of herbicides. State agricultural experiment stations or extension service weed specialists in many states issue recommendations to fit local conditions. Be sure that use of this product conforms to all applicable regulations.

Tank Mixing

Alligare Picloram 22K may be applied in tank mix combination with labeled rates of 2,4-D or other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates. If products containing the same active ingredient are tank mixed, do not exceed the maximum allowable active ingredient use rates.
- For products packaged in water-soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See "Sprayer Clean-Out" under "Precautions for Avoiding Injury to Non-Target Plants".)
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Alligare Picloram 22K and other pesticides or carriers. Use a clear glass jar with lid and mix the tank mix ingredients in their relative proportions. The tank mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if

12/47

separation occurs, should readily mix if agitated. An incompatible mixture is indicated by separation into distinct layers which do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film on the jar.

APPLICATION METHODS

Ground or Aerial Broadcast

Use Alligare Picloram 22K as a broadcast treatment by ground or aerially to control listed broadleaf weeds and woody plants. Apply Alligare Picloram 22K as a coarse low-pressure spray at the recommended rates in a spray volume of 1 or more gallons per acre by air or 10 or more gallons per acre by ground. Aerial applications can be applied using aerial electrostatic spray charging systems.

High-Volume Foliar Applications

Spray to thoroughly wet foliage and stems. An approved agricultural surfactant may be added at the manufacturer's recommended rate. Do not apply more than the maximum application rate of Alligare Picloram 22K specified for a given treatment site.

Spot Treatment

Use application rates as suggested in the "Approved Uses" section of this label or recommended by your area weed control specialist. Apply in a total spray volume of 20 to 100 gallons per acre. To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to directions provided below. Do not exceed maximum application rates for Alligare Picloram 22K for a given treatment site. On rangeland and permanent grass pastures, spot treatments may be applied at an equivalent broadcast rate of up to 2 quarts per acre per annual growing season, but not more than 50% of an acre may be treated. Repeat treatments may be applied as necessary, but total use must not exceed the maximum amount specified.

Hand-Held Sprayers: Hand-held or backpack sprayers may be used for spot applications of Alligare Picloram 22K if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq. ft. Mix the amount of Alligare Picloram 22K (fl. oz. or ml) corresponding to the desired broadcast rate in 0.5 to 2.5 gallons of water, depending on the spray volume required to treat 1,000 sq. ft. To calculate the amount of Alligare Picloram 22K required for larger areas, multiply the table value (fl. oz. or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3,500 sq. ft., multiply the table value of 3.5 (calc. 3,500/1,000 = 3.5). An area of 1,000 sq. ft. is approximately 10.5 x 10.5 yards (strides) in size.

| Amount of Alligare Picloram 22K per 1,000 sq. ft. to Equal Specified Broadcast Rate | | | | | |
|---|-------------------------|-------------------------|-----------------------|------------------------|----------------------|
| ¼ pt./acre | 1/3 pt./acre | ½ pt./acre | 2/3 pt./acre | 1 pt./acre | 1 qt./acre |
| 1/10 fl. oz. (2.7 ml) | 1/8 fl. oz. (3.6 ml) | 1/5 fl. oz. (5.4 ml) | ¼ fl. oz. (7.3 ml) | 3/8 fl. oz. (11 ml) | ¼ fl. oz. (22 ml) |

1 fl. oz. = 29.6 (30) ml

Special Application Methods

Wick Application – Mix 1 part of Alligare Picloram 22K with 2 parts of water to prepare a 33% solution. Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained when applications are made prior to the early bud stage. Drain the wick applicator and clean after each use. Change ropes when flow is reduced from wear, extended use, poor cleaning or intermittent use.

Carpet Roller Application – Alligare Picloram 22K can be applied with carpeted rollers where drift presents a hazard to susceptible crops, surface waters and other sensitive areas. Apply to previously untreated plants less than 6 feet tall, and short enough to pass beneath the tractor without breaking off at the ground. Applications made during periods of extended drought conditions will not provide acceptable control. Do not burn, mow or otherwise disturb the treated plants during the remainder of the growing season. Operate carpeted rollers as close to the ground as possible without breaking the stems, but

above the tallest grasses. Grasses growing adjacent to treated plants may exhibit temporary injury. Maximize herbicide deposition on stems and foliage and minimize drippage losses by rotating the carpeted roller at 30 to 40 rpm with the lower edge moving in the same direction as the direction of travel. Maintain the carpet sufficiently wetted to apply up to 1 gal./acre of herbicide-water mixture to stands of average density (100 to 200 plants/acre), and up to 2 gals./acre in dense stands (300 to 400 plants/acre). Rewet rollers at regular intervals. See the Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River - Carpet Roller Use Recommendations) section of this label for treatment recommendations.

Soil Spot Concentrate: Alligare Picloram 22K may be applied undiluted as a spot concentrate application to control ashe juniper, eastern redcedar and eastern persimmon (see specific user directions for these plant species which follow). Applications should precede periods of expected rainfall. Apply directly to the soil within the dripline and on the upslope side of the tree. Applications to trees taller than 12 feet are not recommended.

Special Ground Sprayer Equipment: To control annual and perennial weed species using special low-volume, minimum drift equipment, such as the hooded Forage Chemical Mower, apply 1 to 2 pts. of Alligare Picloram 22K in total volumes ranging from 1 gallon to 5 gallons per acre in water alone or as an oil-water emulsion at a 1:5 and 1:4 oil-to-water ratio for a 1 gallon and 5 gallon per acre solutions, respectively.

WOODY PLANTS AND BROADLEAF WEEDS CONTROLLED BY ALLIGARE PICLORAM 22K

Woody Plants:

- | | |
|--------------------|-----------------------------|
| acacia, blackbrush | huisache (suppression only) |
| acacia, catclaw | junipers/cedars |
| acacia, twisted | locust |
| aspen | mesquite |
| broom, Scotch | pine; pinyon |
| camelthorn | rabbitbrush, Douglas |
| chaparral sp. | rose, Macartney |
| gorse | rose, multiflora |
| granjeno | sagebrush, fringed |
| guajillo | tallowtree, Chinese |
| | wormwood, absinth |

Annual and Perennial Broadleaf Weeds:

- | | | |
|---------------------------------|--------------------------------|-------------------------------|
| bindweed, field (p) | ironweed (p) | ragwort, tansy (b) |
| bitterweed (a) | knapweed, diffuse (a) | Russian thistle (a) |
| broomweed, annual (a) | knapweed, Russian (p) | skeletonweed, rush (p) |
| buckwheat, wild (a) | knapweed, spotted (p) | smartweed (a) |
| buffalobur (a) | knapweed, squarrose (p) | snakeweed, broom (p) |
| bullnettle (p) | lambsquarters (a) | sneezeweed, bitter (a) |
| bursage (a) | larkspur, geyer (p) | sowthistle, perennial (p) |
| burroweed (p) | larkspur, plains (p) | spurge, leafy (p) |
| camphorweed (a) | larkspur, tall (p) | St. Johnswort (p) |
| carrot, wild (b) | lettuce, prickly (a) | starthistle, Iberian (a) |
| cinquefoil, sulfur (p) | licorice, wild (p) | starthistle, purple (a) |
| cocklebur (a) | locoweeds (p) | starthistle, yellow (a) |
| coneflower, upright prairie (p) | loco, woolly (p) | sunflower (a) |
| croton (a) | loco, Wooten (garbancillo) (p) | tasajillo (p) |
| crupina, common (a) | lupines (p) | thistles, annual or biennial, |
| daisy, ox-eye (p) | marshelder (sumpweed) (a) | including: |

14/47

- | | | |
|-------------------------------|-----------------------------|--------------------------------|
| dock, curly (p) | mayweed (a) | thistle, bull (b) |
| garbancillo (Wooten loco) (p) | milkweed (p) | thistle, distaff (a) |
| goldaster, gray (p) | mustard, wild (a) | thistle, Italian (b) |
| goldaster, narrowleaf (p) | nightshade, silverleaf (p) | thistle, musk (b) |
| goldenrod, common (p) | pennycress (a) | thistle, plumeless (b) |
| goldenweed, Drummond (p) | pigweed (a) | thistle, Scotch (b) |
| groundsel (p) | pricklypear, plains (p) | thistles, perennial, including |
| henbane, black (a,b) | pricklypear, lindheimer (p) | thistle, Canada (p) |
| horsenettle, Carolina (p) | ragweed, bur (a) | thistle, wavy leaf (p) |
| horsenettle, western (p) | ragweed, common (a) | toadflax, dalmation (p) |
| horsenettle, white (p) | ragweed, lanceleaf (a) | toadflax, yellow (p) |
| horseweed (a) | ragweed, western (a) | yankeeweed (p) |
- (a) - annual; (b) - biennial; (p) - perennial

SPECIFIC USE DIRECTIONS FOR RANGELAND, PERMANENT GRASS PASTURES AND NON-CROPLAND

General Requirements for Non-Cropland Areas

Use Alligare Picloram 22K to control susceptible broadleaf weeds and woody plants on non-cropland areas such as roadsides or other rights-of-way, fence rows, and around farm buildings. Up to 2 quarts of Alligare Picloram 22K per acre may be applied. See "Rangeland and Permanent Grass Pastures" section for specific target weed or woody plant species treatment recommendations.

Maximum Use Rates for Non-Cropland Areas: Total use of Alligare Picloram 22K, including retreatments or spot treatments must not exceed 2 quarts per acre per annual growing season.

General Requirements for Rangeland and Permanent Grass Pastures for States West of the Mississippi River

Use Alligare Picloram 22K on rangeland and permanent grass pastures to control susceptible broadleaf weeds and woody plants including, but not limited to those shown in the following tables. Many annual weeds at the seedling stage are controlled at the rate of 1 pt. per acre. Where a rate range is recommended, choose the higher rate for dense weed infestations, and for more dependable, longer-lasting control. Lower rates will perform best when applied under favorable conditions and at the optimum growth stage, but may provide a lower level of control and require retreatment. For best results, treat when weeds are small and actively growing in the spring before full bloom, however, certain weeds may also be treated in late summer or fall. Treatments during full bloom or seed stage of some weeds may not provide acceptable control.

Refer to the "Application Methods" section of this label for information on various methods of application including ground or aerial broadcast, high volume foliar application, spot treatments, and special application methods for certain weeds or woody plants including spot concentrate application or application with special low volume or hooded spray equipment.

Precautions and Restrictions:

Maximum Use Rates for Rangeland and Permanent Grass Pastures: For control of noxious weeds as defined by federal, state, or local authorities, Alligare Picloram 22K may be applied at up to 2 quarts per acre per annual growing season as a broadcast treatment. Spot treatments may be applied at the equivalent broadcast rate of up to 2 quarts per acre.

For control of other broadleaf weeds and woody plants, Alligare Picloram 22K may be applied broadcast (ground, air, or high volume foliar) at up to 1 quart per acre per annual growing season. Spot treatments (hand sprayer, calibrated boom, high volume foliar, or soil spot concentrate) may be applied at an equivalent broadcast rate of up to 2 quarts per acre per annual growing season, but not more than 50% of

ALLIGARE PICLORAM 22K

15/47

an acre may be treated. Repeat treatments may be applied as necessary, but total use, must not exceed the maximum amount specified.

Grazing Restrictions: Do not cut grass for feed within two weeks after treatment when applying more than 1 quart of Alligare Picloram 22K per acre. Meat animals grazing for up to two weeks after treatment should be removed from treated areas three days prior to slaughter. Do not graze lactating dairy animals on treated areas within two weeks after treatment.

Certain established grasses, such as bromegrass, blue gamma, and buffalograss may be suppressed when using rates of Alligare Picloram 22K over 1 quart per acre. However, subsequent grass growth should be improved by release from weed competition.

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River, Except Texas, Oklahoma and New Mexico)

Apply Alligare Picloram 22K broadcast or as a spot application unless otherwise specified (See "Application Methods" section of this label). For additional species or more specific rates, consult your area's current Weed Control Guide and/or your local Alligare, LLC representative. Rates in the following table are advisory in nature and do not supersede the maximum rates for the use site allowed for different methods of application.

Weed Control Prior to Seeding Perennial Grasses:

Weed control with Alligare Picloram 22K fits into the following grass re-vegetation programs. Alligare Picloram 22K may be applied in the spring or early summer, depending on the target weed species and grass seed planted in the fall when conditions are favorable. Alternatively, Alligare Picloram 22K may be applied in the fall and grass seed planted in the winter or fall when conditions are favorable.

Apply Alligare Picloram 22K at 1 qt./acre or less; see rate chart for information on control of target weed species. Depending on grass species sensitivity, there may be temporary injury on new plantings when Alligare Picloram 22K is applied at 1 qt./acre. However, this injury will be insignificant in comparison with the benefit to grasses due to the removal of weed competition. Germination of annual grass species may be suppressed after treatment. To optimize weed control, it is suggested the application area be disturbed as little as possible by the seeding operation. At the very least, the site should be left undisturbed for 14 days prior to seedbed preparation or seeding. To decrease the potential for injury on sensitive grass species, increase the interval between application and seeding.

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|--|---|---|
| Annual and Biennial Weeds: | | |
| bursage (bur ragweed) crupina, common henbane, black horseweed starthistle, Iberian starthistle, purple starthistle, yellow | 1-2 pts. Alligare Picloram 22K | Apply when there is adequate soil moisture and weeds are actively growing. |
| thistles, including, bull distaff Italian musk plumeless scotch | Fall: ½ - ¾ pt. Alligare Picloram 22K Spring: ½ - ¾ pt. Alligare Picloram 22K + 1 lb. a.e. 2,4-D | General: Apply at the rosette stage before bolting in the spring or in the fall prior to soil freeze up. Distaff Thistle: Apply at rosette stage in spring only. Bolted Musk Thistle: Apply before flowering at the rate of ½ - 1 pt. of Alligare Picloram 22K plus 1 lb. a.e. of 2,4-D/acre. |

16/47

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|--|---|--|
| Perennial Weeds: | | |
| pricklypear, plains | ½ - 1 pt. Alligare Picloram 22K | Apply at peak of flowering. Use of an oil-water emulsion spray mixture may improve control. Lower rate will provide partial control (stand reduction) and high rate more complete control. Treatment response is slow and may continue for 2 years or longer. |
| sagebrush, fringed | ½ - 1 pt. Alligare Picloram 22K + 1 lb. a.e. 2,4-D ester | Apply after seed stalk elongation and early flowering and throughout the summer if growing conditions are favorable. |
| cinquefoil, sulfur larkspur, geyer larkspur, plains locoweeds snakeweed, broom | 1 pt. Alligare Picloram 22K | General: Apply when weeds are actively growing. Sulfur cinquefoil: Apply during active growth or fall regrowth. Geyer larkspur: Apply when plant is actively growing between rosette stage and flower bud formation. Locoweeds: Apply from early bud to early bloom stage. See "General Use Precautions" section of this label for note on grazing treated poisonous plants. Broom snakeweed: Apply during active growth between full leaf to early bloom stage. |
| burroweed daisy, ox-eye goldenrod, common knapweed, diffuse knapweed, spotted knapweed, squarrose rabbitbrush, Douglas thistle, Canada thistle, wavy leaf wormwood, absinth | 1-2 pts. Alligare. Picloram 22K | General: Lower rates in rate range may require annual spot treatments. Control with lower rates may be improved by tank mixing with 1.0 lb. a.e. per acre 2,4-D. Goldenrod: Apply during active growth prior to bud stage. Diffuse or spotted knapweed: Optimum application from rosette to mid-bolting stage or to fall regrowth. Thistle (Canada and Wavy Leaf): Apply when most basal leaves have emerged, but before bud stage, or apply to regrowth in the fall. Apply rates less than 1 ½ pts./acre only under favorable conditions and in combination with 1 lb. a.e./acre of 2,4-D; retreatment may be required. Absinth wormwood: Apply in spring or early summer when plants are actively growing. |
| licorice, wild milkweed | 2 pts. Alligare Picloram 22K | Wild Licorice: Apply at bloom stage. Milkweed: Treat during active growth and tank mix recommended rate of Alligare Picloram 22K with 1 lb. a.e./acre 2,4-D. |
| bindweed, field gorse lupines knapweed, Russian | 2-4 pts. Alligare Picloram 22K | General: Annual retreatment of these species will be required at rates at low end of rate range. Control at low end of rate range may be improved by tank mixing |

17/47

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|---|---|--|
| ragwort, tansy skeletonweed, rush spurge, leafy St. Johnswort toadflax, dalmation | | with 1 lb. a.e. 2,4-D. Russian Knapweed: Apply during active growth from bud to mid-flowering, or to fall regrowth. Leafy Spurge: Apply at true flower stage of growth or apply to fall regrowth. Re-apply when level of control falls below 80 percent. Dalmation Toadflax: Apply when plants are actively growing through full bloom stage of growth. |
| larkspur, tall sowthistle, perennial toadflax, yellow | 4 pts. Alligare Picloram 22K | General: A retreatment program may be necessary for satisfactory control of these species. Tall Larkspur: For best results apply from 6 inches tall to late bloom stage. For increased control, apply in tank-mix with Ally [®] or Escort [®] herbicide and non-ionic surfactant. See "General Use Precautions" section of this label for note on grazing treated poisonous plants. |
| Woody Plants: | | |
| juniper | 4 qts. Alligare Picloram 22K per 100 gallons of spray | Apply as a high volume foliar spray/individual plant treatment. |
| redcedar, eastern | Eastern red cedar can be controlled with spot concentrate applications of Alligare Picloram 22K in either the spring (April-May) or fall (September-October). For best results, use 3 ml to 4 ml of Alligare Picloram 22K (undiluted) per 3 feet of plant height. Application should precede periods of expected rainfall. Apply directly to soil within the dripline and on the upslope side of the tree. Application to trees taller than 15 feet is not recommended. Do not use more than 2 pints of Alligare Picloram 22K per acre in any one year. | |

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: Texas, Oklahoma and New Mexico)

Apply Alligare Picloram 22K as a broadcast treatment by ground or air, as a spot application, or high volume foliar application to control listed broadleaf weeds and woody plants, unless otherwise specified (See "Application Methods" section of this label). Refer to the "General Requirements for States West of the Mississippi River" for maximum use rates. To control certain species, Alligare Picloram 22K may be applied alone or in combination with 2,4-D as indicated in the following table. When Alligare Picloram 22K is applied alone, herbicide symptoms will be slower to appear than when it is applied in combination with 2,4-D.

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|--|--|-------------------------------------|---|
| Annual and Biennial Weeds: | | | |
| bitterweed, western broomweed, annual buffalobur | Early Season: ½ - 1 pt. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + | General: Apply when there is adequate soil moisture and weeds are actively growing. |

18/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|---|--|--|---|
| bursage (bur ragweed) camphorweed carrot, wild cocklebur croton horseweed lettuce, prickly marshelder (sumpweed, sulfaweed) ragweed, common ragweed, lanceleaf ragweed, western smartweed sneezeweed, bitter sunflower thistle, bull thistle, musk | + 0.5 lb. a.e. 2,4-D or ¾ - 1 ½ pts. Alligare Picloram 22K Mid to Late Season: 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1-2 pts. Alligare Picloram 22K | 2 lbs. a.e. 2,4-D | Early Season: Recommendations are intended only for very early in the season when weeds are no more than 2 to 3 inches tall. Mid to Late Season: Recommendations are for weeds from 3 inches tall to early flowering. Marshelder: Use a minimum of ¾ pt./acre of Alligare Picloram 22K plus 0.75 lb. a.e./acre 2,4-D + non-ionic surfactant. Apply when plants are no more than 3-6 inches tall. Control may be improved by adding ammonium sulfate at 17 lbs./100 gals. spray solution. Thistles: Apply when thistles are in the rosette stage before bolting. Lanceleaf Ragweed: Use the higher rate within the recommended rate range. |
| Perennial Weeds: | | | |
| snakeweed, broom | Fall, Early Winter 1 pt. Alligare Picloram 22K Late Winter, Early Spring 2 pts. Alligare Picloram 22K | | Fall and Early Winter: If rainfall is less than average prior to flowering, apply after flowering is complete. If rainfall is average to above average prior to or during flowering, apply during full flower and/or active pollination, before resumption of new top growth. Late Winter and Early Spring: Apply following sufficient precipitation (rain or snow) to stimulate active plant growth. Both basal and terminal leaves should be green and active plant growth occurring. |
| bullnettle coneflower, upright prairie dock, curly horsenettle, Carolina horsenettle, western horsenettle, white ironweed nightshade, silverleaf yankeeweed | ½ - 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1-2 pts. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D | General: Apply when there is adequate soil moisture and weeds are actively growing. Nettles and Silverleaf Nightshade: Apply when plants begin to flower in spring. Upright Prairie Coneflower: Apply when plants are 2-6 inches tall, before flowering. Curly Dock: Apply up to bolting. Ironweed: Apply up to bud stage. Yankeeweed: Apply when plants are 8 to 10 inches tall. |

19/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|---|--|--|---|
| goldaster, gray goldaster, narrowleaf goldenweed, common goldenweed, Drummond (Isocoma spp.) | 1-2 pts. Alligare Picloram 22K + 0.5-1 lb. a.e. 2,4-D or 2 pts. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4- -D | Gray and Narrowleaf Goldaster: Apply in oil-water emulsion in spring during bud stage (prebloom). Thorough coverage is essential. Goldenweed: Apply in spring (April-June) when there is substantial canopy development as a result of good growing conditions. Add an agricultural surfactant at 0.25% - 0.5% or apply in oil-water emulsion. Increase spray volume, 4-5 gpa by air or 15-20 gpa by ground, to ensure thorough coverage. |
| Poisonous Plants such as groundsel (Senecio spp.) loco, woolly loco, Wooton (garbancillo) | ¾ - 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1 ½ - 2 pts. Alligare Picloram 22K | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4- D | General: Apply in fall or winter when there is adequate soil moisture and weeds are actively growing. See the "General Use Precautions" section of this label for note on grazing treated poisonous plants. Locoweeds: To improve wetting of locoweeds, use an agricultural surfactant at 0.25%-0.5% or apply in oil-water emulsion. |
| Woody Plants: | Tank Mixing: Within rate ranges for listed products, consult local recommendations | | |
| huisache (suppression) | 2 pts. Alligare Picloram 22K + 1 pt. Remedy™ | 2 qt. Alligare Picloram 22K + 1 qt. Remedy™ | Fall application is recommended, however, fall applications will not provide satisfactory control of other woody species in the South Texas mixed brush complex. Performance can be erratic. |
| juniper, including, alligator redberry Utah one-seeded eastern redcedar pinyon pine | -- | 4 qts. Alligare Picloram 22K | Apply May through July. Complete coverage is essential. Results with ashe juniper may be variable with high volume foliar application. |
| pricklypear, lindheimer (unburned rangeland) | 2 pts. Alligare Picloram 22K | 4 qts. Alligare Picloram 22K | Application may be made anytime, but optimum time is late August to early November. Onset of herbicidal activity is very slow and may continue for two years or longer. Good coverage is essential. |
| pricklypear, lindheimer (burned rangeland) | 1 pt. Alligare Picloram 22K | 2 qts. Alligare Picloram 22K | Conduct intense controlled burns from December through March and apply Alligare Picloram 22K mid-April through May. Rainfall |

20/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|--|---|---|--|
| | | | following burning can also stimulate prolific resprouting of the burned plants. Good coverage is also essential. |
| rose, Macartney rose, multiflora | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D | 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4-D | Apply in the spring or fall when conditions are favorable for plant growth. Use an agricultural surfactant (0.5% v/v) or apply as an oil-water emulsion. Ensure thorough and uniform coverage by applying at higher spray volume, 5 or more gpa by air or 20 or more gpa by ground. Avoid treatment less than 9 to 12 months after mowing when plants have a high percentage of new growth. Repeat treatment as necessary. |
| tallowtree, Chinese | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D or 1 pt. Remedy™ | 2 qts. Alligare Picloram 22K or 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4-D or 1 qt. Remedy™ | Apply in the spring or fall when conditions are favorable for plant growth. Use an agricultural surfactant (0.5% v/v) or use an oil-water emulsion and higher spray volumes, 5 gpa or more by air and 20 gpa or more by ground. |
| South Texas mixed brush, including, acacia, blackbrush acacia, catclaw acacia, twisted granjeno guajillo mesquite prickly pear tasajillo | 2 pts. Alligare Picloram 22K + 2/3-1 1/3 pt. Reclaim™ or 1 to 2 pts. Remedy™ | 2 qts. Alligare Picloram 22K + 2-3 pts. Remedy™ or 1-2 qts. Reclaim™ | Apply in oil-water emulsion. Use 4 or more gpa by air or 20 or more gpa by ground. For application timing for mesquite, see comments in section on mesquite control. Tank mixing with Reclaim™ will provide improved control of pricklypear and legume species such as mesquite and acacias while tank mixing with Remedy™ will provide improved control of non-legume species such as granjeno, oaks and hackberry. |
| mesquite | 1-2 pts. Alligare Picloram 22K + 2/3-1 1/3 pt. Reclaim™ or 2 pts. Alligare Picloram 22K + 1 pt. Remedy™ | 1-2 qts. Alligare Picloram 22K + 1-2 qts. Reclaim™ or 1 1/2-3 pts. Remedy™ | Alligare Picloram 22K Alone: Apply as a water spray or oil-emulsion (see Mixing Instructions) in 4 or more gpa by air or 10 or more gpa by ground. Increase spray volumes with increasing brush density and height to ensure adequate coverage. Where control of pricklypear cactus is desired, use the 2 pint/acre rate of Alligare |

2147

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|---|--|----------------------------------|-------------------------|
| | | | Picloram 22K. |
| <p>Alligare Picloram 22K in Tank Mix: Tank mixing with Reclaim™ will provide control of pricklypear and improved control of legume species such as mesquite and acacias while tank mixing with Remedy™ will provide improved control of non-legume species such as granjeno, oaks and hackberry. Regrowth mesquite should be at least 4 ft. tall prior to treatment. See labels for Reclaim™ and Remedy™ for additional treatment recommendations and information on mesquite control. Within rate ranges given for Alligare Picloram 22K and tank mix products, consult local recommendations.</p> <p>Timing and Factors in Control: The herbicidal response of mesquite is strongly influenced by environmental conditions as well as foliage condition and stage of growth. For best results, apply when new growth foliage has turned from light to dark green, when the soil temperature has reached 75°F to 83°F at a depth of 12-18 inches, and soil moisture is adequate for plant growth. Application should be made within 45 days after the critical soil temperature at the 12-18 inch depth has been reached, or if Alligare Picloram 22K is applied in combination with Reclaim™ within 60 days. Product performance may be adversely affected if application is made before mesquite foliage has turned from light to dark green or if foliage has been injured or removed by late frost, insects, hail or plant diseases. Do not apply if mesquite exhibits new (light green) growth in response to significant rainfall during the growing season. Soil temperatures at the 12-18 inch depth may vary with soil texture and drainage. Coarse-textured (sandy) soils warm up sooner than fine-textured soils (clay) soils and dry soils warm up more quickly than wet soils.</p> <p>Re-application: Do not reapply in the same growing season. Retreatment will not be effective until woody plants develop sufficient new foliage for interception, uptake, and translocation of the herbicide to plant roots.</p> | | | |
| ashe juniper eastern redcedar eastern persimmon | <p>General: Apply Alligare Picloram 22K undiluted as a spot concentrate application prior to periods of expected rainfall. Apply directly to the soil within the dripline and on the upslope side of the tree. Application to trees taller than 12 feet is not recommended. See directions for "Soil Spot Concentrate" in "Application Methods" section.</p> <p>Ashe Juniper: Apply 4 to 6 ml per 3 ft. of plant height in the spring (April-May).</p> <p>Eastern Redcedar: Apply 3 to 4 ml per 3 ft. of plant height in either spring (April-May) or fall (September-October).</p> <p>Eastern Persimmon: Apply 2 to 4 ml per inch of stem diameter in spring (March through May).</p> | | |

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River – Carpet Roller Use Recommendations)

| Brush Species | Amount of Alligare Picloram 22K | Specific Use Directions |
|---|---|---|
| mesquite regrowth | 1 gal. alone or with 2 qts. Reclaim™ herbicide to make 8 gals. of spray mixture | Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply from May through August, but preferably in May and June, when moisture availability is sufficient to allow normal plant growth. |
| huisache/blackbrush | 1 gal. alone or with 2 to 3 qts. of Reclaim™ to make 8 gals. of spray mixture | Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply in the fall. |
| <p>Refer to the Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: Texas, Oklahoma, and New Mexico) table for information on timing and factors in control of mesquite.</p> | | |

22/47

SPECIFIC USE DIRECTIONS FOR SMALL GRAINS, FALLOW CROPLAND, AND CRP ACRES

Spring Seeded Barley, Oats, and Wheat Not Underseeded With a Legume (Which is Not Flood or Sub-Irrigated and Not Rotated to Broadleaf Crops)

Do not apply to winter wheat or barley as crop injury may occur. Do not treat durum wheat since some varieties of durum wheat may be injured.

Use Alligare Picloram 22K for the control of susceptible annual broadleaf weeds such as (but not limited to) volunteer sunflower, wild buckwheat, lambsquarters, pigweed, Russian thistle, and sowthistle. This product may cause shorter straw on some varieties of cereals but grain yields are usually not affected.

Use Restrictions

- Do not apply Alligare Picloram 22K within 50 days before harvest.
- Do not graze or feed forage from treated areas for 2 weeks after treatment. Do not harvest hay from treated grain fields.
- Use only on land that will be planted the following year to grass, barley, oats, wheat, grain sorghum or fallowed. Do not plant grain sorghum within 8 months after application. This product is not intended for use on land planted to sweet sorghum. Do not apply more than 1 ½ fluid ounces of Alligare Picloram 22K per acre during the small grain growing season.

Use Alligare Picloram 22K for the control of susceptible annual broadleaf weeds such as (but not limited to) volunteer sunflower, wild buckwheat, lambsquarters, pigweed, Russian thistle, and sowthistle.

Broadcast Treatment (Ground and Aerial Applications)

Alligare Picloram 22K can be applied as a single broadcast treatment by ground or aerially to control several broadleaf weeds by itself or as a tank mix with 2,4-D, MCPA, or sulfonyleurea herbicides such as Ally®. Apply Alligare Picloram 22K at the rates suggested in the following table in 1 to 5 gallons of water per acre by air or in 5 to 20 gallons of water per acre by ground. The addition of surfactants may aid control under dry conditions, but may cause injury to grain if used over the top. Read and follow directions and precautions on other product labels when tank mixing.

Application Timing

Spring Wheat, Barley and Oats: Apply from the 3, to 5 leaf stage to early jointing stage of growth as indicated in the table below. Applications at the 3 to 5 leaf stage occasionally cause slight head malformations and straw shortening but normally do not affect yield.

Use Rates for Spring Wheat, Barley and Oats¹

| Weed Species | Weed Growth Stage ² | Grain Growth Stage | Amounts of Each Product Per Acre ³ | | |
|--|--------------------------------|--------------------------------|---|------------------------------|------------------------------|
| | | | Alligare Picloram 22K | 4 lb. a.e./gal 2,4-D or MCPA | 6 lb. a.e./gal 2,4-D or MCPA |
| More susceptible species, such as: lambsquarters pennycress wild mustard mayweed | 3 inches | 3 to 5 leaf to early tillering | 1 fl. oz. | ½ pint | 1/3 pint |
| | 3 to 6 inches | Tillering to early jointing | 1 ½ fl. oz. | ¾ pint | ½ pint |

23/47

| Weed Species | Weed Growth Stage ² | Grain Growth Stage | Amounts of Each Product Per Acre ³ | | |
|---|--------------------------------|-----------------------------|---|------------------------------|------------------------------|
| | | | Alligare Picloram 22K | 4 lb. a.e./gal 2,4-D or MCPA | 6 lb. a.e./gal 2,4-D or MCPA |
| Less susceptible species such as: volunteer sunflower wild buckwheat Russian thistle pigweed Canada thistle, top growth suppression | 1 to 6 inches | Tillering to early jointing | 1 ½ fl. oz. | ¾ to 1 pint | ½ to 2/3 pint |

¹For oats, do not tank mix with 2,4-D herbicides.

²For best results, treat when weeds have 2 to 4 leaves, and are actively growing.

³When measuring small amounts of Alligare Picloram 22K, special care should be taken not to exceed suggested rates.

Fallow Cropland (Not Rotated to Broadleaf Crops)

Apply Alligare Picloram 22K as a post harvest or fallow treatment in continuous grain or during the fallow period. Alligare Picloram 22K may be applied alone or in tank mix combination with 2,4-D or other herbicides registered for this use. Apply in 1 or more gallons of water per acre by air or 5 or more gallons per acre by ground.

Application Rates

Annual Weeds: To control annual weeds such as Russian thistle and wild buckwheat, apply ¼ to ½ pint per acre of Alligare Picloram 22K in tank mix combination with ½ to 1 lb. a.e. of 2,4-D or other herbicides registered for use on fallow land. Apply when weeds are actively growing.

Field Bindweed: Apply ½ to 1 pint per acre of Alligare Picloram 22K plus ½ lb. to 1 lb. a.e. per acre of 2,4-D when bindweed is actively growing. Optimum time for treatment is when plant runners reach 8 to 12 inches. Use ½ pint per acre to control light to moderate infestations under good growing conditions or to reduce the potential for crop injury. Use higher rates for heavy infestations and longer term control. Some regrowth will occur the following season and a re-treatment program for successive years is recommended.

Canada thistle: Apply 1 pint per acre of Alligare Picloram 22K plus 1 lb. a.e. per acre of 2,4-D when the majority of thistle plants are emerged but prior to bud stage.

Crop Rotation

Use only on land to be planted the following year to grass, barley, oats, wheat or fallow land. Many broadleaf crops are extremely sensitive to soil residues of Alligare Picloram 22K. Do not plant sensitive broadleaf crops for 36 months after treatment or until soil residues have declined to a safe level as indicated by an adequately sensitive bioassay using the intended broadleaf crop. A bioassay is recommended following treatment prior to planting any sensitive broadleaf crop.

Preplant Interval

A preplant interval following application of Alligare Picloram 22K prior to planting small grains is recommended to reduce or eliminate potential crop injury and/or yield reduction. The possibility for crop injury or yield reduction to occur depends on application rate, soil organic matter, rainfall, temperature and incidence of cereal diseases. Adequate soil moisture and soil temperature during the preplant interval is important in reducing, but may not eliminate, the risk of crop injury. When considering use of Alligare Picloram 22K on fallow land, growers should consider the benefit of weed control against the risk of crop damage and treat only if the risk of injury to small grains can be tolerated. The following preplant intervals are recommended:



For applications up to ½ pint per acre, allow a minimum of 45 days of soil temperatures above 40°F between application and planting.

24/47

For applications of greater than ½ pint and up to 1 pint per acre, allow a minimum of 60 days of soil temperatures above 40°F between application and planting, except in the states of Idaho, North Dakota, Nebraska, Montana, Oregon, South Dakota, Washington and Wyoming, where the minimum preplant interval is 90 days.

Restrictions:

- Do not apply more than 1 pint per acre as a broadcast treatment per annual growing season.
- Spot Treatment: See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing. Spot treatments of Alligare Picloram 22K at rates over 1 pint per acre can be made on fallow, non-irrigated cropland if the treated areas comprise less than 10% of the immediate field in any one year. Alligare Picloram 22K should not be applied to cropland at rates exceeding 2 quarts per acre. When Alligare Picloram 22K is applied at rates above 1 pint per acre, injury to small grains may result for periods up to two years after treatment.

Conservation Reserve Program (CRP) for Seeding to Permanent Grasses Only

Do not use Alligare Picloram 22K if legumes are a desired cover during CRP.

Conditions that stress grasses, such as drought, will increase potential for injury to the grass at all stages of growth.

To reduce potential damage to subsequent small grain crops, use the lower rate or discontinue the use of Alligare Picloram 22K at least 2 years prior to the seeding of small grain crops. After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay shows that no detectable picloram is present in the soil.

Broadcast Treatment (Ground and Aerial Applications)

Applications of Alligare Picloram 22K should be made after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses show improved tolerance to the herbicide at this stage of development.

For control of actively growing perennial weeds, use up to 2 pints per acre of Alligare Picloram 22K after the grass is established. For best results, use in 1 or more gallons of water per acre by air or in 5 or more gallons of water per acre by ground. Increasing the rate of application can increase the risk of injury.

For control of actively growing susceptible annual broadleaf weeds, (including Russian thistle) apply ¼ to ½ pint per acre of Alligare Picloram 22K. Alligare Picloram 22K can also be tank mixed with ½ to 1 pound per acre of 2,4-D where 2,4-D-sensitive species are present. Read and follow all directions for use and use precautions on other product labels.

Spot Treatment

See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing. For spot applications when perennial grasses are established, use 1 to 4 pints per acre of Alligare Picloram 22K. Rates of 2 pints per acre or more should only be used for control of deep-rooted perennial broadleaf weeds.

Precaution: Alligare Picloram 22K at rates over 2 pints per acre may suppress certain established grasses such as brome grass, bluegramma and buffalograss. However, subsequent grass growth should be improved by release from weed competition.

25/47

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
PESTICIDE STORAGE: The active ingredient in this product may crystallize and settle out of solution if product is exposed to subfreezing temperatures. Under these conditions, warm product to at least 40°F and agitate well to dissolve any crystallized material prior to use. Open dumping is prohibited.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Deleted: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Every 2 years starting January 1, 2008, the registrant will offer training to applicators which will cover application techniques and product stewardship particular to their use(s) of this product (Alligare Picloram 22K, Registration No. 81927-18). Applicators of this product must be able to provide certification of such training on demand to the State, Tribal or Federal enforcement agent.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Reclaim™ and Remedy™ are trademarks of Dow AgroSciences LLC.
Ally® and Escort® are registered trademarks of E.I. duPont de Nemours & Co., Inc.

EPA 20080320

26/47

RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible, non-target plants. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

Alligare Picloram 22K

For use in areas west of the Mississippi River for the control of susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, spring seeded wheat, barley and oats not underseeded with a legume, non-cropland, and on Conservation Reserve Program (CRP) acres.

ACTIVE INGREDIENT:

Picloram: 4-amino-3,5,6-trichloropicolinic acid, potassium salt24.4%

OTHER INGREDIENTS:75.6%

TOTAL:100.0%

Alligare Picloram 22K contains the following acid equivalent:

Picloram: 4-amino-3,5,6-trichloropicolinic acid – 21.1% (2 lbs./gal.)

**KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

| | |
|-------------|---|
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
|-------------|---|

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals.

EPA Reg. No. 81927-18

EPA Est. No.

Manufactured For:

Alligare, LLC
13 N. 8th Street
Opelika, AL 36801

Net Contents:

27/47

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves Category A, such as barrier laminate ≥ 14 mils, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to some plants at very low concentrations. Non-target plants may be adversely affected if pesticide is allowed to drift from areas of application. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of wastes or rinsate. Do not allow runoff or spray to contaminate wells, irrigation ditches or any body of water used for irrigation or domestic purposes. Do not make application when circumstances favor movement from treatment site.

This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, picloram may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetative filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not intended for manufacturing or formulating.

28/47

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves Category A, such as barrier laminate ≥ 14 mils, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: For applications on rangeland, permanent grass pastures, and non-cropland, do not enter or allow worker entry into treated areas until sprays have dried, unless applicator and other handler PPE is worn.

GENERAL INFORMATION

In areas west of the Mississippi River use Alligare Picloram 22K to control susceptible broadleaf weeds and woody plants on rangeland and permanent grass pastures, fallow cropland, spring seeded wheat, barley and oats not underseeded with a legume, non-cropland, and on Conservation Reserve Program (CRP) acres. This product is NOT for sale or use in the San Luis Valley of Colorado.

PRECAUTIONS AND RESTRICTIONS

Use this product only as specified on this label.

Observe any special use and application restrictions and limitations, including method of application and permissible areas of use as promulgated by state or local authorities.

Do not mix with dry fertilizer.

Chemigation: Do not apply this product through any type of irrigation system.

Grazing Poisonous Plants: Application of this herbicide may increase the palatability of certain poisonous plants. Do not graze treated areas until poisonous plants are dry and no longer palatable to livestock.

29/47

Maximum Use Rates:

Non-cropland Areas: Total use of picloram, including retreatments or spot treatments, must not exceed 1 lb ae per acre picloram (2 quarts per acre Alligare Picloram 22K) per annual growing season.

Rangeland and Permanent Grass Pastures: For control of noxious weeds as defined by federal, state, or local authorities, picloram may be applied at up to 1 lb ae per acre (2 quarts Alligare Picloram 22K per acre) per annual growing season as a broadcast treatment. Spot treatments may be applied at the equivalent broadcast rate of up to 1 lb ae picloram (2 quarts Alligare Picloram 22K) per acre.

For control of other broadleaf weeds and woody plants, picloram may be applied broadcast at up to 0.5 lb ae per acre (1 quart Alligare Picloram 22K per acre) per annual growing season. Apply spot treatments at an equivalent broadcast rate of up to 1 lb ae per acre (2 quarts Alligare Picloram 22K per acre) per annual growing season, but not more than 50% of an acre may be treated. Repeat treatments may be applied as necessary, but total use must not exceed the maximum amount specified.

Cropland (Spring-Seeded Wheat, Barley and Oats): Do not apply more than 0.09 lb ae picloram (1 1/2 fluid ounces of Alligare Picloram 22K) per acre during the small grain growing season.

Fallow Cropland (Not Rotated to Broadleaf Crops): Do not apply more than 1 lb ae picloram per acre (1 pint per acre of Alligare Picloram 22K) as a broadcast treatment per annual growing season.

Conservation Reserve Program (CRP) for Seeding to Permanent Grasses Only: Do not broadcast apply more than 0.5 lb ae per acre of picloram (1 quart per acre of Alligare Picloram 22K) per annual growing season or apply more than 1 lb ae per acre picloram (2 quarts per acre of Alligare Picloram 22K) per annual growing season as a spot application. To reduce potential damage to subsequent small grain crops, use the lower rate or discontinue the use of Alligare Picloram 22K at least 2 years prior to the seeding of small grain crops. After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay shows that no detectable picloram is present in the soil.

Precautions for Avoiding Injury to Non-Target Plants

- Do not move treated soil to other areas or use it to grow plants if they are not registered for use with picloram until an adequate sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.
- Do not spray if the loss of forage legumes, including clover cannot be tolerated. Alligare Picloram 22K may injure or kill legumes. New legume seedlings may not grow for several years following application of this product.
- Do not make application when circumstances favor movement from treatment site.
- Do not rotate to food or feed crops on treated land if they are not registered for use with picloram until an adequately sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.
- Do not contaminate water intended for irrigation or domestic purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or runoff to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes.
- Do not apply Alligare Picloram 22K on residential or commercial lawns or near ornamental trees and shrubs. Untreated trees can occasionally be affected by root uptake of herbicide through movement into the topsoil or by excretion of the product from the roots of nearby treated trees. Do not apply Alligare Picloram 22K within the root zone of desirable trees unless such injury can be tolerated.
- Allow 7 days of grazing on an untreated grass pasture (or feeding of untreated hay) before transferring livestock from treated grazing areas (or feeding of treated hay) onto sensitive broadleaf crop areas. Otherwise, urine and manure may contain enough picloram to cause injury to sensitive broadleaf plants.
- Do not use manure from animals grazing treated areas or feeding on treated hay on land used for growing broadleaf crops, ornamentals, orchards or other susceptible, desirable plants. Manure may contain enough picloram to cause injury to susceptible plants.

30/47

- Do not use grass or hay from treated areas for composting or mulching of susceptible broadleaf plants.
- Do not apply to snow or frozen ground. Application during very cold (near freezing) weather is not advisable.
- Do not use on flood or sub-irrigated land.

Sprayer Clean-Out: To avoid injury to desirable plants, equipment used to apply Alligare Picloram 22K should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
2. Rinse a second time, adding 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15-20 min.). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed and cleaned separately.

Precautions for Avoiding Injurious Spray Drift

For aerial applications on rights-of-way or other areas near susceptible crops, use drift control additive as recommended by the manufacturer.

Do not apply or otherwise permit Alligare Picloram 22K or spray containing Alligare Picloram 22K to contact crops or other desirable broadleaf plants, including but not limited to alfalfa, beans, grapes, melons, peas, potatoes, safflower, soybeans, sugar beets, sunflower, tomatoes, and other vegetable crops, flowers, fruit plants, ornamentals, or shade trees or the soil containing roots of nearby valuable plants.

Applications should be made to avoid spray drift since very small quantities of spray, which may not be visible, may seriously injure susceptible crops during both growing and dormant periods. To minimize spray drift, use low nozzle pressure; apply as a coarse spray; and use nozzles designed for herbicide application that do not produce a fine droplet spray. To aid in further reducing spray drift, a drift control and deposition aid may be used with this product, especially when water alone is used as the carrier. If a drift control aid is used, follow all use recommendations and precautions on the product label. Do not use a thickening agent with Microfoil or Thru-Valve booms, or other systems that cannot accommodate thick sprays.

Ground Equipment: With ground equipment, spray drift can be reduced by keeping the spray boom as low as possible; by applying 10 gallons or more of spray per acre; by keeping the operating spray pressures at manufacturer's recommended minimum pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); by spraying when wind velocity is low (follow state regulations). Avoid calm conditions which may be conducive to air inversions. Direct sprays no higher than the tops of target vegetation and keep spray pressures low enough to provide coarse spray droplets to minimize drift. A spray thickening agent may be used to further reduce the potential for drift.

Aerial Application:

AERIAL SPRAY DRIFT MANAGEMENT

Spray Drift Management

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

31/47

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations:

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Aerial Drift Reduction Advisory

[This section is advisory in nature and does not supersede the mandatory label requirements.]

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2

32/47

mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

MIXING INSTRUCTIONS

Mix the required amount of Alligare Picloram 22K in water and apply as a coarse, low pressure spray using ground equipment or aircraft. Use enough spray volume to provide uniform coverage of the weeds.

Use with Surfactants: Addition of wetting or penetration agents is not usually necessary when using Alligare Picloram 22K. However, under certain conditions, such as drought, addition of a surfactant may improve efficacy. However, if foliar burn occurs too rapidly, translocation of Alligare Picloram 22K will not occur and control of perennial weeds, such as field bindweed, may be reduced.

Mixing with Water

To prepare the spray, add about half the desired amount of water in the spray tank. Then with agitation, add the recommended amount of Alligare Picloram 22K and other registered tank mix herbicides. Finally, with continued agitation, add the rest of the water and additives such as surfactants or drift control and deposition aids.

Mixing Oil-Water Emulsions (Ground and Aerial Applications)

For aerial application, add oil to the total spray mix at the ratio of 1 part oil to 5 parts water (1:5 ratio). For ground application, add oil to the spray mix at a rate of 5 to 10% of the total mix. Do not use more than 1 gallon of oil per acre for aerial or ground application. Use agricultural spray emulsifiers such as Sponto 712 or Triton X-100 according to mixing instructions given below.

Batch Mixing Instructions

With continuous, vigorous agitation:

1. Add half the amount of water to be used to the spray tank.
2. Add the required amount of water-soluble herbicides such as Alligare Picloram 22K, Reclaim™ herbicide or 2,4-D Amine.
3. With continued, vigorous agitation, slowly add a premix of oil, emulsifier and oil soluble herbicides such as Remedy™ herbicide or a 2,4-D ester as required. Note: Do not add water or mixtures containing water to the premix or oil soluble herbicide since a thick "invert" (water in oil) emulsion may

be formed that will be difficult to break. An invert emulsion will also form if the premix is added to the mixing tank before the addition of water.

- 4. Finish filling the spray tank and maintain sufficient agitation to ensure uniformity of the spray mixture during application.

Mixing with Sprayable Liquid Fertilizer Solutions

Alligare Picloram 22K is compatible with most non-pressurized liquid fertilizer solutions; however, a compatibility test (jar test) should be made prior to mixing. Jar tests are particularly important when a new batch of fertilizer or pesticide is used, when water sources change, or when tank mixture ingredients or concentrations are changed. Compatibility may be determined by mixing the spray components in the desired order and proportions in a clear glass jar before large scale mixing of spray components in the spray tank.

Note: The lower the temperature of the liquid fertilizer, the greater the likelihood of mixing problems. Use of a compatibility aid such as Unite or Compex may help obtain and maintain a uniform spray solution during mixing and application. Compatibility is best with straight liquid nitrogen fertilizer solutions. Mixing with N-P-K fertilizer solutions or suspensions is more difficult and should not be attempted without first conducting a successful jar test. Agitation in the spray tank must be vigorous to be comparable with jar test agitation. For best results, liquid fertilizer rates should not exceed 50% of the total spray volume. Premix Alligare Picloram 22K with water and add to the liquid fertilizer/water mixture while agitating contents of the spray tank. Apply the spray the same day it is prepared while maintaining continuous agitation. Rinse spray tank thoroughly after use.

Note: Foliar applied liquid fertilizers used as carrier for Alligare Picloram 22K can cause yellowing or leaf burn of crop foliage.

Do not use spray equipment used to apply Alligare Picloram 22K for other applications to land planted to, or to be planted to susceptible crops or desirable sensitive plants, unless it has been determined that all phytotoxic residue of this herbicide has been removed by thorough cleaning of equipment.

Local conditions may affect the use of herbicides. State agricultural experiment stations or extension service weed specialists in many states issue recommendations to fit local conditions. Be sure that use of this product conforms to all applicable regulations.

Tank Mixing

Alligare Picloram 22K may be applied in tank mix combination with labeled rates of 2,4-D or other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not exceed recommended application rates. If products containing the same active ingredient are tank mixed, do not exceed the maximum allowable active ingredient use rates.
- For products packaged in water-soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See "Sprayer Clean-Out" under "Precautions for Avoiding Injury to Non-Target Plants".)
- For direct injection or other spray equipment where the product formulations will be mixed in undiluted form, special care should be taken to ensure tank mix compatibility.
- Always perform a (jar) test to ensure the compatibility of products to be used in tank mixture.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Alligare Picloram 22K and other pesticides or carriers. Use a clear glass jar with lid and mix the tank mix ingredients in their relative proportions. The tank mixture is compatible if the materials mix readily when the jar is inverted several times. The mixture should remain stable after standing for 1/2 hour or, if

34/47

separation occurs, should readily mix if agitated. An incompatible mixture is indicated by separation into distinct layers which do not readily remix when agitated and/or the presence of flakes, precipitates, gels, or heavy oily film on the jar.

APPLICATION METHODS

Ground or Aerial Broadcast

Use Alligare Picloram 22K as a broadcast treatment by ground or aerially to control listed broadleaf weeds and woody plants. Apply Alligare Picloram 22K as a coarse low-pressure spray at the recommended rates in a spray volume of 1 or more gallons per acre by air or 10 or more gallons per acre by ground. Aerial applications can be applied using aerial electrostatic spray charging systems.

High-Volume Foliar Applications

Spray to thoroughly wet foliage and stems. An approved agricultural surfactant may be added at the manufacturer's recommended rate. Do not apply more than the maximum application rate of Alligare Picloram 22K specified for a given treatment site.

Spot Treatment

Use application rates as suggested in the "Approved Uses" section of this label or recommended by your area weed control specialist. Apply in a total spray volume of 20 to 100 gallons per acre. To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to directions provided below. Do not exceed maximum application rates for Alligare Picloram 22K for a given treatment site. On rangeland and permanent grass pastures, spot treatments may be applied at an equivalent broadcast rate of up to 2 quarts per acre per annual growing season, but not more than 50% of an acre may be treated. Repeat treatments may be applied as necessary, but total use must not exceed the maximum amount specified.

Hand-Held Sprayers: Hand-held or backpack sprayers may be used for spot applications of Alligare Picloram 22K if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on an area of 1,000 sq. ft. Mix the amount of Alligare Picloram 22K (fl. oz. or ml) corresponding to the desired broadcast rate in 0.5 to 2.5 gallons of water, depending on the spray volume required to treat 1,000 sq. ft. To calculate the amount of Alligare Picloram 22K required for larger areas, multiply the table value (fl. oz. or ml) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3,500 sq. ft., multiply the table value of 3.5 (calc. 3,500/1,000 = 3.5). An area of 1,000 sq. ft. is approximately 10.5 x 10.5 yards (strides) in size.

| Amount of Alligare Picloram 22K per 1,000 sq. ft. to Equal Specified Broadcast Rate | | | | | |
|---|--------------------------------|--------------------------------|------------------------------|-------------------------------|-----------------------------|
| ¼ pt./acre | 1/3 pt./acre | ½ pt./acre | 2/3 pt./acre | 1 pt./acre | 1 qt./acre |
| <u>1/10 fl. oz.</u> (2.7 ml) | <u>1/8 fl. oz.</u> (3.6 ml) | <u>1/5 fl. oz.</u> (5.4 ml) | <u>¼ fl. oz.</u> (7.3 ml) | <u>3/8 fl. oz.</u> (11 ml) | <u>¾ fl. oz.</u> (22 ml) |

¹ 1 fl. oz. = 29.6 (30) ml

Special Application Methods

Wick Application – Mix 1 part of Alligare Picloram 22K with 2 parts of water to prepare a 33% solution. Apply when weeds are actively growing and are above most desirable plants. For ironweed and goldenrod, best results are obtained when applications are made prior to the early bud stage. Drain the wick applicator and clean after each use. Change ropes when flow is reduced from wear, extended use, poor cleaning or intermittent use.

Carpet Roller Application – Alligare Picloram 22K can be applied with carpeted rollers where drift presents a hazard to susceptible crops, surface waters and other sensitive areas. Apply to previously untreated plants less than 6 feet tall, and short enough to pass beneath the tractor without breaking off at the ground. Applications made during periods of extended drought conditions will not provide acceptable control. Do not burn, mow or otherwise disturb the treated plants during the remainder of the growing season. Operate carpeted rollers as close to the ground as possible without breaking the stems, but

35/47

above the tallest grasses. Grasses growing adjacent to treated plants may exhibit temporary injury. Maximize herbicide deposition on stems and foliage and minimize drippage losses by rotating the carpeted roller at 30 to 40 rpm with the lower edge moving in the same direction as the direction of travel. Maintain the carpet sufficiently wetted to apply up to 1 gal./acre of herbicide-water mixture to stands of average density (100 to 200 plants/acre), and up to 2 gals./acre in dense stands (300 to 400 plants/acre). Rewet rollers at regular intervals. See the Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River – Carpet Roller Use Recommendations) section of this label for treatment recommendations.

Soil Spot Concentrate: Alligare Picloram 22K may be applied undiluted as a spot concentrate application to control ashe juniper, eastern redcedar and eastern persimmon (see specific user directions for these plant species which follow). Applications should precede periods of expected rainfall. Apply directly to the soil within the dripline and on the upslope side of the tree. Applications to trees taller than 12 feet are not recommended.

Special Ground Sprayer Equipment: To control annual and perennial weed species using special low-volume, minimum drift equipment, such as the hooded Forage Chemical Mower, apply 1 to 2 pts. of Alligare Picloram 22K in total volumes ranging from 1 gallon to 5 gallons per acre in water alone or as an oil-water emulsion at a 1:5 and 1:4 oil-to-water ratio for a 1 gallon and 5 gallon per acre solutions, respectively.

WOODY PLANTS AND BROADLEAF WEEDS CONTROLLED BY ALLIGARE PICLORAM 22K

Woody Plants:

| | |
|--------------------|-----------------------------|
| acacia, blackbrush | huisache (suppression only) |
| acacia, catclaw | junipers/cedars |
| acacia, twisted | locust |
| aspen | mesquite |
| broom, Scotch | pine, pinyon |
| camelthorn | rabbitbrush, Douglas |
| chaparral sp. | rose, Macartney |
| gorse | rose, multiflora |
| granjeno | sagebrush, fringed |
| guajillo | tallowtree, Chinese |
| | wormwood, absinth |

Annual and Perennial Broadleaf Weeds:

| | | |
|---------------------------------|--------------------------------|-------------------------------|
| bindweed, field (p) | ironweed (p) | ragwort, tansy (b) |
| bitterweed (a) | knapweed, diffuse (a) | Russian thistle (a) |
| broomweed, annual (a) | knapweed, Russian (p) | skeletonweed, rush (p) |
| buckwheat, wild (a) | knapweed, spotted (p) | smartweed (a) |
| buffalobur (a) | knapweed, squarrose (p) | snakeweed, broom (p) |
| bullnettle (p) | lambquarters (a) | sneezeweed, bitter (a) |
| bursage (a) | larkspur, geyer (p) | sowthistle, perennial (p) |
| burweed (p) | larkspur, plains (p) | spurge, leafy (p) |
| camphorweed (a) | larkspur, tall (p) | St. Johnswort (p) |
| carrot, wild (b) | lettuce, prickly (a) | starthistle, Iberian (a) |
| cinquefoil, sulfur (p) | licorice, wild (p) | starthistle, purple (a) |
| cocklebur (a) | locoweeds (p) | starthistle, yellow (a) |
| coneflower, upright prairie (p) | loco, woolly (p) | sunflower (a) |
| croton (a) | loco, Wooten (garbancillo) (p) | tasajillo (p) |
| crupina, common (a) | lupines (p) | thistles, annual or biennial, |
| daisy, ox-eye (p) | marshelder (sumpweed) (a) | including: |

36/47

dock, curly (p)
 garbancillo (Wooten loco) (p)
 goldaster, gray (p)
 goldaster, narrowleaf (p)
 goldenrod, common (p)
 goldenweed, Drummond (p)
 groundsel (p)
 henbane, black (a,b)
 horsenettle, Carolina (p)
 horsenettle, western (p)
 horsenettle, white (p)
 horseweed (a)
 (a) – annual; (b) – biennial; (p) – perennial

mayweed (a)
 milkweed (p)
 mustard, wild (a)
 nightshade, silverleaf (p)
 pennycress (a)
 pigweed (a)
 pricklypear, plains (p)
 pricklypear, lindheimer (p)
 ragweed, bur (a)
 ragweed, common (a)
 ragweed, lanceleaf (a)
 ragweed, western (a)

thistle, bull (b)
 thistle, distaff (a)
 thistle, Italian (b)
 thistle, musk (b)
 thistle, plumeless (b)
 thistle, Scotch (b)
 thistles, perennial, including
 thistle, Canada (p)
 thistle, wavy leaf (p)
 toadflax, dalmation (p)
 toadflax, yellow (p)
 yankeeweed (p)

SPECIFIC USE DIRECTIONS FOR RANGELAND, PERMANENT GRASS PASTURES AND NON-CROPLAND

General Requirements for Non-Cropland Areas

Use Alligare Picloram 22K to control susceptible broadleaf weeds and woody plants on non-cropland areas such as roadsides or other rights-of-way, fence rows, and around farm buildings. Up to 2 quarts of Alligare Picloram 22K per acre may be applied. See "Rangeland and Permanent Grass Pastures" section for specific target weed or woody plant species treatment recommendations.

Maximum Use Rates for Non-Cropland Areas: Total use of Alligare Picloram 22K, including retreatments or spot treatments must not exceed 2 quarts per acre per annual growing season.

General Requirements for Rangeland and Permanent Grass Pastures for States West of the Mississippi River

Use Alligare Picloram 22K on rangeland and permanent grass pastures to control susceptible broadleaf weeds and woody plants including, but not limited to those shown in the following tables. Many annual weeds at the seedling stage are controlled at the rate of 1 pt. per acre. Where a rate range is recommended, choose the higher rate for dense weed infestations, and for more dependable, longer-lasting control. Lower rates will perform best when applied under favorable conditions and at the optimum growth stage, but may provide a lower level of control and require retreatment. For best results, treat when weeds are small and actively growing in the spring before full bloom, however, certain weeds may also be treated in late summer or fall. Treatments during full bloom or seed stage of some weeds may not provide acceptable control.

Refer to the "Application Methods" section of this label for information on various methods of application including ground or aerial broadcast, high volume foliar application, spot treatments, and special application methods for certain weeds or woody plants including spot concentrate application or application with special low volume or hooded spray equipment.

Precautions and Restrictions:

Maximum Use Rates for Rangeland and Permanent Grass Pastures: For control of noxious weeds as defined by federal, state, or local authorities, Alligare Picloram 22K may be applied at up to 2 quarts per acre per annual growing season as a broadcast treatment. Spot treatments may be applied at the equivalent broadcast rate of up to 2 quarts per acre.

For control of other broadleaf weeds and woody plants, Alligare Picloram 22K may be applied broadcast (ground, air, or high volume foliar) at up to 1 quart per acre per annual growing season. Spot treatments (hand sprayer, calibrated boom, high volume foliar, or soil spot concentrate) may be applied at an equivalent broadcast rate of up to 2 quarts per acre per annual growing season, but not more than 50% of

37/47

an acre may be treated. Repeat treatments may be applied as necessary, but total use, must not exceed the maximum amount specified.

Grazing Restrictions: Do not cut grass for feed within two weeks after treatment when applying more than 1 quart of Alligare Picloram 22K per acre. Meat animals grazing for up to two weeks after treatment should be removed from treated areas three days prior to slaughter. Do not graze lactating dairy animals on treated areas within two weeks after treatment.

Certain established grasses, such as bromegrass, blue gamma, and buffalograss may be suppressed when using rates of Alligare Picloram 22K over 1 quart per acre. However, subsequent grass growth should be improved by release from weed competition.

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River, Except Texas, Oklahoma and New Mexico)

Apply Alligare Picloram 22K broadcast or as a spot application unless otherwise specified (See "Application Methods" section of this label). For additional species or more specific rates, consult your area's current Weed Control Guide and/or your local Alligare, LLC representative. Rates in the following table are advisory in nature and do not supersede the maximum rates for the use site allowed for different methods of application.

Weed Control Prior to Seeding Perennial Grasses:

Weed control with Alligare Picloram 22K fits into the following grass re-vegetation programs. Alligare Picloram 22K may be applied in the spring or early summer, depending on the target weed species and grass seed planted in the fall when conditions are favorable. Alternatively, Alligare Picloram 22K may be applied in the fall and grass seed planted in the winter or fall when conditions are favorable.

Apply Alligare Picloram 22K at 1 qt./acre or less; see rate chart for information on control of target weed species. Depending on grass species sensitivity, there may be temporary injury on new plantings when Alligare Picloram 22K is applied at 1 qt./acre. However, this injury will be insignificant in comparison with the benefit to grasses due to the removal of weed competition. Germination of annual grass species may be suppressed after treatment. To optimize weed control, it is suggested the application area be disturbed as little as possible by the seeding operation. At the very least, the site should be left undisturbed for 14 days prior to seedbed preparation or seeding. To decrease the potential for injury on sensitive grass species, increase the interval between application and seeding.

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|--|--|---|
| Annual and Biennial Weeds: | | |
| bursage (bur ragweed) crupina, common henbane, black horseweed starthistle, Iberian starthistle, purple starthistle, yellow | 1-2 pts. Alligare Picloram 22K | Apply when there is adequate soil moisture and weeds are actively growing. |
| thistles, including, bull distaff Italian musk plumeless scotch | Fall: ½ - ¾ pt. Alligare Picloram 22K Spring: ½ - ¾ pt. Alligare Picloram 22K + 1 lb. a.e. 2,4-D | General: Apply at the rosette stage before bolting in the spring or in the fall prior to soil freeze up. Distaff Thistle: Apply at rosette stage in spring only. Bolted Musk Thistle: Apply before flowering at the rate of ½ - 1 pt. of Alligare Picloram 22K plus 1 lb. a.e. of 2,4-D/acre. |

38/47

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|--|--|--|
| Perennial Weeds: | | |
| pricklypear, plains | ½ - 1 pt. Alligare Picloram 22K | Apply at peak of flowering. Use of an oil-water emulsion spray mixture may improve control. Lower rate will provide partial control (stand reduction) and high rate more complete control. Treatment response is slow and may continue for 2 years or longer. |
| sagebrush, fringed | ½ - 1 pt. Alligare Picloram 22K 1 lb. a.e. 2,4-D ester | Apply after seed stalk elongation and early flowering and throughout the summer if growing conditions are favorable. |
| cinquefoil, sulfur larkspur, geyer larkspur, plains locoweeds snakeweed, broom | 1 pt. Alligare Picloram 22K | General: Apply when weeds are actively growing. Sulfur cinquefoil: Apply during active growth or fall regrowth. Geyer larkspur: Apply when plant is actively growing between rosette stage and flower bud formation. Locoweeds: Apply from early bud to early bloom stage. See "General Use Precautions" section of this label for note on grazing treated poisonous plants. Broom snakeweed: Apply during active growth between full leaf to early bloom stage. |
| burweed daisy, ox-eye goldenrod, common knapweed, diffuse knapweed, spotted knapweed, squarrose rabbitbrush, Douglas thistle, Canada thistle, wavy leaf wormwood, absinth | 1-2 pts. Alligare Picloram 22K | General: Lower rates in rate range may require annual spot treatments. Control with lower rates may be improved by tank mixing with 1.0 lb. a.e. per acre 2,4-D. Goldenrod: Apply during active growth prior to bud stage. Diffuse or spotted knapweed: Optimum application from rosette to mid-bolting stage or to fall regrowth. Thistle (Canada and Wavy Leaf): Apply when most basal leaves have emerged, but before bud stage, or apply to regrowth in the fall. Apply rates less than 1 ½ pts./acre only under favorable conditions and in combination with 1 lb. a.e./acre of 2,4-D; retreatment may be required. Absinth wormwood: Apply in spring or early summer when plants are actively growing. |
| licorice, wild milkweed | 2 pts. Alligare Picloram 22K | Wild Licorice: Apply at bloom stage. Milkweed: Treat during active growth and tank mix recommended rate of Alligare Picloram 22K with 1 lb. a.e./acre 2,4-D. |
| bindweed, field gorse lupines knapweed, Russian | 2-4 pts. Alligare Picloram 22K | General: Annual retreatment of these species will be required at rates at low end of rate range. Control at low end of rate range may be improved by tank mixing |

39/47

| Weed Species | Broadcast Application (Rate/acre) | Specific Use Directions |
|---|---|--|
| ragwort, tansy skeletonweed, rush spurge, leafy St. Johnswort toadflax, dalmation | | with 1 lb. a.e. 2,4-D. Russian Knapweed: Apply during active growth from bud to mid-flowering, or to fall regrowth. Leafy Spurge: Apply at true flower stage of growth or apply to fall regrowth. Re-apply when level of control falls below 80 percent. Dalmation Toadflax: Apply when plants are actively growing through full bloom stage of growth. |
| larkspur, tall sowthistle, perennial toadflax, yellow | 4 pts. Alligare Picloram 22K | General: A retreatment program may be necessary for satisfactory control of these species. Tall Larkspur: For best results apply from 6 inches tall to late bloom stage. For increased control, apply in tank-mix with Ally® or Escort® herbicide and non-ionic surfactant. See "General Use Precautions" section of this label for note on grazing treated poisonous plants. |
| Woody Plants: | | |
| juniper | 4 qts. Alligare Picloram 22K per 100 gallons of spray | Apply as a high volume foliar spray/individual plant treatment. |
| redcedar, eastern | Eastern red cedar can be controlled with spot concentrate applications of Alligare Picloram 22K in either the spring (April-May) or fall (September-October). For best results, use 3 ml to 4 ml of Alligare Picloram 22K (undiluted) per 3 feet of plant height. Application should precede periods of expected rainfall. Apply directly to soil within the dripline and on the upslope side of the tree. Application to trees taller than 15 feet is not recommended. Do not use more than 2 pints of Alligare Picloram 22K per acre in any one year. | |

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: Texas, Oklahoma and New Mexico)

Apply Alligare Picloram 22K as a broadcast treatment by ground or air, as a spot application, or high volume foliar application to control listed broadleaf weeds and woody plants, unless otherwise specified (See "Application Methods" section of this label). Refer to the "General Requirements for States West of the Mississippi River" for maximum use rates. To control certain species, Alligare Picloram 22K may be applied alone or in combination with 2,4-D as indicated in the following table. When Alligare Picloram 22K is applied alone, herbicide symptoms will be slower to appear than when it is applied in combination with 2,4-D.

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|--|--|-------------------------------------|---|
| Annual and Biennial Weeds: | | | |
| bitterweed, western broomweed, annual buffalobur | Early Season: ½ - 1 pt. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + | General: Apply when there is adequate soil moisture and weeds are actively growing. |

40/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|---|--|--|---|
| bursage (bur ragweed) camphorweed carrot, wild cocklebur croton horseweed lettuce, prickly marshelder (sumpweed, sulfaweed) ragweed, common ragweed, lanceleaf ragweed, western smartweed sneezeweed, bitter sunflower thistle, bull thistle, musk | + 0.5 lb. a.e. 2,4-D or ¾ - 1 ½ pts. Alligare Picloram 22K Mid to Late Season: 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1-2 pts. Alligare Picloram 22K | 2 lbs. a.e. 2,4-D | Early Season: Recommendations are intended only for very early in the season when weeds are no more than 2 to 3 inches tall. Mid to Late Season: Recommendations are for weeds from 3 inches tall to early flowering. Marshelder: Use a minimum of ¾ pt./acre of Alligare Picloram 22K plus 0.75 lb. a.e./acre 2,4-D + non-ionic surfactant. Apply when plants are no more than 3-6 inches tall. Control may be improved by adding ammonium sulfate at 17 lbs./100 gals. spray solution. Thistles: Apply when thistles are in the rosette stage before bolting. Lanceleaf Ragweed: Use the higher rate within the recommended rate range. |
| Perennial Weeds: | | | |
| snakeweed, broom | Fall, Early Winter 1 pt. Alligare Picloram 22K Late Winter, Early Spring 2 pts. Alligare Picloram 22K | - | Fall and Early Winter: If rainfall is less than average prior to flowering, apply after flowering is complete. If rainfall is average to above average prior to or during flowering, apply during full flower and/or active pollination, before resumption of new top growth. Late Winter and Early Spring: Apply following sufficient precipitation (rain or snow) to stimulate active plant growth. Both basal and terminal leaves should be green and active plant growth occurring. |
| bullnettle coneflower, upright prairie dock, curly horsenettle, Carolina horsenettle, western horsenettle, white ironweed nightshade, silverleaf yankeeweed | ½ - 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1-2 pts. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D | General: Apply when there is adequate soil moisture and weeds are actively growing. Nettles and Silverleaf Nightshade: Apply when plants begin to flower in spring. Upright Prairie Coneflower: Apply when plants are 2-6 inches tall, before flowering. Curly Dock: Apply up to bolting. Ironweed: Apply up to bud stage. Yankeeweed: Apply when plants are 8 to 10 inches tall. |

4/1/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|---|--|---|--|
| goldaster, gray goldaster, narrowleaf goldenweed, common goldenweed, Drummond (Isocoma spp.) | 1-2 pts. Alligare Picloram 22K + 0.5-1 lb. a.e. 2,4-D or 2 pts. Alligare Picloram 22K | 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4-D | Gray and Narrowleaf Goldaster: Apply in oil-water emulsion in spring during bud stage (prebloom). Thorough coverage is essential. Goldenweed: Apply in spring (April-June) when there is substantial canopy development as a result of good growing conditions. Add an agricultural surfactant at 0.25% - 0.5% or apply in oil-water emulsion. Increase spray volume, 4-5 gpa by air or 15-20 gpa by ground, to ensure thorough coverage. |
| Poisonous Plants such as groundsel (Senecio spp.) loco, woolly loco, Wooton (garbancillo) | ¾ - 1 pt. Alligare Picloram 22K + 0.5-1.0 lb. a.e. 2,4-D or 1 ½ - 2 pts. Alligare Picloram 22K | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D | General: Apply in fall or winter when there is adequate soil moisture and weeds are actively growing. See the "General Use Precautions" section of this label for note on grazing treated poisonous plants. Locoweeds: To improve wetting of locoweeds, use an agricultural surfactant at 0.25%-0.5% or apply in oil-water emulsion. |
| Woody Plants: | Tank Mixing: Within rate ranges for listed products, consult local recommendations | | |
| huisache (suppression) | 2 pts. Alligare Picloram 22K + 1 pt. Remedy™ | 2 qt. Alligare Picloram 22K + 1 qt. Remedy™ | Fall application is recommended, however, fall applications will not provide satisfactory control of other woody species in the South Texas mixed brush complex. Performance can be erratic. |
| juniper, including, alligator redberry Utah one-seeded eastern redcedar pinyon pine | -- | 4 qts. Alligare Picloram 22K | Apply May through July. Complete coverage is essential. Results with ashe juniper may be variable with high volume foliar application. |
| pricklypear, lindheimer (unburned rangeland) | 2 pts. Alligare Picloram 22K | 4 qts. Alligare Picloram 22K | Application may be made anytime, but optimum time is late August to early November. Onset of herbicidal activity is very slow and may continue for two years or longer. Good coverage is essential. |
| pricklypear, lindheimer (burned rangeland) | 1 pt. Alligare Picloram 22K | 2 qts. Alligare Picloram 22K | Conduct intense controlled burns from December through March and apply Alligare Picloram 22K mid-April through May. Rainfall |

42/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|--|---|---|--|
| | | | following burning can also stimulate prolific resprouting of the burned plants. Good coverage is also essential. |
| rose, Macartney rose, multiflora | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D | 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4-D | Apply in the spring or fall when conditions are favorable for plant growth. Use an agricultural surfactant (0.5% v/v) or apply as an oil-water emulsion. Ensure thorough and uniform coverage by applying at higher spray volume, 5 or more gpa by air or 20 or more gpa by ground. Avoid treatment less than 9 to 12 months after mowing when plants have a high percentage of new growth. Repeat treatment as necessary. |
| tallowtree, Chinese | 1 qt. Alligare Picloram 22K + 2 lbs. a.e. 2,4-D or 1 pt. Remedy™ | 2 qts. Alligare Picloram 22K or 1-2 qts. Alligare Picloram 22K + 2-4 lbs. a.e. 2,4-D or 1 qt. Remedy™ | Apply in the spring or fall when conditions are favorable for plant growth. Use an agricultural surfactant (0.5% v/v) or use an oil-water emulsion and higher spray volumes, 5 gpa or more by air and 20 gpa or more by ground. |
| South Texas mixed brush, including, acacia, blackbrush acacia, catclaw acacia, twisted granjeno guajillo mesquite prickly pear tasajillo | 2 pts. Alligare Picloram 22K + 2/3-1 1/3 pt. Reclaim™ or 1 to 2 pts. Remedy™ | 2 qts. Alligare Picloram 22K + 2-3 pts. Remedy™ or 1-2 qts. Reclaim™ | Apply in oil-water emulsion. Use 4 or more gpa by air or 20 or more gpa by ground. For application timing for mesquite, see comments in section on mesquite control. Tank mixing with Reclaim™ will provide improved control of pricklypear and legume species such as mesquite and acacias while tank mixing with Remedy™ will provide improved control of non-legume species such as granjeno, oaks and hackberry. |
| mesquite | 1-2 pts. Alligare Picloram 22K + 2/3-1 1/3 pt. Reclaim™ or 2 pts. Alligare Picloram 22K + 1 pt. Remedy™ | 1-2 qts. Alligare Picloram 22K + 1-2 qts. Reclaim™ or 1 1/2-3 pts. Remedy™ | Alligare Picloram 22K Alone: Apply as a water spray or oil-emulsion (see Mixing Instructions) in 4 or more gpa by air or 10 or more gpa by ground. Increase spray volumes with increasing brush density and height to ensure adequate coverage. Where control of pricklypear cactus is desired, use the 2 pint/acre rate of Alligare |

43/47

| Weed Species | Broadcast Application (Rate/acre) | High Vol. Foliar (Rate/100 gal.) | Specific Use Directions |
|--|--|----------------------------------|-------------------------|
| | | | Picloram 22K. |
| <p>Alligare Picloram 22K in Tank Mix: Tank mixing with Reclaim™ will provide control of pricklypear and improved control of legume species such as mesquite and acacias while tank mixing with Remedy™ will provide improved control of non-legume species such as granjeno, oaks and hackberry. Regrowth mesquite should be at least 4 ft. tall prior to treatment. See labels for Reclaim™ and Remedy™ for additional treatment recommendations and information on mesquite control. Within rate ranges given for Alligare Picloram 22K and tank mix products, consult local recommendations.</p> | | | |
| <p>Timing and Factors in Control: The herbicidal response of mesquite is strongly influenced by environmental conditions as well as foliage condition and stage of growth. For best results, apply when new growth foliage has turned from light to dark green, when the soil temperature has reached 75°F to 83°F at a depth of 12-18 inches, and soil moisture is adequate for plant growth. Application should be made within 45 days after the critical soil temperature at the 12-18 inch depth has been reached, or if Alligare Picloram 22K is applied in combination with Reclaim™ within 60 days. Product performance may be adversely affected if application is made before mesquite foliage has turned from light to dark green or if foliage has been injured or removed by late frost, insects, hail or plant diseases. Do not apply if mesquite exhibits new (light green) growth in response to significant rainfall during the growing season. Soil temperatures at the 12-18 inch depth may vary with soil texture and drainage. Coarse-textured (sandy) soils warm up sooner than fine-textured soils (clay) soils and dry soils warm up more quickly than wet soils.</p> | | | |
| <p>Re-application: Do not reapply in the same growing season. Retreatment will not be effective until woody plants develop sufficient new foliage for interception, uptake, and translocation of the herbicide to plant roots.</p> | | | |
| ashe juniper eastern redcedar eastern persimmon | <p>General: Apply Alligare Picloram 22K undiluted as a spot concentrate application prior to periods of expected rainfall. Apply directly to the soil within the dripline and on the upslope side of the tree. Application to trees taller than 12 feet is not recommended. See directions for "Soil Spot Concentrate" in "Application Methods" section.</p> <p>Ashe Juniper: Apply 4 to 6 ml per 3 ft. of plant height in the spring (April-May).</p> <p>Eastern Redcedar: Apply 3 to 4 ml per 3 ft. of plant height in either spring (April-May) or fall (September-October).</p> <p>Eastern Persimmon: Apply 2 to 4 ml per inch of stem diameter in spring (March through May).</p> | | |

Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: All Areas West of the Mississippi River – Carpet Roller Use Recommendations)

| Brush Species | Amount of Alligare Picloram 22K | Specific Use Directions |
|---|---|---|
| mesquite regrowth | 1 gal. alone or with 2 qts. Reclaim™ herbicide to make 8 gals. of spray mixture | Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply from May through August, but preferably in May and June, when moisture availability is sufficient to allow normal plant growth. |
| huisache/blackbrush | 1 gal. alone or with 2 to 3 qts. of Reclaim™ to make 8 gals. of spray mixture | Include 1 ounce of a recommended agricultural surfactant per gallon of herbicide-water mixture (0.75% vol/vol). Apply in the fall. |
| <p>Refer to the Rangeland, Permanent Grass Pastures and Non-Cropland (Specific Use Directions: Texas, Oklahoma, and New Mexico) table for information on timing and factors in control of mesquite.</p> | | |

44/47

SPECIFIC USE DIRECTIONS FOR SMALL GRAINS, FALLOW CROPLAND, AND CRP ACRES

Spring Seeded Barley, Oats, and Wheat Not Underseeded With a Legume (Which is Not Flood or Sub-Irrigated and Not Rotated to Broadleaf Crops)

Do not apply to winter wheat or barley as crop injury may occur. Do not treat durum wheat since some varieties of durum wheat may be injured.

Use Alligare Picloram 22K for the control of susceptible annual broadleaf weeds such as (but not limited to) volunteer sunflower, wild buckwheat, lambsquarters, pigweed, Russian thistle, and sowthistle. This product may cause shorter straw on some varieties of cereals but grain yields are usually not affected.

Use Restrictions

- Do not apply Alligare Picloram 22K within 50 days before harvest.
- Do not graze or feed forage from treated areas for 2 weeks after treatment. Do not harvest hay from treated grain fields.
- Use only on land that will be planted the following year to grass, barley, oats, wheat, grain sorghum or fallowed. Do not plant grain sorghum within 8 months after application. This product is not intended for use on land planted to sweet sorghum. Do not apply more than 1 1/2 fluid ounces of Alligare Picloram 22K per acre during the small grain growing season.

Use Alligare Picloram 22K for the control of susceptible annual broadleaf weeds such as (but not limited to) volunteer sunflower, wild buckwheat, lambsquarters, pigweed, Russian thistle, and sowthistle.

Broadcast Treatment (Ground and Aerial Applications)

Alligare Picloram 22K can be applied as a single broadcast treatment by ground or aurally to control several broadleaf weeds by itself or as a tank mix with 2,4-D, MCPA, or sulfonyleurea herbicides such as Ally[®]. Apply Alligare Picloram 22K at the rates suggested in the following table in 1 to 5 gallons of water per acre by air or in 5 to 20 gallons of water per acre by ground. The addition of surfactants may aid control under dry conditions, but may cause injury to grain if used over the top. Read and follow directions and precautions on other product labels when tank mixing.

Application Timing

Spring Wheat, Barley and Oats: Apply from the 3 to 5 leaf stage to early jointing stage of growth as indicated in the table below. Applications at the 3 to 5 leaf stage occasionally cause slight head malformations and straw shortening but normally do not affect yield.

Use Rates for Spring Wheat, Barley and Oats¹

| Weed Species | Weed Growth Stage ² | Grain Growth Stage | Amounts of Each Product Per Acre ³ | | |
|--|--------------------------------|--------------------------------|---|------------------------------|------------------------------|
| | | | Alligare Picloram 22K | 4 lb. a.e./gal 2,4-D or MCPA | 6 lb. a.e./gal 2,4-D or MCPA |
| More susceptible species, such as: lambsquarters pennycress wild mustard mayweed | 3 inches | 3 to 5 leaf to early tillering | 1 fl. oz. | 1/2 pint | 1/3 pint |
| | 3 to 6 inches | Tillering to, early jointing | 1 1/2 fl. oz. | 3/4 pint | 1/2 pint |

45/47

| Weed Species | Weed Growth Stage ² | Grain Growth Stage | Amounts of Each Product Per Acre ³ | | |
|---|--------------------------------|-----------------------------|---|------------------------------|------------------------------|
| | | | Alligare Picloram 22K | 4 lb. a.e./gal 2,4-D or MCPA | 6 lb. a.e./gal 2,4-D or MCPA |
| Less susceptible species such as: volunteer sunflower wild buckwheat Russian thistle pigweed Canada thistle, top growth suppression | 1 to 6 inches | Tillering to early jointing | 1 ½ fl. oz. | ¾ to 1 pint | ½ to 2/3 pint |

¹For oats, do not tank mix with 2,4-D herbicides.

²For best results, treat when weeds have 2 to 4 leaves and are actively growing.

³When measuring small amounts of Alligare Picloram 22K, special care should be taken not to exceed suggested rates.

Fallow Cropland (Not Rotated to Broadleaf Crops)

Apply Alligare Picloram 22K as a post harvest or fallow treatment in continuous grain or during the fallow period. Alligare Picloram 22K may be applied alone or in tank mix combination with 2,4-D or other herbicides registered for this use. Apply in 1 or more gallons of water per acre by air or 5 or more gallons per acre by ground.

Application Rates

Annual Weeds: To control annual weeds such as Russian thistle and wild buckwheat, apply ¼ to ½ pint per acre of Alligare Picloram 22K in tank mix combination with ½ to 1 lb. a.e. of 2,4-D or other herbicides registered for use on fallow land. Apply when weeds are actively growing.

Field Bindweed: Apply ½ to 1 pint per acre of Alligare Picloram 22K plus ½ lb. to 1 lb. a.e. per acre of 2,4-D when bindweed is actively growing. Optimum time for treatment is when plant runners reach 8 to 12 inches. Use ½ pint per acre to control light to moderate infestations under good growing conditions or to reduce the potential for crop injury. Use higher rates for heavy infestations and longer term control. Some regrowth will occur the following season and a re-treatment program for successive years is recommended.

Canada thistle: Apply 1 pint per acre of Alligare Picloram 22K plus 1 lb. a.e. per acre of 2,4-D when the majority of thistle plants are emerged but prior to bud stage.

Crop Rotation

Use only on land to be planted the following year to grass, barley, oats, wheat or fallow land. Many broadleaf crops are extremely sensitive to soil residues of Alligare Picloram 22K. Do not plant sensitive broadleaf crops for 36 months after treatment or until soil residues have declined to a safe level as indicated by an adequately sensitive bioassay using the intended broadleaf crop. A bioassay is recommended following treatment prior to planting any sensitive broadleaf crop.

Preplant Interval

A preplant interval following application of Alligare Picloram 22K prior to planting small grains is recommended to reduce or eliminate potential crop injury and/or yield reduction. The possibility for crop injury or yield reduction to occur depends on application rate, soil organic matter, rainfall, temperature and incidence of cereal diseases. Adequate soil moisture and soil temperature during the preplant interval is important in reducing, but may not eliminate, the risk of crop injury. When considering use of Alligare Picloram 22K on fallow land, growers should consider the benefit of weed control against the risk of crop damage and treat only if the risk of injury to small grains can be tolerated. The following preplant intervals are recommended:

46/47

For applications up to ½ pint per acre, allow a minimum of 45 days of soil temperatures above 40°F between application and planting.

For applications of greater than ½ pint and up to 1 pint per acre, allow a minimum of 60 days of soil temperatures above 40°F between application and planting, except in the states of Idaho, North Dakota, Nebraska, Montana, Oregon, South Dakota, Washington and Wyoming, where the minimum preplant interval is 90 days.

Restrictions:

- Do not apply more than 1 pint per acre as a broadcast treatment per annual growing season.
- Spot Treatment: See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing. Spot treatments of Alligare Picloram 22K at rates over 1 pint per acre can be made on fallow, non-irrigated cropland if the treated areas comprise less than 10% of the immediate field in any one year. Alligare Picloram 22K should not be applied to cropland at rates exceeding 2 quarts per acre. When Alligare Picloram 22K is applied at rates above 1 pint per acre, injury to small grains may result for periods up to two years after treatment.

Conservation Reserve Program (CRP) for Seeding to Permanent Grasses Only

Do not use Alligare Picloram 22K if legumes are a desired cover during CRP.

Conditions that stress grasses, such as drought, will increase potential for injury to the grass at all stages of growth.

To reduce potential damage to subsequent small grain crops, use the lower rate or discontinue the use of Alligare Picloram 22K at least 2 years prior to the seeding of small grain crops. After CRP, do not plant broadleaf crops in treated acres until an adequately sensitive bioassay shows that no detectable picloram is present in the soil.

Broadcast Treatment (Ground and Aerial Applications)

Applications of Alligare Picloram 22K should be made after perennial grasses are well established (have developed a good secondary root system and show good vigor). Most perennial grasses show improved tolerance to the herbicide at this stage of development.

For control of actively growing perennial weeds, use up to 2 pints per acre of Alligare Picloram 22K after the grass is established. For best results, use in 1 or more gallons of water per acre by air or in 5 or more gallons of water per acre by ground. Increasing the rate of application can increase the risk of injury.

For control of actively growing susceptible annual broadleaf weeds, (including Russian thistle) apply ¼ to ½ pint per acre of Alligare Picloram 22K. Alligare Picloram 22K can also be tank mixed with ½ to 1 pound per acre of 2,4-D where 2,4-D sensitive species are present. Read and follow all directions for use and use precautions on other product labels.

Spot Treatment

See "Spot Treatment" in "Mixing and Application Methods" section for directions for calibration, spray volume determination and mixing. For spot applications when perennial grasses are established, use 1 to 4 pints per acre of Alligare Picloram 22K. Rates of 2 pints per acre or more should only be used for control of deep-rooted perennial broadleaf weeds.

Precaution: Alligare Picloram 22K at rates over 2 pints per acre may suppress certain established grasses such as bromegrass, bluegramma and buffalograss. However, subsequent grass growth should be improved by release from weed competition.

47/47

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: The active ingredient in this product may crystallize and settle out of solution if product is exposed to subfreezing temperatures. Under these conditions, warm product to at least 40°F and agitate well to dissolve any crystallized material prior to use. Open dumping is prohibited.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Every 2 years starting January 1, 2008, the registrant will offer training to applicators which will cover application techniques and product stewardship particular to their use(s) of this product (Alligare Picloram 22K, Registration No. 81927-18). Applicators of this product must be able to provide certification of such training on demand to the State, Tribal or Federal enforcement agent.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Reclaim™ and Remedy™ are trademarks of Dow AgroSciences LLC.
 Ally® and Escort® are registered trademarks of E.I. duPont de Nemours & Co., Inc.

EPA 20080320