

81927-22

11-02-2011

1/20



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Registration
 Number:
 81927-22

Date of Issuance:
 NOV 2 2011

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: **Unconditional**

Name of Pesticide Product:
Alligare Ecomazapyr 2 SL

Name and Address of Registrant (include ZIP Code):
Alligare, LLC c/o **Pyxis Regulatory Consulting, Inc.**
13 N. 8th Street **4110 136th Street, NW**
Opelika, AL 36801 **Gig Harbor, WA 98332**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted on November 2, 2011. EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision (RED) for **Imazapyr** have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

The Basic Confidential Statement of Formula (CSF) dated October 24, 2008 is acceptable, and supersedes all previously-accepted CSFs.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after twelve (12) months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

If you have any questions regarding this Notice, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Signature of Approving Official:

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

A handwritten signature in black ink, appearing to read "Kable Bo Davis".

Date:

NOV 2 2011

2
20

Alligare Ecomazapyr 2 SL

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. Alligare Ecomazapyr 2 SL may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water. Alligare Ecomazapyr 2 SL may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites.

ACTIVE INGREDIENT:	% BY WT.
Isopropylamine salt of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)*	27.8%
OTHER INGREDIENTS	<u>72.2%</u>
TOTAL	100.0%

*Equivalent to 22.6% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 2 pounds acid equivalent per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor for further treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to so by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call 1-800-424-9300.	

EPA Reg. No. 81927-22

EPA Est. No.

Manufactured For:
Alligare, LLC
13 N. 8th Street
Opelika, AL 36801

Net Contents:

ACCEPTED

NOV 2 2011

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 81927-22

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION! Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (except for pilots)
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are given for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Do not apply to water except as specified in this label. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss may cause the suffocation of some aquatic organism. Do not contaminate water when disposing of equipment washwaters or rinsate. This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift precautions on the label.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of **Alligare Ecomazapyr 2 SL** must be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply **Alligare Ecomazapyr 2 SL** or spray solutions of **Alligare Ecomazapyr 2 SL** in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Keep containers closed to avoid spills and contamination.

PRODUCT INFORMATION

Alligare Ecomazapyr 2 SL is an aqueous solution intended to be mixed with water and surfactant(s) for application to control floating and emergent undesirable vegetation (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section) in or near bodies of water which may be

4
20

flowing, non-flowing or transient. **Alligare Ecomazapyr 2 SL** may be applied to aquatic sites that include lakes, rivers, streams, ponds, seeps, drainage ditches, canals, reservoirs, swamps, bogs, marshes, estuaries, bays, brackish water, transitional areas between terrestrial and aquatic sites and seasonal wet areas. See **AQUATIC USES, PRECAUTIONS AND RESTRICTIONS** section of this label for precautions, restrictions, and instructions on aquatic uses. **Alligare Ecomazapyr 2 SL** may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites (see **AERIAL APPLICATIONS AND GROUND APPLICATIONS** sections for additional details).

Read and observe the following directions if aquatic sites are present in terrestrial non-crop areas and are part of the intended treatment area:

Alligare Ecomazapyr 2 SL controls most annual and perennial grasses, broadleaf weeds and many brush and vine species. **Alligare Ecomazapyr 2 SL** will provide some residual control of undesirable species that germinate above the waterline. **Alligare Ecomazapyr 2 SL** controls vegetation by absorption through emergent leaves and stems, from which it is translocated throughout the plant, where it accumulates in rapidly-growing meristematic tissue. Treated plants stop growing soon after treatment. Chlorosis (yellowing of plant tissue) first appears in the newest leaves, and necrosis spreads from this point. In perennials, **Alligare Ecomazapyr 2 SL** is translocated into and kills underground or submerged storage organs to prevent regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two or more weeks after application. Complete kill of plants may not occur for several weeks. Applications of **Alligare Ecomazapyr 2 SL** are rainfast within one hour after treatment.

Alligare Ecomazapyr 2 SL will not control plants which are completely submerged or have a majority or their foliage under water.

Alligare Ecomazapyr 2 SL must be applied to the emergent foliage of the target vegetation, since it has little-to-no activity on submerged vegetation. Residual concentrations of **Alligare Ecomazapyr 2 SL** from direct application to water are not expected to be sufficient to provide control of undesirable vegetation. Apply **Alligare Ecomazapyr 2 SL** so as to maximize spray contact onto the target vegetation while minimizing overspray onto adjacent or underlying water. For optimum performance, include a surfactant (see **ADJUVANTS** section for specific directions) in the spray solution and apply to actively growing weeds. **Alligare Ecomazapyr 2 SL** may be applied in spot treatments by using low-volume directed application techniques or broadcast by using ground equipment, watercraft or helicopter.

Apply Alligare Ecomazapyr 2 SL with surface or helicopter application equipment in at least 5 gallons of water per acre. When applying by helicopter, follow directions under the **AERIAL APPLICATIONS** section of this label, otherwise refer to section on **GROUND APPLICATIONS** when using surface equipment.

Applications to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. **DO NOT** apply to bodies of water or portions of bodies of water where emergent and/or floating weeds do not exist.

When target vegetation covers a large percentage of the surface area of impounded water, apply **Alligare Ecomazapyr 2 SL** to alternating strips (i.e., **DO NOT** apply to every swath) to avoid oxygen depletion due to decaying vegetation, since oxygen depletion may result in the suffocation of sensitive aquatic organisms. **DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days before retreatment. Begin treatment along the shore and proceed outward in alternating swaths as described above to allow aquatic organisms to move into untreated areas.

Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash for one hour after application.

Apply **Alligare Ecomazapyr 2 SL** at 2 to 6 pints per acre, depending on weed species and weed density. Use higher rates for heavy weed pressure. Refer to the **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE**

5
29

ECOMAZAPYR 2 SL section of this label for specifics. **DO NOT** exceed 6 pints per acre (1.5 lb ai/A) per year.

Alligare Ecomazapyr 2 SL may also be applied as a draw down treatment in areas described above. Apply **Alligare Ecomazapyr 2 SL** to weeds after water has been drained and allow 14 days before reintroduction of water.

PRECAUTIONS FOR AVOIDING INJURY TO NON-TARGET PLANTS

Alligare Ecomazapyr 2 SL can occasionally affect non-target or desirable vegetation by root uptake of the herbicide from treated soil. Injury or loss of non-target plants may result if **Alligare Ecomazapyr 2 SL** is applied onto or near desirable plants, or to areas where their roots extend, or in areas where treated soil may be washed or moved into contact with their root zone.

When making applications along shorelines where desirable plants may be present, exercise caution to avoid spray contact onto their foliage or spray application to the soil in which they are rooted. Shoreline plants with roots that extend into the waters where **Alligare Ecomazapyr 2 SL** has been applied will generally not be adversely affected.

If treated vegetation is to be removed from the application site, **DO NOT** use the vegetative matter as mulch or compost on or around desirable species.

IMPORTANT

DO NOT use on food crops. **DO NOT** apply this product within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water, such as a lake, pond or reservoir. **DO NOT** apply to water used for irrigation except as described in **APPLICATION TO WATERS USED FOR IRRIGATION** section of this label. Keep away from fertilizers, insecticides, fungicides, and seeds. **DO NOT** drain or flush equipment on or near desirable plants, or onto areas where their roots may extend, or in locations where the chemical or treated soil may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, driveways, tennis courts or similar areas where roots of desirable vegetation may extend and be exposed to potential injury and/or mortality from root uptake of **Alligare Ecomazapyr 2 SL**. **DO NOT** side trim desirable vegetation with this product unless severe injury or plant death is acceptable. Prevent drift of spray to desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

Do not enter or allow others to enter treated areas until sprays have dried.

SPRAY DRIFT MANAGEMENT

Spray Drift: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the entity authorizing spraying are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. **DO NOT** apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, and temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces the drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **WIND, TEMPERATURE AND HUMIDITY, AND TEMPERATURE INVERSIONS**).

Controlling Droplet Size:

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. **DO NOT** use nozzles producing a mist droplet spray.

Application Height: Making applications at the lowest possible height (helicopter, ground driven spray bloom) that is safe and practical reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the treatment area, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind: Drift potential is lowest between wind speeds of 3-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during temperature inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind Erosion: Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Aerial Applications: Applicators are required to use a Coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater for release heights below 10 feet. Applicators are required to use a Very Coarse or coarser droplet size or, if specifically using a spinning atomizer nozzle, applicators are required to use a VMD of 475 microns or greater for release heights above 10 feet. Applicators must consider the effects of nozzle orientation and flight speed when determining droplet size.

Applicators are required to use upwind swath displacement.

The boom length must not exceed 60% of the wingspan or 90% of the rotor blade diameter to reduce spray drift.

Applications with wind speeds less than 3 mph and with wind speeds greater than 10 mph are prohibited.

Applications into temperature inversions are prohibited.

Ground Boom Applications: Applicators are required to use a nozzle height below 4 feet above the ground or plant canopy and Coarse or coarser droplet size (ASABE S572) or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater.

Applications with wind speeds greater than 10 mph are prohibited.

Applications into temperature inversions are prohibited.

AQUATIC SPECIES CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

Alligare Ecomazapyr 2 SL will control the following target species when used as directed in the table below. Rate specifications are expressed as product volume for broadcast applications and as a % solution for directed applications including spot treatments. For % solution applications, **DO NOT** apply more than the equivalent of 3 quarts of **Alligare Ecomazapyr 2 SL** per acre.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Floating Species		
*Duckweed	<i>Lemna minor</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Duckweed, Giant	<i>Spirodela polyrriza</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Frogbit	<i>Limnobium spongia</i>	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Spatterdock	<i>Nuphar luteum</i>	Apply a tank-mix of 2-4 pints/acre Alligare Ecomazapyr 2 SL plus 4-6 pints/acre glyphosate (0.5% Alligare Ecomazapyr 2 SL plus 1.5% glyphosate) in 100 GPA water for best control. Completely cover all actively-growing emergent foliage.
*Water Hyacinth	<i>Eichhornia crassipes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water to actively growing foliage.
*Water Lettuce	<i>Pistia stratiotes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Emerged Species		
*Alligatorweed	<i>Alternanthera philoxeroides</i>	1-4 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.
*Arrowhead, Duck-potato	<i>Sagittaria</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Bacopa, lemon	<i>Bacopa</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Parrot feather	<i>Myriophyllum aquaticum</i>	Apply 2-4 pints in 100 GPA water to actively growing emergent foliage. Foliage must be above water for sufficient uptake.
*Pennywort	<i>Hydrocotyle</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Pickerelweed	<i>Pontederia cordata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Taro, wild; Dasheen; Elephant's Ear; Coco Yam	<i>Colocasia esculentum</i>	4-6 pints/acre (1.5% solution) in 100 GPA with a high quality 'sticker' adjuvant. Completely cover all actively-growing emergent foliage.
*Water lily	<i>Nymphaea odorata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Water Primrose	<i>Ludwigia uruguayensis</i>	4-6 pints/acre (1.5% solution) in 100 GPA water, Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.
Terrestrial/Marginal		
*Soda Apple, aquatic; Nightshade	<i>Solanum tampicense</i>	2 pints/acre applied to foliage.
*Bamboo, Japanese	<i>Phyllostachys</i> spp.	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Brazilian Pepper; Christmasberry	<i>Schinus terebinthifolius</i>	2-4 pints/acre applied to foliage.
Cattail	<i>Typha</i> spp.	2-4 pints (1% solution) applied to actively-growing, green foliage after full leaf elongation. Lower rates will control cattail in the north; higher rates are needed in the south.
Chinese Tallow Tree	<i>Sapium sebiferum</i>	16-24 fl oz applied to foliage.
Cogon grass	<i>Imperata cylindrica</i>	Burn foliage, till area, then treat in fall at 2 quarts/acre Alligare Ecomazapyr 2 SL plus MSO applied to new growth.
Cordgrass, prairie	<i>Spartina</i> spp.	4-6 pints applied to actively growing foliage.
*Cutgrass	<i>Zizaniopsis miliacea</i>	4-6 pints applied to actively growing foliage.
*Elephant Grass; Napier Grass	<i>Pennisetum purpureum</i>	3 pints/acre applied to actively growing foliage.
*Flowering rush	<i>Butumu typla</i>	2-3 pints applied to actively growing foliage.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Giant Reed, Wild Cane	<i>Arundo donax</i>	4-6 pints/acre applied in spring to actively growing foliage.
*Golden Bamboo	<i>Phyllostachys aurea</i>	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Junglerice Knapweeds	<i>Echinolchloa colonum</i> <i>Centaurea species</i>	3-4 pints applied to actively growing foliage. Russian Knapweed- Apply 2-3 pints plus 1 quart/acre MSO in the fall after senescence begins.
Knotweed, Japanese (see <i>Fallopia japonica</i>) Melaleuca; Paperbark Tree	<i>Polygonum cuspidatum</i> <i>Melaleuca quinquenervia</i>	3-4 pints/acre applied postemergence to actively growing foliage. For established stands, apply 6 pints/acre Alligare Ecomazapyr 2 SL plus 6 pints/acre glyphosate plus adjuvant. For best results, use 4 quarts/A methylated seed oil for adjuvant. For ground foliar application, uniformly apply to completely cover all actively-growing foliage. For aerial foliar application, apply in at least two passes in a cross pattern at 10 GPA. For spot treatment, apply a 25% solution of Alligare Ecomazapyr 2 SL plus 25% solution of glyphosate plus 1.25% MSO in water as a frill or stump treatment.
*Nutgrass; Kili'p'opu	<i>Cyperus rotundus</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied early postemergence.
*Nutsedge	<i>Cyperus</i> spp.	2-3 pints postemergence to foliage or pre-emergence incorporated. Non-incorporated preemergence applications will not provide control.
Phragmites; Common Reed	<i>Phragmites australis</i>	4-6 pints/acre applied to actively-growing, green foliage after leaf elongation. Completely cover all actively-growing foliage. If stand has a substantial amount of old stem tissue, mow or burn, allow to regrow to approximately 5' tall before treatment. Lower rates will control phragmites in the north; higher rates are needed in the south.
*Poison Hemlock	<i>Conium maculatum</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied preemergence or early postemergence up to rosette stage before flowering.
Purple Loosestrife Reed canarygrass	<i>Lythrum salicaria</i> <i>Phalaris arundinacea</i>	1 pint/acre applied to actively growing foliage. 3-4 pints/acre applied to actively growing foliage.
Rose, swamp	<i>Rosa palustris</i>	2-3 pints/ acre applied to actively growing foliage.
Russian-Olive Saltcedar; Tamarisk	<i>Elaeagnus angustifolia</i> <i>Tamarix species</i>	2-4 pints/acre or 1% solution applied to foliage. Aerially apply 2 quarts Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS to actively growing foliage during flowering. For spot spraying, use a 1% solution of Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS and spray to wet foliage. After application wait at least two years before

10
20

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
		disturbing treated saltcedar. Earlier disturbance can reduce overall control.
Smartweed	<i>Polygonum</i> spp.	2 pints/acre applied early postemergence.
Sumac	<i>Rhus</i> spp.	2-3 pints/acre applied to foliage.
Swamp Morning Glory;	<i>Ipomoea aquatica</i>	1-2 pints/acre Alligare Ecomazapyr 2 SL plus
Water Spinach;		1 quart/acre MSO applied at early
Kangkong		postemergence.
Torpedo grass	<i>Panicum repens</i>	4 pints/acre (1-1.5% solution). Completely cover all actively growing foliage.
*White Top; Hoary	<i>Cardaria draba</i>	1-2 pints/acre applied in spring to foliage during flowering.
Cress		
Willow	<i>Salix</i> spp.	2-3 pints/acre Alligare Ecomazapyr 2 SL . Completely cover all actively growing foliage.

* Not approved for use in California

ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

When used as directed on terrestrial sites, **Alligare Ecomazapyr 2 SL** will provide preemergence or postemergence control with residual control of the following vegetation species. Annual weeds may be controlled by preemergence or postemergence applications of **Alligare Ecomazapyr 2 SL**. Use postemergence treatments of **Alligare Ecomazapyr 2 SL** for established biennials and perennial vegetation control.

The tables below show broadcast rates and indicate relative weed sensitivity. It is important to consider relative weed sensitivity when preparing low volume spray solutions (see **LOW VOLUME** section of **GROUND APPLICATIONS** section), since low volume treatments apply less **Alligare Ecomazapyr 2 SL** per acre than is shown for the broadcast treatments.

The relative sensitivity of the species listed below can also be used to determine the relative risk of causing non-target plant injury if any of those species are considered to be desirable within the treatment area.

Resistant Biotypes: Some weeds listed below may have naturally-occurring biotypes (plants within a given species that have a slightly different but distinct genetic makeup from other plants of that species) that are not effectively controlled by this and/or other herbicides (Oust®) with the ALS/AHAS enzyme-inhibiting mode of action. If naturally-occurring ALS/AHAS-resistant biotypes are present in area, tank mix or apply **Alligare Ecomazapyr 2 SL** sequentially with a registered herbicide that depends on a different mode of action to ensure control.

GRASSES

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 2-3 pints per acre¹		
Annual bluegrass	<i>Poa annua</i>	A
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	A
Canada bluegrass	<i>Poa compressa</i>	P
Downy brome	<i>Bromus tectorum</i>	A
Fescue	<i>Festuca</i> spp.	A/P
Foxtail	<i>Setaria</i> spp.	A
Italian ryegrass	<i>Lolium multiflorum</i>	A
Johnsongrass	<i>Sorghum halepense</i>	P
Kentucky bluegrass	<i>Poa pratensis</i>	P

COMMON NAME	SPECIES	GROWTH HABIT ²
Lovegrass	<i>Eragrostis</i> spp	A/P
*Napier grass	<i>Pennisetum purpureum</i>	P
Orchardgrass	<i>Dactylis glomerata</i>	P
Paragrass	<i>Brachiaria mutica</i>	P
Quackgrass	<i>Agropyron repens</i>	P
Sandbur	<i>Cenchrus</i> spp	A
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Smooth brome	<i>Bromus inermis</i>	P
Vaseygrass	<i>Paspalum urvillei</i>	P
Wild Oats	<i>Avena fatua</i>	A
Witchgrass	<i>Panicum capillare</i>	A

Apply 3-4 pints per acre¹

Barnyardgrass	<i>Echinochloa crus-gali</i>	A
Beardgrass	<i>Andropogon</i> spp.	P
Bluegrass, Annual	<i>Poa annua</i>	A
*Bulrush	<i>Scirpus validus</i>	P
Cheat	<i>Bromus secalinus</i>	A
Crabgrass	<i>Digitaria</i> spp	A
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	A
Fall Panicum	<i>Panicum dichotomiflorum</i>	A
Goosegrass	<i>Eleusine indica</i>	A
Itchgrass	<i>Rottboellia exaltata</i>	A
Lovegrass	<i>Eragrostis</i> spp.	A
*Maidencane	<i>Panicum hemitomon</i>	A
Panicum, Browntop	<i>Panicum fasciculatum</i>	A
Panicum, Texas	<i>Panicum texanum</i>	A
Prairie threeawn	<i>Aristida oligantha</i>	P
Sandbur, Field	<i>Cenchrus incertus</i>	A
Signalgrass	<i>Brachiaria platyphylla</i>	A
Wild barley	<i>Hordeum</i> spp.	A
Wooly Cupgrass	<i>Eriochloa villosa</i>	A

Apply 4-6 pints per acre¹

Bahiagrass	<i>Paspalum notatum</i>	P
Bermudagrass ³	<i>Cynodon dactylon</i>	P
Big bluestem	<i>Andropogon gerardii</i>	P
Dallisgrass	<i>Paspalum dilatatum</i>	P
Feathertop	<i>Pennisetum villosum</i>	P
Guineagrass	<i>Panicum maximum</i>	P
Saltgrass ³	<i>Distichlis stricta</i>	P
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Sprangletop	<i>Leptochloa</i> spp.	A
Timothy	<i>Phleum pretense</i>	P
Wirestem muhly	<i>Muhlenbergia frondosa</i>	P

BROADLEAF WEEDS

Apply 2-3 pints per acre¹

Burdock	<i>Arctium</i> spp.	B
Carpetweed	<i>Mollugo verticillata</i>	A
Carolina geranium	<i>Geranium carolinianum</i>	A

COMMON NAME	SPECIES	GROWTH HABIT ²
Clover	<i>Trifolium</i> spp.	A/P
Common chickweed	<i>Stellaria media</i>	A
Common ragweed	<i>Ambrosia artemisiifolia</i>	A
Dandelion	<i>Taraxacum officinale</i>	P
Dog fennel	<i>Eupatorium capillifolium</i>	A
Filaree	<i>Erodium</i> spp.	A
Fleabane	<i>Erigeron</i> spp.	A
Hoary vervain	<i>Verbena stricta</i>	P
Indian mustard	<i>Brassica juncea</i>	A
Kochia	<i>Kochia scoparia</i>	A
Lambsquarters	<i>Chenopodium album</i>	A
*Lespedeza	<i>Lespedeza</i> spp.	P
Miners lettuce	<i>Montia perfoliata</i>	A
Mullein	<i>Verbascum</i> spp.	B
Nettleleaf goosefoot	<i>Chenopodium murale</i>	A
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>	P
Pepperweed	<i>Lepidium</i> spp.	A
Pigweed	<i>Amaranthus</i> spp.	A
Puncturevine	<i>Tribulus terrestris</i>	A
Russian thistle	<i>Salsola kali</i>	A
Smartweed	<i>Polygonum</i> spp.	A/P
Sorrell	<i>Rumex</i> spp.	P
Sunflower	<i>Helianthus</i> spp.	A
Sweet clover	<i>Melilotus</i> spp.	A/B
Tansymustard	<i>Descurainia pinnata</i>	A
Western ragweed	<i>Ambrosia psilostachya</i>	P
Wild carrot	<i>Daucus carota</i>	B
Wild lettuce	<i>Lactuca</i> spp.	A/B
Wild parsnip	<i>Pastinaca sativa</i>	B
Wild turnip	<i>Brassica campestris</i>	B
Woollyleaf bursage	<i>Franseria tomentosa</i>	P
Yellow woodsorrel	<i>Oxalis stricta</i>	P

Apply 3-4 pints per acre¹

Broom snakeweed ⁴	<i>Gutierrezia sarothrae</i>	P
Bull thistle	<i>Cirsium vulgare</i>	B
Burclover	<i>Medicago</i> spp.	A
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	A
Clover, Hop	<i>Trifolium procumbens</i>	A
Cocklebur	<i>Xanthium strumarium</i>	A
Cudweed	<i>Gnaphalium</i> spp.	A
Desert Camelthorn	<i>Alhagi pseudalhagi</i>	P
Dock	<i>Rumex</i> spp.	P
Fiddleneck	<i>Amsinckia intermedia</i>	A
Goldenrod	<i>Solidago</i> spp.	P
Henbit	<i>Lamium aplexicaule</i>	A
Knotweed, prostrate	<i>Polygonum aviculare</i>	A/P
Pokeweed	<i>Phytolacca americana</i>	P
Purslane	<i>Portulaca</i> spp.	A
Pusley, Florida	<i>Richardia scabra</i>	A
Rocket, London	<i>Sisymbrium irio</i>	A
Rush skeletonweed ⁴	<i>Chondrilla juncea</i>	B
Saltbush	<i>Atriplex</i> spp.	A

13
20

COMMON NAME	SPECIES	GROWTH HABIT ²
Sheperd's-purse	<i>Capsella bursa-pastoris</i>	A
Spurge, Annual	<i>Euphorbia</i> spp.	A
Stinging nettle ⁴	<i>Urtica dioica</i>	P
Velvetleaf	<i>Abutilon theophrasti</i>	A
Yellow starthistle	<i>Centaurea solstitialis</i>	A

Apply 4-6 pints per acre¹

Arrowwood	<i>Pluchea sericea</i>	A
Canada thistle	<i>Cirsium arvense</i>	P
Giant ragweed	<i>Ambrosia trifida</i>	A
Grey rabbitbrush	<i>Chrysothamus nauseosus</i>	P
Little mallow	<i>Malva parviflora</i>	B
Milkweed	<i>Asclepias</i> spp.	P
Primrose	<i>Oenothera kunthiana</i>	P
Silverleaf nightshade	<i>Solanum eleagnifolium</i>	P
Sowthistle	<i>Sonchus</i> spp.	A
Texas thistle	<i>Cirsium texanum</i>	P

VINES AND BRAMBLES

Apply 1 pint per acre

Field bindweed	<i>Convolvulus arvensis</i>	P
Hedge bindweed	<i>Calystegia sequium</i>	A

Apply 2-3 pints per acre¹

Wild buckwheat	<i>Polygonum convolvulus</i>	P
----------------	------------------------------	---

Apply 3-4 pints per acre¹

Greenbriar	<i>Smilax</i> spp.	P
Honeysuckle	<i>Lonicera</i> spp.	P
Morningglory	<i>Ipomoea</i> spp.	A/P
Poison ivy	<i>Rhus radicans</i>	P
Redvine	<i>Brunnichia cirrhosa</i>	P
Wild rose	<i>Rosa</i> spp.	P
Including:		P
Multiflora rose	<i>Rosa multiflora</i>	
McCartney rose	<i>Rosa bracteata</i>	P

Apply 4-6 pints per acre¹

*Kudzu ³	<i>Pueraria lobata</i>	P
Trumpet creeper	<i>Campsis radicans</i>	P
Virginia creeper	<i>Parthenocissus quinquefolia</i>	P
Wild grape	<i>Vitis</i> spp.	P

BRUSH SPECIES

Apply 4-6 pints per acre¹

American beech	<i>Fagus grandifolia</i>	P
Ash	<i>Fraxinus</i> spp.	P
Bald cypress	<i>Taxodium distichum</i>	P

COMMON NAME	SPECIES	GROWTH HABIT ²
Bigleaf maple	<i>Acer macrophyllum</i>	P
Black locust ⁵	<i>Robinia pseudoacacia</i>	P
Black gum	<i>Nyssa sylvatica</i>	P
Box elder	<i>Acer negundo</i>	P
Cherry	<i>Prunus</i> spp.	P
Chinaberry	<i>Melia azadarach</i>	P
Dogwood	<i>Cornus</i> spp.	P
Elm ⁶	<i>Ulmus</i> spp.	P
Hawthorn	<i>Crataegus</i> spp.	P
Hickory	<i>Carya</i> spp.	P
Honeylocust ⁵	<i>Gleditsia triacanthos</i>	P
Maple	<i>Acer</i> spp.	P
Mulberry	<i>Morus</i> spp.	P
Oak	<i>Quercus</i> spp.	P
Persimmon	<i>Diospyros virginiana</i>	P
*Pine ⁵	<i>Pinus</i> spp.	P
Poplar	<i>Populus</i> spp.	P
Privet	<i>Ligustrum vulgare</i>	P
Red Alder	<i>Alnus rubra</i>	P
Red Maple	<i>Acer rubrum</i>	P
Russian Olive	<i>Eleagnus angustifolia</i>	P
Sassafras	<i>Sassafras albidum</i>	P
Sourwood	<i>Oxydendrum arboreum</i>	P
Sweetgum	<i>Liquidambar styraciflua</i>	P
*Water willow	<i>Justica americana</i>	P
Willow	<i>Salix</i> spp.	P
Yellow poplar	<i>Liriodendron tulipifera</i>	P

¹The higher rates should be used where heavy or well-established infestations occur.

²Growth Habit – A = Annual, B = Biennial, P = Perennial

³Use a minimum of 75 GPA - Control of established stands may require repeat applications.

⁴For best results early postemergence applications are required.

⁵Tank mix with glyphosate or triclopyr.

⁶Tank-mix with glyphosate.

*Not approved for use in California.

ADJUVANTS

For optimal postemergence performance of **Alligare Ecomazapyr 2 SL**, the addition of an adjuvant to the spray solution is essential to aid in the deposition and uptake of the herbicide. For this purpose, **ONLY** use spray adjuvants that are approved or appropriate for aquatic use.

Nonionic Surfactants: Use a nonionic surfactant at 0.25% v/v or higher (depending on surfactant manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product. Do not use alcohols, fatty acids, horticultural spray oils, ethylene glycol or diethylene as surfactants to meet these requirements.

Methylated Seed Oils or Vegetable Oil Concentrates: Methylated seed oil or vegetable oil concentrate may be used at 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, mix methylated seed oil or vegetable oil concentrate at a rate of 1% of the total spray volume.

Silicone Based Surfactants: Silicone-based surfactants allow greater spreading of the spray droplet on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based

surfactants may dry too quickly and limit herbicide uptake. Refer to the surfactant manufacturer's label for specific directions.

Invert emulsions: Alligare Ecomazapyr 2 SL can be applied as an invert emulsion (water-in-oil emulsion) designed to minimize spray drift and spray run-off, thereby delivering more herbicide to the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Refer to the invert chemical label for proper mixing directions.

Other: Anti-foam agents, spray pattern indicators or drift reduction agents may also be used if necessary or desired. Refer to the adjuvant manufacturer's label for specific directions.

AQUATIC USES, PRECAUTIONS AND RESTRICTIONS

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. Alligare Ecomazapyr 2 SL may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water.

DO NOT apply more than 6 pints of product (1.5 pounds acid equivalent) per acre per year.

Aerial application is restricted to helicopter equipment only.

Alligare Ecomazapyr 2 SL can only be applied by federal or state agencies, such as Water Management District personnel, municipal officials and the U.S. Army Corps of Engineers, or those applicators who are licensed or certified as aquatic pest control applicators and are authorized by the state or local government.

Treatment of other-than-non-native, invasive species is limited to those that have been determined to be nuisance vegetation by a federal or state government entity.

Applications to private waters: Alligare Ecomazapyr 2 SL may be applied to private non-flowing waters, such as ponds, lakes and drainage ditches where there is minimal or no outflow to public waters.

Applications to public waters: Alligare Ecomazapyr 2 SL may be applied to public waters such as ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers and other slow-moving or non-moving waters for control of aquatic weeds or of riparian and wetland weed species.

Consult local state fish and game agency and water control authorities before applying this product to any public waters. Permits may be required to treat such water.

Recreational Use of Water in Treated Areas: There are no restrictions on the use of water in treated areas for recreational purposes, including swimming and fishing.

Livestock Use of Water in/from Treated Areas:
There are no restrictions on livestock consumption of water from treated areas.

Precautions for Potable Water Intakes: **DO NOT** apply Alligare Ecomazapyr 2 SL directly to waters within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within one-half mile of active potable water intakes, the water intake must be turned off during application and for at least 48 hours after the treatment. Such aquatic applications may be made only where there are alternative water sources or holding ponds that can be used while active potable water intakes are turned off during the minimum 48 hours post-treatment period. **NOTE:** Existing potable water intakes which are no longer in use, such as those replaced by connections to wells or municipal water systems, are not considered to be active potable

water intakes. This restriction does not apply to intermittent, inadvertent overspray of water in terrestrial use sites.

APPLICATION TO WATERS USED FOR IRRIGATION

DO NOT use water treated with **Alligare Ecomazapyr 2 SL** for irrigation purposes for 120 days after treatment or until **Alligare Ecomazapyr 2 SL** residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Seasonal Irrigation Waters: Alligare Ecomazapyr 2 SL may be applied during the off-season to surface waters that are used for irrigation on a seasonable basis, provided at least 120 days elapses between **Alligare Ecomazapyr 2 SL** treatment and the first use of treated water for irrigation or until **Alligare Ecomazapyr 2 SL** residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Irrigation Canals/Ditches: DO NOT apply **Alligare Ecomazapyr 2 SL** to irrigation canals/ditches unless the 120-day restriction on irrigation water usage can be observed or **Alligare Ecomazapyr 2 SL** residue levels are determine by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less. **DO NOT** apply **Alligare Ecomazapyr 2 SL** to dry irrigation canals/ditches.

Quiescent or Slow Moving Waters: In lakes and reservoirs **DO NOT** apply **Alligare Ecomazapyr 2 SL** within one (1) mile of an active irrigation water intake during the irrigation season. Applications less than one (1) mile from an inactive irrigation water intake may be made during the off-season, provided the irrigation intake will remain inactive for at least 120 days after treatment or until **Alligare Ecomazapyr 2 SL** residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Moving Waters: DO NOT apply **Alligare Ecomazapyr 2 SL** within one-half mile downstream of an active irrigation water intake. When making applications upstream from an active irrigation water intake, the intake must be turned off for a period of time sufficient to allow the upstream treated water to completely flow past the irrigation intake. Shut off time will be determined by the speed of water flow and the distance and length of water treated upstream from the intake. Consult local, state and/or federal authorities before making any applications upstream from an active irrigation water intake.

TANK MIXES

Alligare Ecomazapyr 2 SL may be tank-mixed with other aquatic-use herbicides to broaden the spectrum of control of emergent and floating aquatic vegetation provided that the tank mix product label does not prohibit such mixing. Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank-mixes.

AERIAL APPLICATIONS

Exercise all precautions to minimize or eliminate spray drift. Aerial applications may only be made by helicopter; applications may not be made by airplane. Helicopters can be used to apply **Alligare Ecomazapyr 2 SL**; however, **DO NOT** apply by helicopter unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or where damage to desirable vegetation can be tolerated. Helicopters equipped with a Microfoil™ boom, Thru-Valve™ boom or raindrop nozzles must be used and calibrated. Unless applying with a Microfoil boom, use a drift control agent at the specified label rate. To avoid drift, **DO NOT** make applications during inversion conditions, when winds are gusty, or during any other conditions that promote spray drift. Do not use side trimming with **Alligare Ecomazapyr 2 SL** unless death of treated vegetation is acceptable.

Uniformly apply **Alligare Ecomazapyr 2 SL** in 5 to 30 gallons of water per acre. Use a nonionic surfactant, methylated seed oil or silicone-based surfactant (see the **ADJUVANTS** section of this label for specific directions). An anti-foam agent may be added, if needed.

Thoroughly clean application equipment, including landing gear, by thoroughly flushing with water immediately after using this product. Prolonged exposure of uncoated/unpainted steel (except stainless steel) surfaces to this product may result in corrosion and failure of the exposed part. Maintaining painted surfaces may prevent corrosion.

GROUND APPLICATIONS

FOLIAR APPLICATIONS:

For Low Volume Foliar Applications: Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. Thoroughly mix 0.5 to 5% (v/v) **Alligare Ecomazapyr 2 SL** in water plus surfactant (see the **ADJUVANTS** section of this label for directions). Use an anti-foam agent at the specified rate, if needed. For difficult to control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weed species), apply the higher concentrations of herbicide and/or spray volumes but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Do not excessively wet foliage. See the **SPRAY SOLUTION MIXING GUIDE** below for some suggested volumes of **Alligare Ecomazapyr 2 SL** and water.

Select appropriate nozzles to avoid over-application. Proper application is critical to ensure desirable results. Optimum results are achieved when the spray covers the crown and approximately 70 percent of the plant. The use of a flat fan nozzle tip with a spray angle of 40 degrees or less will aid in proper deposition.

Specific nozzle tip sizes include 4004E or 1504E. For a straight stream and cone pattern, use adjustable cone nozzles such as 5500 X3 or 5500 X4. Attaching a rollover valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Moisten, but do not drench target vegetation. **DO NOT** spray to run off.

For Low Volume Foliar Applications by Backpacks: For low-growing species, spray downward to cover approximately 70% of the plant and the crown.

For target species 4 to 8 feet tall, apply a directed spray in a smooth vertical motion from the crown upward on at least two sides of the target vegetation, making sure to cover the crown whenever possible.

For target species over 8 feet tall, apply a directed spray in a smooth zig-zag motion from the crown upward on at least two sides of the target brush.

For Low Volume Foliar Applications with Hydraulic Handgun Equipment: Use same technique as described above for **Low Volume with Backpacks**.

For broadcast applications, simulate a gentle rain near the top of target vegetation, allowing spray to penetrate the target foliage and contact the crown without run-off onto understory vegetation. **DO NOT** spray to run-off. Herbicide spray that contacts understory vegetation may result in severe injury or death of understory plants.

SPRAY SOLUTION MIXING GUIDE FOR LOW-VOLUME FOLIAR APPLICATIONS

Amount of spray solution	Desired Concentration (fluid volume)				
	0.5%	0.75%	1%	1.5%	5%
(amount of Alligare Ecomazapyr 2 SL to use)					
1 gallon	0.6 fl oz	0.9 fl oz	1.3 fl oz	1.9 fl oz	6.5 fl oz
3 gallons	1.9 fl oz	2.8 fl oz	3.8 fl oz	5.8 fl oz	1.2 pint
4 gallons	2.5 fl oz	3.8 fl oz	5.1 fl oz	7.7 fl oz	1.6 pint
5 gallons	3.2 fl oz	4.8 fl oz	6.5 fl oz	9.6 fl oz	2 pints
50 gallons	2 pints	3 pints	4 pints	6 pints	10 pints
100 gallons	4 pints	6 pints	8 pints	6 quarts	5 gallons

2 tablespoons = 1 fluid ounce

For High Volume Foliar Applications: For optimum performance when spraying medium to high-density vegetation, use equipment calibrated to deliver up to 100 gallons of finished spray per acre (GPA). Application volumes exceeding 100 gallons GPA may result in excessive spray run-off, causing injury to desirable ground cover species. Thoroughly mix **Alligare Ecomazapyr 2 SL** in water and include a surfactant (see **ADJUVANTS** section for surfactant specifications). Use an anti-foam agent according to its label, if needed. For hard-to-control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weeds), use the higher concentrations of the herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Uniformly cover the foliage of the target vegetation but **DO NOT** apply to run-off.

Side Trimming: **DO NOT** side trim with **Alligare Ecomazapyr 2 SL** unless severe injury or death of the treated vegetation is acceptable. **Alligare Ecomazapyr 2 SL** is readily translocated and can result in the death of the vegetation.

CUT SURFACE TREATMENTS

Alligare Ecomazapyr 2 SL will control undesirable woody vegetation when applied as a water solution to the cambium area of freshly-cut stump surfaces or to cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the spring. Tree injection and cut stem treatments are most effective in late summer and early fall. **DO NOT** over apply solution causing run-off from the cut surface.

NOTE: Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

CUT SURFACE APPLICATIONS WITH DILUTE AND CONCENTRATE SOLUTIONS:

Mix **Alligare Ecomazapyr 2 SL** as either a concentrated or dilute solution for cut surface treatments. Apply dilute solutions to the cut surface of the stump or to cuts on the stem of the target woody vegetation. Apply concentrate solutions to cuts on the stem. Use of concentrated solutions permits application to fewer cuts on the stem, especially for large diameter trees. Follow the application directions below to determine proper application techniques for each type of solution.

To prepare a dilute solution, mix 8 to 12 fluid ounces of **Alligare Ecomazapyr 2 SL** with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be added according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve herbicide uptake through partially callused cambium tissues.

To prepare a concentrated solution, mix 2 quarts of **Alligare Ecomazapyr 2 SL** with no more than 1 quart water.

For cut stump treatments: Spray or brush the solution onto the cambium area of the freshly cut stump surface. Thoroughly wet the entire cambium area (the wood just inside the bark of the stump).

For cut stem (injection, hack and squirt) treatments:

Dilute Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Ensure that the injector completely penetrates the bark at each injection site.

Concentrate Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut while a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site, place the injection cuts at approximately equal intervals around the tree.

Frill or girdle treatments: Use a hatchet, machete or chainsaw to make cuts through the bark around the tree to expose the cambium. The cuts must angle downward into the cambium sufficiently deep to expose at least two growth rings. Use a spray applicator or brush to apply a 25% to 100% solution of **Alligare Ecomazapyr 2 SL** into each cut until thoroughly wet. Avoid applying so much herbicide that runoff to the ground or water occurs.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: DO NOT store below 10°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable container: Refill this container with imazapyr only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. For final disposal, offer for

20
20

recycling or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT: Read the entire **DIRECTIONS FOR USE** and the **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Microfoil is a trademark of Rhone Poulenc Ag. Company.
Thru-Valve is a trademark of Waldrum Specialties.

[EPA approval date]