

81927-32

04/20/2009

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND  
TOXIC SUBSTANCES

Michael Kellogg  
Pyxis Regulatory Consulting, Inc.  
4110 136<sup>th</sup> St. NW  
Gig Harbor, WA 98332

4.20.09

Subject: EPA Reg. 81927-32 / Alligare VOLT Herbicide Label Amendment

Dear Mr. Kellogg:

The labeling referred to above as amended is acceptable provided you make the following changes:

1. The size of KEEP OUT OF REACH OF CHILDREN and CAUTION must be changed to be in compliance with 40 CFR 15.10.
2. Under the product name on page 1 and page 9, remove "Not for homeowner use".
3. On page 7 and 8, change "rate recommendations" to "rate directions".
4. On page 9, first paragraph change "such as" to "including" and delete "other similar industrial noncrop areas".

Submit one copy of the revised final printed label for the record. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. A stamped copy of the label is enclosed for your records. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Jim Tompkins".

Jim Tompkins  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

2  
12

# Alligare VOLT Herbicide

(Not for homeowner use)

**ACTIVE INGREDIENTS:**

Pyraflufen ethyl (ethyl 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorophenoxyacetate .....	0.16%
Glyphosate (N-(phosphonomethyl)glycine, in the form of it's isopropylamine salt) .....	30.00%
<b>OTHER INGREDIENTS:</b> .....	<b>69.84%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

Contains 0.015 lb pyraflufen ethyl and 2.8 lbs glyphosate acid equivalent per U.S. gallon

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain to you in detail.)

FIRST AID	
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing.

EPA Reg. No. 81927-32

EPA Est. No.

**Manufactured For:**  
Alligare, LLC  
13 N. 8<sup>th</sup> Street  
Opelika, AL 36801

Net Contents:

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**4-20-09**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

81927-32

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant (such as nitrile or butyl) gloves
- Shoes plus socks
- Protective eyewear
- For overhead exposure, wear chemical resistant headgear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove and wash contaminated clothing before reuse.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. This product may contaminate water through drift of spray in wind or via runoff events. Use care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift from treated areas.

### SPRAY DRIFT

Avoid spray drift to all other crops and nontarget areas. Do not apply when weather conditions may cause drift. Do not allow this product to drift onto nontarget areas. Drift may result in illegal residues or injury to adjacent crops and vegetation, in the form of leaf yellowing and defoliation. To avoid spray drift, DO NOT apply aerially when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet size will also reduce spray drift.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Droplet size, boom height, and wind speed are the primary factors determining drift. The specific application conditions required for the use of this product are described below.

#### Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under

unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

### **Controlling Droplet Size**

**Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure** – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation** – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

**Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Maintenance of Nozzles** – Periodic inspection and subsequent replacement of nozzles to ensure proper chemical application is recommended.

### **Boom Length**

For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

### **Application Height**

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

### **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Applications must be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### **Temperature Inversions**

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light and variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant (such as nitrile or butyl) gloves
- Shoes plus socks
- Protective eyewear

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

For other uses, including interiorscapes and other nonagricultural uses, do not enter treated areas without protective clothing until sprays have dried.

**INFORMATION**

Alligare VOLT Herbicide is designed for nonselective grass and broadleaf weed control. This product contains two active ingredients; one of which provides systemic grass and broadleaf control and one which provides contact activity against broadleaf weeds.

Only certified applicators are permitted to apply Alligare VOLT Herbicide for turf and ornamental sites. This product is nonselective and should not be applied to sensitive crops or foliage, including turfgrass.

**USE RESTRICTIONS**

- Do not apply more than 32 fl. oz./acre for preplant burndown use prior to seeding crops listed on this label (corn, wheat, soybean, cotton, root and tuber vegetables, leafy vegetables, cole crops, legumes, fruiting vegetables, cucurbits, and small grains).
- Do not apply more than 76 fl. oz./acre for preplant fallow and crop stubble.
- Do not apply more than 76 fl. oz./acre for nonbearing and bearing deciduous fruit and nut trees and vines.
- Do not apply more than 76 fl. oz./acre for noncrop land and uncultivated agricultural areas per year.
- Do not apply more than 96 fl. oz./acre per growing season for noncrop weeds per year.
- Do not apply this product through any type of irrigation system.

**WEEDS CONTROLLED**

The following broadleaf and grass weed species can be controlled by applications of Alligare VOLT Herbicide at rates of 24-38 fl. oz. per acre. Tankmixes of Alligare VOLT Herbicide with other herbicides may be needed for control of these weed species if larger than 4 inches tall or rosettes of greater than 3 inches in diameter.

Barley Bedstraw Beggartick, hairy Bindweed, field (suppression) Bittercress Bluegrass, annual Bluegrass, bulbous Brome, downy Brome, Japanese Browntop panicum Cheatgrass Cocklebur Corn speedwell	Crabgrass Dwarf dandelion Fall panicum Field pennycress Fleabane, annual Fleabane, rough Foxtail, giant Foxtail, green Foxtail, yellow Goatgrass, jointed Grain sorghum Johnsongrass, seedling Little Barley	London rocket Mustard, blue Mustard, tansy Mustard, tumble Mustard, wild Sandbur, field Shepherd's-purse Sprangletop Sunflower, common Texas panicum Wild Oats
--	--	--

The following broadleaf and grass weed species can be controlled by applications of Alligare VOLT Herbicide at rates of 32-38 fl. oz. per acre. Tankmixes of Alligare VOLT Herbicide with other herbicides may be needed for control of these weed species if larger than 4 inches tall or rosettes of greater than 3 inches in diameter.

Anoda, spurred Barnyardgrass Beggarweed, Florida Buckwheat, wild Bur cucumber Canola Carolina geranium Carpetweed Celery, wild Chickweed Copperleaf, hornbeam Copperleaf, Virginia Culteaf Evening Primrose Dandelion Dock, curly Eclipta False dandelion Fiddleneck Filaree Fleabane, hairy Florida pusley	Goosegrass Groundcherry Groundsel, common Hemp sesbania Henbit Marestalk (Horseweed) Jimsonweed Knotweed Kochia Lambsquarters Mallow, common Morningglory Nightshade, black Nightshade, hairy Nettle, stinging Pigweed (Palmer Amaranth) Pigweed species Poinsettia, wild Poison-ivy Prickly lettuce Purslane Mayweed	Ragweed, common Ragweed, giant Sicklepod Signalgrass, broadleaf Smellmelon Smartweed (ladysthumb) Smartweed, Pennsylvania Sowthistle, annual Spurge, leafy (suppression) Spurge, prostrate Spurge, spotted Swinecress Prickly sida Thistle, Russian Toadflax, dalmation Velvetleaf Virginia pepperweed Waterhemp, tall Wild proso millet Woolly cupgrass Yellow rocket
---	--	--

### MIXING DIRECTIONS

Add ½ to ¾ of the required amount of water to the spray tank. Start agitation. Add the required amount of Alligare VOLT Herbicide and the remaining amount of water. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

**Use an approved agricultural buffering agent buffering to pH 7.0 or less if using Alligare VOLT Herbicide in a water source of greater than pH 7.0.**

### TANK MIXTURES

Alligare VOLT Herbicide may be applied as a tankmix or in sequential application with other herbicide products. Weather, crop conditions, or the presence of certain weeds, crop damaging insects, or diseases will indicate the inclusion of other pesticides in the defoliation or desiccation application.

**Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.**

**Note:** It is recommended that the compatibility of Alligare VOLT Herbicide in any tankmix combination be tested before use. To determine the physical compatibility with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

### EQUIPMENT CLEANING

**Do not allow the spray solution to dry in the application equipment. After application and before using the sprayer equipment for any other applications, the sprayer must be thoroughly cleaned. Applicators must ensure proper equipment clean-out for any other products mixed with Alligare VOLT Herbicide as provided on the other product label(s). Immediately following application, clean all equipment thoroughly with detergent or a spray tank cleaner and water as described below. Should residues of Alligare VOLT Herbicide remain in inadequately cleaned equipment, they may be released in subsequent applications and cause injury to crops.**

1. Drain sprayer tank, hoses, and spray boom and thoroughly rinse with clean water the inside of the spray tank, sprayer hoses, boom, and nozzles to remove any sediment or residues.
2. Fill the tank ½ full with clean water, add the appropriate detergent (follow manufacturer's directions for use). Fill tank to capacity and operate the sprayer with agitation for 15 minutes to flush hoses, boom, and nozzles.
3. Drain the sprayer tank, lines, and booms. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray nozzles, tips, and screens.
4. Dispose of all cleaning solutions, rinsate, and washwaters in accordance with Federal, state, and local regulations.

### ROTATIONAL CROP RESTRICTIONS

Do not plant rotational crops other than crops listed on this label for at least 30 days after application.

8  
12

### APPLICATION AND DOSAGE

#### PREPLANT BURNDOWN

Alligare VOLT Herbicide may be used for weed control preplant to the following crops:

**Field Corn, Wheat, Soybeans, Cotton, Root and Tuber Vegetables, Leafy Vegetables, Cole Crops, Legumes, Fruiting Vegetables, Cucurbits, and Small Grains (Limited to Preplant Burndown)**

Thorough, uniform spray coverage is essential for good control. Alligare VOLT Herbicide may be applied preplant burndown to control weeds or in tank mixtures with other labeled herbicides.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Crop	Pest	Rate/Acre	Use Restrictions and Comments
Field Corn, Soybeans, Wheat and Cotton	Annual grass and/or broadleaf weeds	24-32 fl. oz./A in a minimum of 5 gallons water per acre by air or 10 gallons water per acre by ground*.	<ul style="list-style-type: none"> <li>▪ Use the higher rate and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>▪ Allow a minimum of 30 days between applications.</li> <li>▪ Treated areas may be replanted immediately with any crop listed on this label.</li> <li>▪ Do not plant any other rotational food crops for 30 days after the last application.</li> <li>▪ Do not apply more than 32 fl. oz./A for this use.</li> </ul>
Root and Tuber Vegetables, Leafy Vegetables, Cole Crops, Legumes, Fruiting Vegetables, Cucurbits, Small Grains			<ul style="list-style-type: none"> <li>▪ Use the higher rate and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>▪ Allow a minimum of 30 days between applications.</li> <li>▪ For crops listed on this label, do not apply within 2 days of planting.</li> <li>▪ Do not plant any other rotational food crops for 30 days after the last application.</li> <li>▪ Do not apply more than 32 fl. oz./A for this use.</li> </ul>
*Refer to Weeds Controlled section for specific rate recommendations. Alligare VOLT Herbicide may be tank mixed with other labeled herbicides, including residual products for expanded weed control.			



**PREPLANT FALLOW BEDS AND CROP STUBBLE**

This product may be applied preplant to fallowland in preparation for planting or postharvest to crop stubble. Preplant applications may be made prior to planting during the fallow period for any crop listed on this label. For crops not listed on this label, applications must be made at least 30 days prior to planting.

Thorough, uniform spray coverage is essential for good control. Alligare VOLT Herbicide may be applied after the harvest of any crop to control late emerging broadleaf weeds or in tank mixtures with other labeled herbicides for broad spectrum weed control.

Addition of a nonionic surfactant or crop oil concentrate (COC) is recommended for optimum control. Follow manufacturer's use rates.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Crop	Pest	Rate/Acre	Use Restrictions and Comments
Preplant fallowbeds and crop stubble	Annual grass and/or broadleaf weeds	24-32 fl. oz./A in a minimum of 10 gallons water per acre by ground; 5 gallons water per acre by air*	<ul style="list-style-type: none"> <li>• Use the higher rate and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>• Allow a minimum of 30 days between applications.</li> <li>• For crops not listed on this label, applications must be made at least 30 days prior to planting.</li> </ul>
*Refer to Weeds Controlled section for specific rate recommendations.			

**NONCROP LAND AND UNCULTIVATED AGRICULTURAL AREAS (nonfood producing)**

Alligare VOLT Herbicide may be used for broad spectrum weed control in noncrop situations. Thorough, uniform spray coverage is essential for good control.

Addition of a nonionic surfactant or crop oil concentrate (COC) is recommended for optimum control. Follow manufacturer's use rates.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Crop	Pest	Rate/Acre	Use Restrictions and Comments
Noncrop Land and Uncultivated Agricultural Areas	Annual grass and/or broadleaf weeds	24-32 fl. oz./A plus other labeled herbicides in a minimum of 5 gallons water per acre by air or 10 gallons water per acre by ground	<ul style="list-style-type: none"> <li>• Use the higher rates and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>• Allow a minimum of 30 days between applications.</li> </ul>

10  
12

**NONCROP WEED CONTROL (Not for homeowner use)**

For use in noncrop areas where control of weeds is desired, such as airports; commercial plants; storage and lumber yards; barrier strips and firebreaks; equipment areas; nurseries and ornamental plantings; sodfarms; Christmas trees; established ornamental turf; railroad, roadside and utility rights-of-way; fuel tank farms and pumping stations; other similar industrial noncrop areas.

For applications to ornamental turf and plantings, do not allow people (other than the applicator) or pets on treatment area during application and until sprays have dried (refer to Nonagricultural Use Requirements box). Apply Alligare VOLT Herbicide at rates specified in the dosage table below for control of broadleaf and grass weeds. **Avoid contact with desirable vegetation.**

Addition of a nonionic surfactant or crop oil concentrate (COC) is recommended for optimum control. Follow manufacturer's use rates.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Use	Rate/Acre	Use Restrictions and Comments
(See directions for use above for explanation of appropriate use sites)	24-38 fl. oz./A in 20 to 40 gpa using ground or backpack or similar spray equipment or 5 gpa by air	<ul style="list-style-type: none"> <li>Do not make more than 3 applications.</li> <li>Allow a minimum of 30 days between applications.</li> </ul>

**NONBEARING DECIDUOUS FRUIT AND NUT TREES AND VINES (EXCLUDING CITRUS)**

Thorough, uniform spray coverage is essential for adequate control.

Addition of a nonionic surfactant or crop oil concentrate (COC) is recommended for optimum control. Follow manufacturer's use rates.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Crop	Pest	Rate/Acre	Comments and Restrictions
Nonbearing Tree Fruit, Nut, and Vine crops	Annual grasses and/or broadleaf weeds	24-32 fl. oz./A in a minimum of 10 gallons of water per acre by ground	<ul style="list-style-type: none"> <li>Use the higher rate and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>Allow a minimum of 30 days between applications.</li> <li>Do not harvest edible crops for 12 months following application.</li> <li>Addition of labeled residual herbicides to extend weed control is permissible.</li> <li>Do not apply by air.</li> </ul>

11  
12

**DECIDUOUS FRUIT AND NUT TREES AND VINES (EXCLUDING CITRUS)**  
**(Postharvest, Dormant and Prebloom applications)**

[dates, feijoa, figs, grapes, kiwi fruit, mango, olives, persimmons, pome fruit, pomegranates, stone fruit, and tree nuts] (excluding citrus)

Alligare VOLT Herbicide may be applied as a postharvest or preplant burndown treatment for control of emerged winter and summer annual broadleaf and grass weeds and burndown or suppression of certain perennial broadleaf and grass weeds during the dormant period prior to bloom. Thorough coverage of target weeds is essential for optimum performance. Addition of a nonionic surfactant or crop oil concentrate (COC) is recommended for optimum control. Follow manufacturer's use rates. Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

Crop	Pest	Rate/Acre	Use Restrictions and Comments
Dates, Feijoa, Figs, Grapes, Kiwi Fruit, Mango, Olives, Persimmons, Pome and Stone Fruit, Pomegranate Tree Nuts	Annual grass and broadleaf weeds	24-32 fl. oz./A in a minimum of 10 gallons water per acre in a broadcast or band directed application	<ul style="list-style-type: none"> <li>• Use the higher rate and spray volumes for control of larger weeds (4" tall). Control may be reduced with weeds larger than 4 inches tall.</li> <li>• Allow a minimum of 30 days between applications.</li> <li>• Do not allow spray to contact green bark of trunk area on young grape vines and fruit or nut trees.</li> <li>• Do not apply by air.</li> </ul>

**STORAGE AND DISPOSAL**

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** Store in a cool place.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Nonrefillable > 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

12  
12

**CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

**Warranty:** Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

**Terms of Sale:** The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

**Limitation of Liability:** To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

EPA [approval date]