



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 26, 2026

Bonnie Bieber
bonnie.bieber@alligare.com
ALLIGARE, LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Adding Subset B section into label because it was mistakenly absent from the label that was approved on January 7, 2026.
Product Name: ALLIGARE FLUMI SC
Admin Number: 81927-78
EPA Receipt Date: 02/09/2026
Action Case Number: 00684358

Dear Bonnie Bieber:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Jeffrey Chang via email at chang.jeffrey@epa.gov.

Sincerely,

Kable Bo Davis

Kable Bo Davis, Senior Advisor
FHB, RD
Office of Pesticide Programs

[Note to reviewer: [Text] in brackets denotes optional text].
[Note to reviewer: {Text} in braces denotes where in the final label text will appear].

FLUMIOXAZIN	GROUP	14	HERBICIDE
--------------------	--------------	-----------	------------------

Alligare FLUMI SC

[Alternate Brand Names: Alligare FLUMIGARD® SC Herbicide; PROMENADE® SC Herbicide; PROMENADE® SC Specialty Herbicide]

- **[Sublabel A – Aquatics]** – FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC VEGETATION IN SLOW MOVING OR QUIESCENT WATERS
- **[Sublabel B – Non-Crop/IVM]** – FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS, CONIFER AND POPLAR RE-FORESTATION SITES

[Sublabel C – Aquatics, Non-Crop/IVM, and T&O] – [FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC VEGETATION IN SLOW MOVING OR QUIESCENT WATERS] [FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS, CONIFER AND POPLAR RE-FORESTATION SITES] [FOR USE IN CONTAINER AND FIELD GROWN CONIFERS (INCLUDING CHRISTMAS TREES) AND DECIDUOUS TREES, AROUND ESTABLISHED WOODY ORNAMENTALS IN LANDSCAPES AND MAINTAIN NON-CROP AREAS AND DORMANT BERMUDAGRASS]

ACTIVE INGREDIENT:

Flumioxazin* 42.0%

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*[2-[7-flouro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione]

Alligare FLUMI SC contains 4 pounds flumioxazin per gallon.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)


[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

[For ≤ 5 Gallon Containers:] [Shake Well Before Use]
[For ≥ 5 Gallon Containers:] [Shake Well, Agitate or Recirculate Before Use]

EPA Reg. No. 81927-78

Manufactured for:
Alligare, LLC
PO Box 97
Deer Park, TX 77536

EPA Est. No.



Net Contents:

[SUBLABEL A]

[Note to reviewer: [Text] in brackets denotes optional text].

[Note to reviewer: {Text} in braces denotes where in the final label text will appear].

{[BOOKLET FRONT PANEL LANGUAGE]}

FLUMIOXAZIN	GROUP	14	HERBICIDE
-------------	-------	----	-----------

Alligare FLUMI SC

[Alternate Brand Names: Alligare FLUMIGARD® SC Herbicide; PROMENADE® SC Herbicide; PROMENADE® SC Specialty Herbicide]

FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC VEGETATION IN SLOW MOVING OR QUIESCENT WATERS

ACTIVE INGREDIENT:

Flumioxazin* 42.0%

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*(2-[7-flouro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione)

Alligare FLUMI SC contains 4 pounds flumioxazin per gallon.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

[For ≤ 5 Gallon Containers:] [Shake Well Before Use]

[For ≥ 5 Gallon Containers:] [Shake Well, Agitate or Recirculate Before Use]

EPA Reg. No. 81927-78

EPA Est. No.

Manufactured for:

Alligare, LLC
PO Box 97
Deer Park, TX 77536

Net Contents:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes and socks

Mixers, loaders and applicators making aquatic subsurface applications using a mechanically-pressurized handgun must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

If not used in accordance with directions on the label, this product can be toxic to non-target plants and aquatic invertebrates. Do not apply directly to treated, finished drinking water reservoirs or drinking water receptacles when the water is intended for human consumption. Drift and runoff may be hazardous to non-target plants and aquatic organisms in water adjacent to treated areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants. Use strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard, treat 1/3 to 1/2 of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is needed.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System

(NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions, and with applicable state and federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as “Buyer”) of this product must be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Alligare. The Buyer must be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that the additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, **THEN DO NOT APPLY THIS PRODUCT**. By applying this product Buyer acknowledges and accepts these inherent unintended risks **AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER**.

Alligare shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

See also **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** sections of the label for additional information.

WEED RESISTANCE MANAGEMENT

This product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population

if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Avoid the consecutive use of Alligare FLUMI SC or other target site of action Group 14 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Alligare LLC retailer, representative or call 888-252-4427. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PRODUCT INFORMATION

This product is a fast-acting contact herbicide that controls selected submersed, emergent, and floating aquatic weeds. It is most effective when applied to young, actively growing weeds in water with a pH of less than 8.5.

This product may be applied to the following quiescent or slow-moving bodies of water:

- Bayous
- Canals
- Drainage ditches
- Lakes
- Marshes
- Ponds (including golf course ponds)
- Reservoirs

Application of this product to public aquatic areas may require special approval and/or permits. Consult with local state agencies, if required.

USE RESTRICTIONS

- Do not apply to intertidal or estuarine areas.
- Do not exceed 400 ppb of this product during any one application.
- Do not re-treat the same section of water with this product more than 6 times per year.
- Do not retreat the same section of water within 28 days of application, except in areas with dense weed vegetation. In these areas, treat the remaining weeds within 10 to 14 days.
- In high density weed populations only treat 1/2 the water body at one time.
- Treated water may not be used for irrigation purposes on food crops until at least five (5) days after application.
- Do not use in water utilized for crawfish farming.

USE PRECAUTIONS

- There is no post-application holding restriction against use of treated water for drinking or recreational purposes (e.g., swimming, fishing).
- Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the *Irrigation Restrictions Following Application* table.

IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

Application Method	Application Rate	Average Water Depth	Turf and Landscape Ornamentals	Ornamentals grown for production in Greenhouse and Nursery
Surface Spray	6 to 12 oz per surface acre	Greater than 3 feet	None	5 days
		Less than 3 feet	12 hours	5 days
Subsurface	Less than 200 ppb	N/A	1 day	5 days
	200 to 300 ppb	N/A	2 days	5 days
	300 to 400 ppb	N/A	3 days	5 days

SPRAY DRIFT MANAGEMENT

Aerial Applications

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the application site. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the application site.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or vegetative canopy.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.Controlling Droplet Size - Aircraft
 - Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
For ground equipment, the boom should remain level with the target area and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

APPLICATION AND SPRAYER INFORMATION

Mixing Instructions

- Mix with water having pH of 5 to 7. If pH is higher than 7, use an appropriate buffer to reduce pH to desirable range.
- Fill clean spray tank 1/2 full of desired level with water and add buffering agent if necessary.
- Add the required amount of this product to the spray tank while agitating.
- Fill spray tank to desired level with water. Ensure that this product is thoroughly mixed before making applications. Continue agitation until spray solution has been applied.
- Mix only the amount of spray solution that can be applied the day of mixing. Apply this product within 48 hours of mixing.

ADDITIVES

When applying this product to the foliage of floating or emerged aquatic weeds, mix with an adjuvant approved for use in aquatic sites. Mix this product with a non-ionic surfactant containing at least 80% active ingredient. Follow adjuvant manufacturer's label rates. Verify mixing compatibility by a jar test before using.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND THIS PRODUCT

Perform a jar test before mixing commercial quantities of this product, when using this product for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pint of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 milliliter of this product to the quart jar for every 3 fl oz of this product per acre being applied (4 ml if 12 fl oz per acre is the desired rate of this product), gently mix until product goes into suspension.
3. Add 1 milliliter on non-ionic surfactant, gently mix.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An ideal tank mix combination will be uniform. If any of the following conditions are observed question the choice of adjuvant:
 - a) Layer of oil or globules on the mixture's surface,
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar,
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

Sprayer Cleanup

If spray equipment is dedicated to application of aquatic herbicides, the following steps are to clean the spray equipment:

- Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.

If spray equipment will be used for purposes other than applying aquatic herbicides, it must be thoroughly cleaned following application of this product. The following steps must be used to clean the spray equipment:

1. Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank with clean water.

4. Circulate through sprayer for 5 minutes.
5. Then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
6. Drain tank completely.
7. Remove all nozzles and screens and rinse them with clean water.

AERIAL APPLICATION

To obtain satisfactory weed control, aerial application of this product, must provide uniform coverage of surface weeds and sufficient contact time. When applied by air, this product may not provide adequate control of some submersed weeds. **Do not** apply by air when significant drift on to non-target plants may occur or when wind velocity is more than 10 mph. Avoid spraying this product within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory application and avoid drift, the following direction must be observed:

- **Volume Pressure**
Apply this product in a minimum of 5 gallons of water per acre, with a maximum spray pressure of 40 PSI. Application at less than 5 gallons per acre may not provide adequate weed control. Higher gallonage applications provide more consistent weed control.
- **Nozzles and Nozzle Operation**
Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles including diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. **Do not** place nozzles on the outer 25% of the wings or rotors.
- **Adjuvants**
Refer to the additive section or the tank mix partners label for adjuvant specifications.

DIRECTIONS FOR USE TO CONTROL FLOATING AND EMERGED WEEDS USING SURFACE APPLICATION

This product will control weeds and algae listed in Floating and Emerged Weeds when applied as a broadcast spray with appropriate equipment. For best results, apply this product to the foliage of actively growing weeds.

Floating and Emerged Weeds

Common Name	Scientific Name
Alligator Weed	<i>Alternanthera philoxeroides</i>
Duckweed*	<i>Lemna</i> spp.
Frog's-bit	<i>Limnobium spongia</i>
Mosquito Fern	<i>Azolla</i> spp.
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal*	<i>Wolffia</i> spp.
Water Pennywort	<i>Hydrocotyle</i> spp.
Filamentous algae	<i>Pithophara</i>
Filamentous algae	<i>Cladophora</i>

*Coverage is essential for effective duckweed and watermeal control. Any duckweed and/or watermeal escapes left in the water column will quickly re-infest the water body. Apply 200 ppb concentration throughout the water body to control duckweed and watermeal. – see **DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATIONS** section for additional application information.

Surface Application

Apply this product as a broadcast spray at 6 to 12 fl oz of formulated product (0.19 to 0.38 lb ai) per acre plus an adjuvant approved for use in aquatics.

This product is a contact herbicide that quickly degrades in the water column so plants that do not initially come in contact with the herbicide will not be controlled. Apply this product in a minimum of 30 gallons of water per acre to all areas of the water body where weeds exist. Coverage is essential for effective control as all floating weeds need to be exposed to lethal concentrations in all parts of the water body. Any untreated escapes or re-introductions of plants that were not treated will reestablish in areas where surface weeds had previously been controlled. If a second application is required to provide control, make a treatment once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Application of this product during early morning hours may enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage. Rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat floating surface weeds in section to avoid a rapid decrease in dissolved oxygen.

This product may be tank mixed with 2,4-D, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds.

Consult a manufacturer's label for specific rate restrictions and weeds controlled. Always follow the most restrictive label restrictions and precautions for all products used when making an application involving tank mixes.

Application Equipment

Apply this product with sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Apply by backpack or handgun sprayer, airboat, helicopter, airplane or other application equipment that will ensure thorough coverage of target plant foliage.

DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATIONS

This product will control submersed and floating weeds listed in *Submersed and Floating Weeds Controlled by Subsurface Application*, when applied subsurface with appropriate equipment.

Submersed and Floating Weeds Controlled by Subsurface Application

Common Name	Scientific Name
Coontail	<i>Ceratophyllum demersum</i>
Duckweed	<i>Lemna</i> spp.
Fanwort	<i>Cabomba caroliniana</i>
Hydrilla	<i>Hydrilla verticillata</i>
Hygrophila	<i>Hygrophila polysperma</i>
Naiad, Southern	<i>Najas guadalupensis</i>
Pondweed, Curlyleaf	<i>Potamogeton crispus</i>
Pondweed, Sago	<i>Potamogeton pectinatus</i>
Pondweed, Variable-leaf	<i>Potamogeton diversifolius</i>
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal	<i>Wolffia</i> spp.
Watermilfoil, Eurasian	<i>Myriophyllum spicatum</i>
Watermilfoil, Variable-Leaf	<i>Myriophyllum heterophyllum</i>

Subsurface Treatment

Apply this product at a rate that will produce an initial concentration of 200 to 400 ppb (of active ingredient flumioxazin) in the water column.

This product is rapidly absorbed by target plants, but also breaks down quickly in water with a pH greater than 8.5. The pH of water surrounding mats of submersed vegetation can exceed 8.5 by early to mid-day, due to photosynthetic processes. Application of this product under these conditions may provide only partial weed control, and regrowth is likely. For best control, apply this product in a minimum of 30 gallons of water per acre in the early morning to actively growing weeds and early in the season before surface matting occurs. Complete coverage and sufficient contact time of submersed weeds with this product is required for optimal performance. Application of this product with subsurface trailing hoses designed to distribute the herbicide within the plant stand will provide more effective and longer-term control of submersed weeds. Use *Subsurface Application Rates* to determine the amount of this product needed to achieve desired concentration at different water depths. Use higher concentrations when weed biomass is heavy and/or weeds are more mature and topped out. Any untreated plants that are left in the water column can re-infest treated areas that had previously been controlled. If a second application is required to provide control, make a treatment once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

When applying this product to densely packed actively growing submersed weeds, a rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat submersed weeds in sections to avoid a rapid decrease in dissolved oxygen.

This product may be tank mixed with other registered submersed applied herbicides for enhanced control of submersed and floating weeds.

Application Equipment for Water Column Treatment

To improve distribution in the water column and ensure adequate coverage, when possible apply this product with subsurface trailing hoses in order to place the herbicide under the surface and throughout the biomass of aquatic vegetation. Keep swath width to a minimum in order to maximize contact with submersed aquatic vegetation. In small shallow water bodies, surface sprays may be required to apply this product. Apply by backpack or handgun sprayer or other application equipment that will ensure adequate coverage of target plant.

Information on Hydrilla Control in Florida

Apply this product as a subsurface treatment for hydrilla control. For best control of hydrilla apply during the late Winter/early Spring and/or early to late Fall. Efficacy of this product will be enhanced at these timings due to lower potential biomass present and lower pH of the water. If applied to mature topped out hydrilla, this product will cause some discoloration and loss of growing tips, but regrowth will be rapid.

Tank mixing this product with other registered herbicides is advised, especially if hydrilla is approaching maturity or biomass is heavy.

Subsurface Application Rates

Water Depth (feet)	Pints of This Product Required Per Surface Acre to Achieve Desired Water Concentration		
	200 ppb	300 ppb	400 ppb
1	1.1	1.6	2.1
2	2.1	3.2	4.2
3	3.2	4.8	6.4
4	4.2	6.4	8.5
5	5.3	8.0	10.6

Example: to achieve an initial concentration of 200 ppb of flumioxazin in a 4-foot deep water column, apply 4.2 pints of this product per surface acre.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Store in a cool, dry, secure place. Do not put formulation or dilute spray solution into food or drink containers. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. Not for use or storage in or around the home. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC (800) 424-9300.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Nonrefillable Containers 5 gallons or less:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Nonrefillable containers larger than 5 gallons:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Refillable containers larger than 5 gallons:] [Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose.

To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

[PROMENADE® is a registered trademark of ADAMA group company.]

[FLUMIGARD® is a registered trademark of ADAMA group company.]

[EPA Approval Date]

REV020926

[SUBLABEL B]

[Note to reviewer: [Text] in brackets denotes optional text].

[Note to reviewer: {Text} in braces denotes where in the final label text will appear].

{[BOOKLET FRONT PANEL LANGUAGE]}

FLUMIOXAZIN	GROUP	14	HERBICIDE
-------------	-------	----	-----------

Alligare FLUMI SC

[Alternate Brand Names: Alligare FLUMIGARD® SC Herbicide; PROMENADE® SC Herbicide; PROMENADE® SC Specialty Herbicide]

FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS, †CONIFER AND POPLAR RE-FORESTATION SITES

†Not for use in California

ACTIVE INGREDIENT:

Flumioxazin* 42.0%

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*(2-[7-flouro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione)

Alligare FLUMI SC contains 4 pounds flumioxazin per gallon.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

[For ≤ 5 Gallon Containers:] [Shake Well Before Use]

[For ≥ 5 Gallon Containers:] [Shake Well, Agitate or Recirculate Before Use]

EPA Reg. No. 81927-78

EPA Est. No.

Manufactured for:

Alligare, LLC
PO Box 97
DeerPark, TX 77536

Net Contents:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

If not used in accordance with directions on the label, this product is toxic to non-target plants and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to non-target plants and aquatic organisms in water adjacent to treated areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants. Use strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to run off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, including no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation strips along rivers, creeks, streams, wetlands or on the downhill side of fields where run-off could occur will minimize water run-off.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site.

Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions, and with applicable state and federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural crops on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

Do not enter or allow others to enter the treated area until sprays have dried.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product must be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Alligare. The Buyer must be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that the additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN DO NOT APPLY THIS PRODUCT. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Alligare shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

See also **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** sections of the label for additional information.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PRODUCT INFORMATION

This product is a selective herbicide to maintain bare ground non-crop areas when used in accordance with this label. This product is effective as a preemergence and/or postemergence herbicide for control of selected grass and broadleaf weeds.

This product controls weeds by inhibiting protoporphyrinogen oxidase, an essential enzyme required by plants for chlorophyll biosynthesis. Seedling weeds are controlled preemergence when exposed to sunlight following contact with the soil applied herbicide.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply Alligare FLUMI SC within 30 days.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply to moist or wet desirable plant foliage.
- Do not apply within 300 feet of non-dormant pome or stone fruit crops.

USE PRECAUTIONS

- Treatment of powdery, dry soil or light sandy soil, or light sandy soil when there is little to no likelihood of rainfall soon after may result in off target movement and possible damage to actively growing susceptible crops when soil particles are moved by wind or water. Do not apply when these soil and environmental conditions are present.

WEED RESISTANCE MANAGEMENT

This product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Avoid the consecutive use of Alligare FLUMI SC or other target site of action Group 14 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Alligare LLC retailer, representative or call 888-252-4427. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

PREEMERGENCE APPLICATION

Make the preemergence application of this product prior to weed emergence. Moisture is necessary to activate this product for residual weed control. Moisture is needed to move this product into the soil for preemergence weed control. Dry weather following application of this product may reduce effectiveness.

POSTEMERGENCE APPLICATION

For best results, apply this product to actively growing weeds. Applying this product under conditions that do not promote active weed growth will reduce herbicide effectiveness.

Do not apply this product when the weeds are under stress due to drought, excessive water and extremes in temperatures or disease. This product is most effective when applied under sunny conditions at temperatures above 65° F.

This product is rainfast one hour after application. **Do not** make applications if rain is expected within one hour of application or efficacy may be reduced.

APPLICATION EQUIPMENT

Important: Thoroughly clean spray equipment, including all tanks, hoses, booms, screens and nozzles. **Do not use spray equipment used to apply this product to apply other materials to any desirable plant foliage.** Equipment with this product's residue remaining in the system may result in crop injury to subsequently treated crops.

SPRAYER PREPARATION

Before applying this product, start with clean, well maintained application equipment. Clean the spray tank, as well as all hoses and booms to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to the sulfonylurea and phenoxy herbicides, are active at very small amounts and can cause crop injury when applied to susceptible crops. Clean the spray equipment according to the manufacturer's directions for the last product used before the equipment is used to apply this product. If two or more products were tank mixed prior to this product's application, follow the most restrictive cleanup procedure.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. Agitate solution. Ensure agitation creates a rippling or rolling action on the water surface.
3. If tank mixing this product with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants.
5. Fill spray tank to desired level with water. **Continue agitation until all spray solution has been applied.**
6. Mix only the amount of spray solution that can be applied the day of mixing. Apply this product within 48 hours of mixing.

SPRAYER CLEANUP

If spray equipment is dedicated to herbicide applications, use the following steps are to clean the spray equipment:

- Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.

If spray equipment will be used for purposes other than applying herbicides, it must be thoroughly cleaned following application of this product. Use the following steps to clean the spray equipment:

- Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
- Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
- Top off tank, add suitable commercial spray tank cleaning material, following label directions, or add 1 gallon of 3% household ammonia for every 100 gallons of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
- Drain tank completely.
- Add enough clean water to the spray tank to allow all hoses, booms, screens and nozzles to be flushed for 2 minutes.
- Remove all nozzles and screens and rinse them with clean water.

SPRAY DRIFT MANAGEMENT

Aerial Applications

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the application site. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the application site.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or vegetative canopy.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.Controlling Droplet Size - Aircraft
 - Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
For ground equipment, the boom should remain level with the target area and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

WEEDS CONTROLLED

When this product is applied preemergence or postemergence at directed rates and weed stages, the following grasses and broadleaf weeds are controlled:

WEEDS CONTROLLED BY THIS PRODUCT

COMMON NAME	SCIENTIFIC NAME
Alyssum, Hoary	<i>Berteroa incana</i>
Amaranth Palmer Spiny	<i>Amaranthus palmeri</i> <i>Amaranthus spinosus</i>
American Burnweed	<i>Erechetities hieracifolia</i>
Barnyardgrass*	<i>Echinochloa crus-galli</i>
Beggarweed, Florida	<i>Desmodium Tortuosum</i>
Bittercress, Hairy	<i>Cardamine hirsute</i>
Bluegrass, Annual	<i>Poa annua</i>
Burclover, California	<i>Medicago Polymorpha</i>
Carpetweed	<i>Mollugo verticillata</i>
Chamberbitter	<i>Phyllanthus urinaria</i>
Chickweed Common Mouseear	<i>Stellaria media</i> <i>Cerastium vulgatum</i>
Crabgrass Large* Smooth* Southern*	<i>Digitaria sanguinalis</i> <i>Digitaria ischaemum</i> <i>Digitaria ciliaris</i>
Croton, Tropic	<i>Croton glandulosus var. septentrionalis</i>
Dandelion*	<i>Taraxacum officinale</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Doveweed	<i>Murdannia nudiflora</i>
Eclipta	<i>Eclipta prostrate</i>
Filaree, Redstem*	<i>Erodium cicutarium</i>
Foxtail Bristly* Giant* Green* Yellow*	<i>Setaria verticillata</i> <i>Setaria faberi</i> <i>Setaria viridis</i> <i>Setaria glauca</i>
Galinsoga, Hairy	<i>Galinsoga ciliate</i>
Geranium, Carolina	<i>Geranium carolinianum</i>
Goosegrass*	<i>Eleusine indica</i>
Groundsel, Common	<i>Senecio vulgaris</i>
Groundsel, Tree	<i>Baccharis halimifolia</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed*	<i>Conyza Canadensis</i>
Indigo, Hairy	<i>Indigofera hirsute</i>
Ivy, Ground*	<i>Glechoma hederacea</i>
Jimsonweed	<i>Datura stramonium</i>

COMMON NAME	SCIENTIFIC NAME
Kochia	<i>Kochia scoparia</i>
Kyllinga, Green*	<i>Kyllinga brevifolia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Lovegrass, California*	<i>Eragrostis diffusa</i>
Liverwort	<i>Marchantia polymorpha</i>
Mallow Common Little Venice	<i>Malva neglecta</i> <i>Malva parviflora</i> <i>Hibiscus trionum</i>
Marsh Parsley	<i>Apium leptophyllum</i>
Mayweed*	<i>Anthemis cotula</i>
Morningglory Entireleaf Ivyleaf Red/Scarlet Smallflower Tall	<i>Ipomoea hederacea</i> var. <i>integriuscula</i> <i>Ipomoea hederacea</i> <i>Ipomoea coccinea</i> <i>Jacquemontia tamnifolia</i> <i>Ipomoea purpurea</i>
Moss	<i>Bryum</i> spp.
Mulberry Weed	<i>Fatua villosa</i>
Mustard Tumble Wild	<i>Sisymbrium altissimum</i> <i>Brassica kaber</i>
Nightshade Black Eastern Black Hairy	<i>Solanum nigrum</i> <i>Solanum ptycanthum</i> <i>Solanum sarrachoides</i>
Northern Willowherb	<i>Epilobium cillatum</i>
Panicum Fall* Texas*	<i>Panicum dichotomiflorum</i> <i>Panicum texanum</i>
Parsley-Peirt	<i>Alchemilla arvensis</i>
Pearlwort, Birdseye*	<i>Sagina procumbens</i>
Pennycress, Field	<i>Thlaspi arvense</i>
Phyllanthus, Longstalked	<i>Phyllanthus tenellus</i>
Pigweed Prostrate Redroot Smooth Tumble	<i>Amaranthus blitoides</i> <i>Amaranthus retroflexus</i> <i>Amaranthus hybridus</i> <i>Amaranthus albus</i>
Pineapple-weed*	<i>Matricaria matricarioides</i>
Plantain Broadleaf* Buckhorn*	<i>Plantago major</i> <i>Plantago lanceolata</i>
Poinsettia, Wild	<i>Euphorbia heterophylla</i>
Pondweed, Sago	<i>Potamogeton pectinatus</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, Common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed Common Giant	<i>Ambrosia artemisiifolia</i> <i>Ambrosia trifida</i>
Redmaids	<i>Calandrinia ciliate</i>

COMMON NAME	SCIENTIFIC NAME
Redweed	<i>Melochia corchorifolia</i>
Rocket, Yellow	<i>Barbarea vulgaris</i>
Senna, Coffee	<i>Cassia occidentalis</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Shepherd's-Purse	<i>Capsella bursa-pastoris</i>
Sida, Prickly (Teaweed)	<i>Sida spinosa</i>
Signalgrass*	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sowthistle, Annual	<i>Sonchus oleraceus</i>
Spiderwort, Tropical	<i>Commelina benghalensis</i>
Spurge Petty Prostrate Spotted	<i>Euphorbia peplus</i> <i>Euphorbia humistrata</i> Engelm <i>Euphorbia maculata</i>
Starbur, Bristly*	<i>Acanthospermum hispidum</i>
Tassle-flower	<i>Emilia</i> spp.
Thistle Canada* Russian	<i>Cirsium arvense</i> <i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp Common Tall	<i>Amaranthus rudis</i> <i>Amaranthus tuberculatus</i>
Woodsorrel, Yellow*	<i>Oxalis stricta</i>

*Preemergence control only

DIRECTIONS FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS

This product, when used as directed, can be used for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed-free. Apply this product only to:

- Bare ground under guard rails, above-ground pipelines, and railroad beds, railroad yards and surrounding areas
- Bare ground in parking and storage areas, plant sites, substations, pumping stations, and tank farms
- Bare ground areas of airports, brick yards, industrial plant sites, lumber yards, military installations, and storage areas
- Bare ground around farm buildings, and along ungrazed fence rows, wind breaks and shelter belts
- Road surfaces, improved roadside areas and gravel shoulders.

Follow all applicable directions as outlined above under Product Information. See table below for a list of broadleaf weeds and grasses controlled by this product.

This product offers residual and postemergence control of susceptible broadleaf and grass weeds as well as additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. The length of residual control is dependent on the rate applied as well as on the rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase.

PREEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of this product to a weed free soil surface. Preemergence applications of this product must be completed prior to weed emergence. Moisture

is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

POSTEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 quart per acre crop oil concentrate). The addition of an adjuvant enhances this product activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product. Emerged weeds are controlled postemergence with this product, however, translocation of this product within a weed is limited, and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with this product occurs when applied in combination with a surfactant to weeds less than 2 inches in height.

SOIL CHARACTERISTICS

Application of this product to soils with high organic matter and/or high clay content may require higher dosages than with soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

CARRIER VOLUME AND SPRAY PRESSURE

PREEMERGENCE APPLICATION

To ensure uniform coverage, use 10 to 30 gallons of spray solution per acre. Nozzle selection must meet manufacturer's gallonage and pressure directions for preemergence herbicide application.

POSTEMERGENCE APPLICATION

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre. Use 20 to 30 gallons per acre if dense vegetation or heavy residue is present on the soil surface. Nozzle selection must meet manufacturer's gallonage and pressure directions for postemergence herbicide application.

ADDITIVES

POSTEMERGENCE

When applying this product after weed emergence, mix with an agronomically approved adjuvant. Use a crop oil concentrate which contains at least 15% emulsifiers and 80% oil or a non-ionic surfactant containing at least 80% active ingredient when applying this product as part of a postemergence weed control program. Verify mixing compatibility by a jar test before using.

A spray grade nitrogen source (either ammonium sulfate at 2.0 to 2.5 pounds per acre or a 28 to 32% nitrogen solution at 1 to 2 quarts per acre) may be added to the spray mixture along with a crop oil concentrate or non-ionic surfactant to enhance weed control. The addition of a nitrogen source does not replace the need for crop oil concentrate or non-ionic surfactant.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND THIS PRODUCT

When using this product and an adjuvant, including in stale seed bed, layby, hooded/shielded or reduced tillage situations, perform a jar test before mixing commercial quantities of this product, when using for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pt of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 ml of this product to the quart jar for every 3 fl oz of this product per acre being applied (4 mls if 12 fl oz per acre is the desired rate of this product), gently mix until product goes into suspension.

3. Add 60 mls of crop oil to the quart jar or 1 ml of non-ionic surfactant if it is being used in place of oil, gently mix.
4. If nitrogen is being used, add 16 mls of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 g AMS to the quart jar in place of the 28 to 32% nitrogen.
5. Place a cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
6. An ideal tank mix combination will be uniform. If any of the following conditions are observed question the choice of adjuvant:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: Fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

APPLICATION EQUIPMENT

Application equipment must be clean and in good repair. Nozzles must be uniformly spaced on boom and frequently checked for accuracy.

BROADCAST APPLICATION

Apply this product, and this product's tank mixes, with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume.

BAND APPLICATION

When banding, use proportionately less water and this product per acre.

HANDGUN APPLICATION

Applications may also be made using a handgun sprayer. Use a spray volume of at least 40 gallons per acre to ensure uniform coverage.

AERIAL APPLICATION

To obtain satisfactory weed control with aerial applications of this product, uniform coverage must be obtained. **Do not** spray when drift is possible or when wind velocity is more than 10 mph. Avoid spraying this product within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory application and drift, the following directions must be observed:

Volume Pressure

Use this product in 5 to 10 gallons of water per acre with a maximum spray pressure of 40 PSI. Application at less than 5 gallons per acre will provide inadequate weed control. Higher gallonage applications provide more consistent weed control.

Nozzles and Nozzle Operation

Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles, including diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. **Do not** place nozzles on the outer 25% of the wings or rotors.

Adjuvants

Refer to the additive section or the tank mix partner's label for adjuvant specifications.

TANK MIX APPLICATION

In addition to weeds controlled by this product used alone, tank mixtures with other preemergence and postemergence herbicides registered for use in non-crop areas provide a broader spectrum of weed control. This product must be tank mixed with other non-crop herbicides including, but not limited to those products listed below.

TANK MIX COMBINATIONS FOR NON-SELECTIVE VEGETATION CONTROL

2,4-D	hexazinone	picloram
bromacil	imazapic	pramitol
chlorsulfuron	imazapyr	prodiamine
dicamba	metsulfuron-methyl	simazine
diuron	norflurazon	sulfometuron-methyl
clopyralid	oryzalin	tebuthiuron
glyphosate	pendimethalin	triclopyr

IMPORTANT: Completely read and follow the label of any potential tank mix partner. When using tank mixtures, use conditions must be in accordance with the most restrictive of the label limitations and precautions on either herbicide label.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.

DIRECTIONS FOR USE

†IN CONIFER RE-FORESTATION SITES FOLLOWING TIMBER HARVEST

This product is a preemergence and postemergence herbicide for control of selected grass and broadleaf weeds in conifer re-forestation sites following timber harvest operations. This product may be used as a site preparation treatment prior to transplanting of conifers or as a conifer release treatment after stand establishment.

†Not for use in California

Site Preparation-Application Before Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre. Transplant operations are permitted as soon as 4 weeks after application. To obtain optimal weed control, apply this product before weed emergence or after a burndown herbicide has controlled existing vegetation. If existing weed canopy is less than 40%, this product may be tank mixed with a burndown herbicide to provide preemergence weed control.

Apply this product in at least 10 gallons of water per acre to achieve uniform spray coverage using ground or aerial spray equipment.

Conifer Release Treatments – Applications only within 3 years after transplanting.

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre over the top of trees prior to budbreak in the spring or after dormancy in fall. **Do not** apply this product over the top of trees after budbreak or needle spotting and defoliation may occur. This product will not affect new growth of trees. See table below for a list of tolerant conifers for over the top treatments. Do not apply Flumi SC over the top unless trees are more than one year past seedling emergence.

TANK MIXING – Conifer Release Treatments

Certain liquid formulations of other pesticides may increase the postemergence activity of this product, but may also increase the potential for injury when applied over the top of various plants. Therefore, tank mixtures of these materials with this product may be more injurious than this product applied alone and need to be tested to determine if they can be used safely on a widespread basis.

ADJUVANTS – Conifer Release Treatments

When applying as a Conifer Release Treatment, **do not** mix this product with any adjuvant or fertilizer.

IMPORTANT: When applied as directed, the conifers listed in table below have shown tolerance to this product. However, this product is a very active herbicide. Exercise responsible judgment and caution until familiarity is gained with this product. If a desired conifer species is not listed in table below, evaluate the safety of this product on a small number of plants under commercial growing conditions, and monitor plant response for four to six weeks for phytotoxicity. Test this product on a small number of plants to determine if this product can be used safely on a widespread basis. **Do not** apply this product over the top of conifers until trees have been growing in the treated area for at least one year. The use of nylon mesh wraps, commonly used to deter animal browsing, may increase plant injury if placed on plants after over the top application of this product.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.

TOLERANT CONIFER TREE SPECIES

COMMON NAME	SCIENTIFIC NAME
Arborvitae American Oriental	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
Fir Concolor Cork Bark Douglas Fraser Grand Noble Turkish	<i>Abies concolor</i> <i>Abies lasiocarpa</i> <i>Pseudotsuga menziesii</i> <i>Abies fraseri</i> <i>Abies grandis</i> <i>Abies procera</i> <i>Abies bommuelleriana</i>
Hemlock Eastern Western	<i>Tsuga Canadensis</i> <i>Tsuga heterophylla</i>
Juniper Blue Star Creeping Japanese Garden Tamarix	<i>Juniperus scopularum</i> <i>Juniperus horizontalis</i> <i>Juniperus chinensis</i> <i>Juniperus Sabina</i>
Pine Austrian Eastern White Jack Japanese Black Loblolly Lodgepole Longleaf Mugo Ponderosa Sand Scotch Shortleaf	<i>Pinus nigra</i> <i>Pinus strobes</i> <i>Pinus banksiana</i> <i>Pinus thunbergiana</i> <i>Pinus taeda</i> <i>Pinus contorta</i> <i>Pinus palustris</i> <i>Pinus mugo</i> <i>Pinus ponderosa</i> <i>Pinus clausa</i> <i>Pinus sylvestris</i> <i>Pinus echinata</i>

COMMON NAME	SCIENTIFIC NAME
Slash Virginia	<i>Pinus elliotii</i> <i>Pinus virginiana</i>
Spruce Blue Dwarf Alberta Norway Sitka	<i>Picea pungens</i> <i>Picea glauca conica</i> <i>Picea abies</i> <i>Picea sitchensis</i>
Yew English Japanese	<i>Taxus baccata</i> <i>Taxus cuspidata</i>

DIRECTIONS FOR USE

†IN POPLAR PLANTATIONS AND TIMBER RE-FORESTATION SITES

This product is a preemergence and postemergence herbicide for control of selected grass and broadleaf weeds in poplar plantations and timber re-forestation sites following timber harvest operations. This product may be used as a site preparation treatment prior to transplanting of trees or as a release treatment after stand establishment.

†Not for use in California

Site Preparation – Application Before Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre. Transplant operations are permitted as soon as 4 weeks after application. To obtain optimal weed control, apply this product before weed emergence or after a burndown herbicide has controlled existing vegetation. If existing weed canopy is less than 40%, this product may be tank mixed with a burndown herbicide to provide preemergence weed control.

Apply this product in at least 10 gallons of water per acre to achieve uniform spray coverage using ground or aerial spray equipment.

Release Treatments – Applications Within 3 Years After Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre over the top of trees prior to budbreak in the spring or after dormancy in the fall. **Do not** apply this product over the top of trees after budbreak or leaf spotting and defoliation may occur. This product will not affect new growth of trees of tolerant poplars for over the top treatments. Do not apply Flumi SC over the top unless trees are more than one year past seedling emergence..

TANK MIXING – Poplar Release Treatments

Certain liquid formulations of other pesticides may increase the postemergence activity of this product, but may also increase the potential for injury when applied over the top of various plants. Therefore, tank mixtures of these materials with this product may be more injurious than this product applied alone and need to be tested to determine if they can be used safely on a widespread basis.

ADJUVANTS – Poplar Release Treatments

When applying Release Treatments, **do not** mix this product with any adjuvant or fertilizer.

IMPORTANT: When applied as directed, poplars (*Populus balsamifera*, *P. niger* and *P. tremuloides*), hybrid poplars (*P. sp. x sp.*), and cottonwoods (*P. deltoids* and *P. trichocarpa*) have shown tolerance to this product. However, this product is a very active herbicide. Exercise responsible judgment and caution until familiarity is gained with this product. Test this product on a small number of plants to determine if this product can be used safely on a widespread basis. **Do not** apply this product over the top unless trees are more than one year old.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Store in a cool, dry, secure place. Do not put formulation or dilute spray solution into food or drink containers. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. Not for use or storage in or around the home. For help with any spill, leak, fire or exposure involving this material, call day or night **CHEMTREC (800) 424-9300**.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Nonrefillable Containers 5 gallons or less:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Nonrefillable containers larger than 5 gallons:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Refillable containers larger than 5 gallons:] [Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

[PROMENADE® is a registered trademark of ADAMA group company.]

[FLUMIGARD® is a registered trademark of ADAMA group company.]

[EPA Approval Date]

REV020926

[SUBLABEL C]

[Note to reviewer: [Text] in brackets denotes optional text].

[Note to reviewer: {Text} in braces denotes where in the final label text will appear].

{[BOOKLET FRONT PANEL LANGUAGE]}

FLUMIOXAZIN	GROUP	14	HERBICIDE
-------------	-------	----	-----------

Alligare FLUMI SC

[Alternate Brand Names: Alligare FLUMIGARD® SC Herbicide; PROMENADE® SC Herbicide; PROMENADE® SC Specialty Herbicide]

[FOR THE MANAGEMENT OF UNDESIRABLE AQUATIC VEGETATION IN SLOW MOVING OR QUIESCENT WATERS]

[FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS, †CONIFER AND POPLAR RE-FORESTATION SITES]

[FOR USE IN CONTAINER AND FIELD GROWN CONIFERS (INCLUDING CHRISTMAS TREES) AND DECIDUOUS TREES, AROUND ESTABLISHED WOODY ORNAMENTALS IN LANDSCAPES AND MAINTAIN NON-CROP AREAS AND DORMANT BERMUDAGRASS][†Not for use in California]

ACTIVE INGREDIENT:

Flumioxazin* 42.0%

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

* (2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione)

Alligare FLUMI SC contains 4 pounds flumioxazin per gallon.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

[For ≤ 5 Gallon Containers:] [Shake Well Before Use]

[For ≥ 5 Gallon Containers:] [Shake Well, Agitate or Recirculate Before Use]

EPA Reg. No. 81927-78

EPA Est. No.

Manufactured for:

Alligare, LLC
PO Box 97
DeerPark, TX 77536

Net Contents:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes and socks

Mixers, loaders and applicators making aquatic subsurface applications using a mechanically-pressurized handgun must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves (made of any waterproof material)
- Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

If not used in accordance with directions on the label, this product is toxic to non-target plants and aquatic invertebrates. For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply directly to treated, finished drinking water reservoirs or drinking water receptacles when the water is intended for human consumption. Drift and runoff may be hazardous to non-target plants and aquatic organisms in water adjacent to treated areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants. Use strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Treatment of aquatic weeds can result in oxygen loss from decomposition of dead weeds. This loss can cause fish suffocation. Therefore, to minimize this hazard, treat 1/3 to 1/2 of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is needed.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with label precautionary statements and directions, and with applicable state and federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to users of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is:

- Coveralls
- Chemical-resistant gloves (made of waterproof material)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural crops on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

Do not enter or allow others to enter the treated area until sprays have dried.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product must be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These

risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Alligare. The Buyer must be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that the additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN DO NOT APPLY THIS PRODUCT. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND, TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Alligare shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

See also **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** sections of the label for additional information.

PRODUCT INFORMATION

This product is a fast-acting contact herbicide for use in the management of undesirable aquatic vegetation in slow moving or quiescent waters, to maintain non-crop areas, conifer and poplar re-forestation sites, container and field grown conifers (including Christmas trees) and deciduous trees, around established woody ornamentals in landscapes and dormant Bermudagrass.

This product is also effective as a preemergence and/or postemergence herbicide for control of selected grass and broadleaf weeds.

This product controls weeds by inhibiting protoporphyrinogen oxidase, an essential enzyme required by plants for chlorophyll biosynthesis. Seedling weeds are controlled preemergence when exposed to sunlight following contact with the soil applied herbicide.

This product may cause spotting or speckling on foliage if the spray solution directly contacts actively growing plant foliage or green bark. Leaves that receive indirect (drift) spray contact may be affected in a similar manner. Translocation of this product is limited, and under most conditions established and vigorously growing woody ornamentals will rapidly outgrow any injury symptoms. **However, direct application to actively growing foliage can cause severe injury or death with sensitive ornamental plant species, especially in herbaceous bedding plants and flowers.**

IMPORTANT: When applied as directed, plants listed on this label have shown tolerance to this product. However, this product is a very active herbicide. Exercise responsible judgment and caution until familiarity is gained with this product. Due to variability within species, crop growth stage, environmental conditions and application techniques, it is directed that users test this product under local growing conditions on a small number of plants and evaluate for 4 to 6 weeks for phytotoxicity. Testing this product on a small number of plants will determine if the herbicide can be used safely on a widespread application. Neither the seller nor the manufacturer of this product has investigated the safety to plants not listed on the label.

WEED RESISTANCE MANAGEMENT

This product is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Avoid the consecutive use of Alligare FLUMI SC or other target site of action Group 14 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Alligare, LLC retailer, representative or call 888-252-4427. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

PREEMERGENCE APPLICATION

Make the preemergence application of this product prior to weed emergence. Moisture is necessary to activate this product for residual weed control. Moisture is needed to move this product into the soil for preemergence weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

When adequate moisture is not received soon after this product is applied to soil, weed control may be improved by using shallow cultivation. If weeds begin to emerge, irrigate (1/2" of water) or cultivate uniformly with shallow tillage equipment that will not damage the crop. Deep cultivation reduces the effectiveness of this product.

POSTEMERGENCE APPLICATION

For best results, apply this product to actively growing weeds. The most effective postemergence weed control with this product occurs when applied in combination with a surfactant to weeds less than 2 inches in height. Applying this product under conditions that do not promote active weed growth will reduce herbicide effectiveness. **Do not** apply this product when the weeds are under stress due to drought, excessive water and extremes in temperatures or disease. This product is most effective when applied under sunny conditions at temperatures above 65° F. This product is rainfast one hour after application. **Do not** make applications if rain is expected within one hour of application or efficacy may be reduced.

SOIL CHARACTERISTICS

Application of this product to soils with high organic matter and/or high clay content may require higher dosages than with soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TANK MIX APPLICATION

In addition to weeds controlled by this product used alone, tank mixtures with other herbicides provides a broader spectrum of weed control. This product can be tank mixed with other herbicides including, but not limited to those products listed below.

TANK MIX COMBINATIONS FOR NON-SELECTIVE VEGETATION CONTROL

2,4-D	hexazinone	picloram
bromacil	imazapic	pramitol
chlorsulfuron	imazapyr	prodiamine
dicamba	metsulfuron-methyl	simazine
diuron	norflurazon	sulfometuron-methyl
clopyralid	oryzalin	tebuthiuron
glyphosate	pendimethalin	triclopyr

*Tank Mixing – Conifer and Poplar Release Treatments

Certain liquid formulations of other pesticides may increase the postemergence activity of this product, but may also increase the potential for injury when applied over the top of various plants. Therefore, tank mixtures of these materials with this product may be potential for injury when applied over the top of various plants. Therefore, tank mixtures of these materials with this product may be more injurious than this product applied alone and need to be tested to determine if they can be used safely on a widespread basis.

**Not for use in California*

Tank Mixing – Container and Field Grown Conifers

This product may be tank mixed with product containing the following active ingredients labeled for use in conifers:

clethodim
glyphosate*
oryzalin
prodiamine
simazine*

***Do not** apply glyphosate or simazine to containerized ornamentals.

Tank Mixing – Field and Container Grown Deciduous Trees

This product may be tank mixed with products containing the following active ingredient labeled for use in deciduous trees:

clethodim
glyphosate*
metolachlor
oryzalin
pendimethalin
prodiamine
simazine*

***Do not** apply glyphosate or simazine to containerized plants.

Tank Mixing – With Other Turfgrass Herbicides

This product may be tank mixed with appropriately labeled metsulfuron-methyl products.

SPRAY DRIFT MANAGEMENT

Aerial Applications

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the windspeed is 10 miles per hour or less, applicators must use ½ swath displacement upwind at the downwind edge of the application site. When the windspeed is between 11-15 miles per hour, applicators must use ¾ swath displacement upwind at the downwind edge of the application site.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- Controlling Droplet Size - Aircraft
 - Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
For ground equipment, the boom should remain level with the target area and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND
Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.
- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

APPLICATION AND SPRAYER INFORMATION

Apply this product with sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Apply by backpack or handgun sprayer, airboat, helicopter, airplane, or other application equipment that will ensure thorough coverage of plant foliage. **Important:** Thoroughly clean spray equipment, including all tanks, hoses, booms, screens and nozzles. **Do not use spray equipment used to apply this product to apply other materials or to any desirable plant foliage.** Equipment with this product's residue remaining in the system may result in crop injury to subsequently treated crops.

BROADCAST APPLICATION

Apply this product, and this product's tank mixes, with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume.

BAND APPLICATION

When banding, use proportionately less water and this product per acre.

HANDGUN APPLICATION

Applications may also be made using a handgun sprayer. Use a spray volume of at least 40 gallons per acre to insure uniform coverage.

BACKPACK APPLICATION

When applying this product with a backpack sprayer, follow all above restrictions. Calibrate backpack sprayers to deliver 1 gallon of spray solution per 500 to 1,000 square feet.

Mixing Rate for This Product in 1 Gallon of Spray Solution for Backpack Applications

Application Volume	Rate (fl oz/A)	Fl oz to Mix in 1 gal Water	Teaspoons to Mix in 1 gal Water	Mls to Mix in 1 gal Water
1 gal per 500 sq ft (87 GPA)	8	0.09	0.6	2.7
	10	0.11	0.7	3.4
	12	0.14	0.8	4.1
1 gal per 750 sq ft (65 GPA)	8	0.14	0.8	4.1
	10	0.17	1	5.1
	12	0.21	1.2	6.1
1 gal per 1,000 sq ft (44 GPA)	8	0.18	1.1	5.3
	10	0.23	1.4	6.8
	12	0.27	1.6	8.1

Example: Applicator wants to spray 1 gallon of this product solution per 1,000 square feet of ground bed at a rate of 12 fl oz/A. Mix 0.27 fl oz (1.6 teaspoons or 8.1 mls) of this product in 1 gallon of water.

AERIAL APPLICATION

To obtain satisfactory weed control, aerial application of this product must provide uniform coverage of surface weeds and sufficient contact time. When applied by air, this product may not provide adequate control of some submersed weeds. **Do not** apply by air when significant drift on to non-target plants may occur or when wind velocity is more than 10 mph. Avoid spraying this product within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory application and avoid drift, the following directions must be observed:

- Volume Pressure

Apply this product in a minimum of 5 gallons of water per acre, with a maximum spray pressure of 40 PSI. Application at less than 5 gallons per acre may not provide adequate weed control. Higher gallonage applications provide more consistent weed control.

- Nozzles and Nozzle Operation

Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles including diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. **Do not** place nozzles on the outer 25% of the wings or rotors.

SPRAYER PREPARATION

Before applying this product, start with clean, well maintained application equipment. Clean the spray tank, as well as all hoses and booms to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to the sulfonyleurea and phenoxy herbicides, are active at very small amounts and can cause crop injury when applied to susceptible crops. Clean the spray equipment according to the manufacturer's directions for the last product used before the equipment is used to apply this product. If two or more products were tank mixed prior to this product's application, follow the most restrictive cleanup procedure.

Mixing Instructions

- Mix with water having pH of 5 to 7. If pH is higher than 7, use an appropriate buffer to reduce pH to desirable range.
- Fill clean spray tank 1/2 full of desired level with water and add buffering agent if necessary.
- Add the required amount of this product to the spray tank while agitating.
- Fill spray tank to desired level with water. Ensure that this product is thoroughly mixed before making applications. Continue agitation until spray solution has been applied.
- If tank mixing this product with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
- Mix only the amount of spray solution that can be applied the day of mixing. Apply this product within 48 hours of mixing.

CARRIER VOLUME AND SPRAY PRESSURE

PREEMERGENCE APPLICATION

To ensure uniform coverage, use 10 to 40 gallons of spray solution per acre. When making backpack applications, apply 50 to 100 gallons of spray solution per acre. Nozzle must meet manufacturer's gallonage pressure directions for preemergence herbicide application.

POSTEMERGENCE APPLICATION

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre. Use 20 to 30 gallons per acre if dense vegetation or heavy residue is present on the soil surface. When applying with a backpack sprayer, apply 1 gallon of spray solution per 500 to 1,000 square feet. Nozzle selection must meet manufacturer's gallonage and pressure directions for postemergence herbicide application.

ADDITIVES

When applying this product to the foliage of floating or emerged aquatic weeds, mix with an adjuvant approved for use in aquatic sites. Follow adjuvant manufacturer's label rates. Verify mixing compatibility by a jar test before using.

When applying this product after weed emergence in terrestrial settings, mix with an agronomically approved adjuvant. A non-ionic surfactant containing at least 80% active ingredient must be used when applying this product as part of a postemergence weed control program. Verify mixing compatibility by a jar test before using.

ADJUVANTS

Refer to the additive section or the tank mix partners label for adjuvant specifications.

When applying Release Treatments, **do not** mix this product with any adjuvant or fertilizer.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND THIS PRODUCT

Perform a jar test before mixing commercial quantities of this product, when using this product for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pint of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 ml of this product to the quart jar for every 3 fl oz of this product per acre being applied (4 ml if 12 fl oz per acre is the desired rate of this product), gently mix until product goes into suspension.
3. Add 1 ml of non-ionic surfactant or 60 ml of crop oil concentrate, gently mix.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An ideal tank mix combination will be uniform. If any of the following conditions are observed question the choice of adjuvant:
 - a) Layer of oil or globules on the mixture's surface,
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar,
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

Sprayer Cleanup

If spray equipment is dedicated to application of aquatic herbicides, use the following steps to clean the spray equipment:

- Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.

If spray equipment will be used for purposes other than applying aquatic herbicides, it must be thoroughly cleaned following application of this product. Use the following steps to clean the spray equipment:

1. Completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank with clean water and household ammonia. Use 1 gallon of 3% household ammonia for every 100 gallons of water.
4. Circulate through sprayer for 5 minutes.
5. Then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
6. Loosen any diaphragms before flushing the spray system, allowing cleaning solution to spray through the open diaphragm.
7. Drain tank completely.
8. Add enough clean water to the spray tank to flush hoses, booms, screens and nozzles for 2 minutes.
9. Remove all nozzles and screens and rinse them with clean water.

WEEDS CONTROLLED

When this product is applied preemergence or postemergence at directed rates and weed stages, the following grasses, broadleaf weeds are controlled.

WEEDS CONTROLLED

Common Name	Scientific Name
Alyssum, Hoary	<i>Berteroa incana</i>
Amaranth	
Palmer	<i>Amaranthus palmeri</i>
Spiny	<i>Amaranthus spinosus</i>
American Burnweed	<i>Erechetities hieracifolia</i>
Barnyardgrass*	<i>Echinochloa crus-galli</i>
Beggarweed, Florida	<i>Desmodium Tortuosum</i>
Bittercress, Hairy	<i>Cardamine hirsute</i>
Bluegrass, Annual	<i>Poa annua</i>
Burclover, California	<i>Medicago polymorpha</i>
Carpetweed	<i>Mollugo verticillata</i>
Chamberbitter	<i>Phyllanthus urinaria</i>
Chickweed	
Common	<i>Stellaria media</i>
Mouseear	<i>Cerastium vulgatum</i>
Crabgrass	
Large*	<i>Digitaria sanguinalis</i>
Smooth*	<i>Digitaria ischaemum</i>
Southern*	<i>Digitaria ciliaris</i>
Croton, Tropic	<i>Croton glandulosus var. septentrionalis</i>
Dandelion*	<i>Taraxacum officinale</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Doveweed	<i>Murdannia nudiflora</i>
Eclipta	<i>Eclipta prostrata</i>
Filaree, Redstem*	<i>Erodium cicutarium</i>
Foxtail	
Bristly*	<i>Setaria verticillata</i>
Giant*	<i>Setaria faberi</i>
Green*	<i>Setaria viridis</i>
Yellow*	<i>Setaria glauca</i>
Galinsoga, Hairy	<i>Galinsoga ciliata</i>
Geranium, Carolina	<i>Geranium carolinianum</i>
Goosegrass*	<i>Eleusine indica</i>
Groundsel, Common	<i>Senecio vulgaris</i>
Groundsel Tree	<i>Baccharis halimifolia</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed*	<i>Conyza Canadensis</i>
Indigo, Hairy	<i>Indigofera hirsuta</i>
Ivy, Ground*	<i>Glechoma hederacea</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Kyllinga, Green*	<i>Kyllinga brevifolia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Liverwort	<i>Marchantia polymorpha</i>
Lovegrass, California*	<i>Eragrostis diffusa</i>
Mallow	
Common	<i>Malva neglecta</i>
Little	<i>Malva parviflora</i>

Common Name	Scientific Name
Venice	<i>Hibiscus trionum</i>
Marsh Parsley	<i>Apium leptophyllum</i>
Mayweed*	<i>Anthemis cotula</i>
Morningglory	
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integruscula</i>
Ivyleaf	<i>Ipomoea hederacea</i>
Red/Scarlet	<i>Ipomoea coccinea</i>
Smallflower	<i>Jacquemontia tamnifolia</i>
Tall	<i>Ipomoea purpurea</i>
Moss	<i>Bryum</i> spp.
Mulberry Weed	<i>Fatoua villosa</i>
Mustard	
Tumble	<i>Sisymbrium altissimum</i>
Wild	<i>Brassica kaber</i>
Nightshade	
Black	<i>Solanum nigrum</i>
Eastern Black	<i>Solanum ptycanthum</i>
Hairy	<i>Solanum sarrachoides</i>
Northern Willowherb	<i>Epilobium cillatum</i>
Panicum	
Fall*	<i>Panicum dichotomiflorum</i>
Texas*	<i>Panicum texanum</i>
Parsley-Piert	<i>Alchemilla arvensis</i>
Pearlwort, Birdseye*	<i>Sagina procumbens</i>
Pennycress, Field	<i>Thlaspi arvense</i>
Phyllanthus, Longstalked	<i>Phyllanthus tenellus</i>
Pigweed	
Prostrate	<i>Amaranthus blitoides</i>
Redroot	<i>Amaranthus retroflexus</i>
Smooth	<i>Amaranthus hybridus</i>
Tumble	<i>Amaranthus albus</i>
Pineappleweed*	<i>Matricaria matricarioides</i>
Plantain	
Broadleaf*	<i>Plantago major</i>
Buckhorn*	<i>Plantago lanceolata</i>
Poinsettia, Wild	<i>Euphorbia heterophylla</i>
Pondweed, Sago	<i>Potamogeton pectinatus</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, Common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed	
Common	<i>Ambrosia artemisiifolia</i>
Giant	<i>Ambrosia trifida</i>
Redmaids	<i>Calandrinia ciliata</i>
Redweed	<i>Melochia corchorifolia</i>
Rocket, Yellow	<i>Barbarea vulgaris</i>
Senna, Coffee	<i>Cassia occidentalis</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Shepherd's-Purse	<i>Capsella bursa-pastoris</i>
Sida, Prickly (Teaweed)	<i>Sida spinosa</i>
Signalgrass*	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Sowthistle, Annual	<i>Sonchus oleraceus</i>
Spiderwort, Tropical	<i>Commelina benghalensis</i>
Spurge	

Common Name	Scientific Name
Petty	<i>Euphorbia peplus</i>
Prostrate	<i>Euphorbia humistrata</i> Engelm
Spotted	<i>Euphorbia maculata</i>
Starbur, Bristly*	<i>Acanthospermum hispidum</i>
Tassel-flower	<i>Emilia</i> spp.
Thistle	
Canada*	<i>Cirsium arvense</i>
Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp	
Common	<i>Amaranthus rudis</i>
Tall	<i>Amaranthus tuberculatus</i>
Woodsorrel, Yellow*	<i>Oxalis stricta</i>

*Preemergence control only.

AQUATIC WEED CONTROL

This product may be applied to the following quiescent or slow-moving bodies of water:

- Bayous
- Canals
- Drainage ditches
- Lakes
- Marshes
- Ponds (including golf course ponds)
- Reservoirs

This product is most effective when applied to young, actively growing weeds in water with a pH of less than 8.5. Application of this product to public aquatic areas may require special approval and/or permits. Consult with local state agencies, if required.

USE RESTRICTIONS

- Do not apply to intertidal or estuarine areas.
- Do not exceed 400 ppb of this product during any one application.
- Do not re-treat the same section of water with this product more than 6 times per year.
- Do not retreat the same section of water within 28 days of application, except in areas with dense weed vegetation. In these areas, treat the remaining weeds within 10 to 14 days.
- In high density weed populations only treat 1/2 the water body at one time.
- Treated water may not be used for irrigation purposes on food crops until at least five (5) days after application.
- Do not use in water utilized for crawfish farming.

USE PRECAUTIONS

- There is no post-application holding restriction against use of treated water for drinking or recreational purposes (e.g., swimming, fishing).
- Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the *Irrigation Restrictions Following Application* table.

IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

Application Method	Application Rate	Average Water Depth	Turf and Landscape Ornamentals	Ornamentals grown for production in Greenhouse and Nursery
Surface Spray	6 to 12 oz per surface acre	Greater than 3 feet	None	5 days
		Less than 3 feet	12 hours	5 days
Subsurface	Less than 200 ppb	N/A	1 day	5 days
	200 to 300 ppb	N/A	2 days	5 days
	300 to 400 ppb	N/A	3 days	5 days

DIRECTIONS FOR USE

TO CONTROL FLOATING AND EMERGED WEEDS USING SURFACE APPLICATION

This product will control weeds and algae listed in table below when applied as a broadcast spray with appropriate equipment. For best results, apply this product to the foliage of actively growing weeds.

Floating and Emerged Weeds

Common Name	Scientific Name
Alligator Weed	<i>Alternanthera philoxeroides</i>
Duckweed*	<i>Lemna</i> spp.
Frog's-bit	<i>Limnobium spongia</i>
Mosquito Fern	<i>Azolla</i> spp.
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal*	<i>Wolffia</i> spp.
Water Pennywort	<i>Hydrocotyle</i> spp.
Filamentous algae	<i>Pithophara</i>
Filamentous algae	<i>Cladophora</i>

*Coverage is essential for effective duckweed and watermeal control. Any duckweed and/or watermeal escapes left in the water column will quickly re-infest the water body. Apply 200 ppb concentration throughout the water body to control duckweed and watermeal. – see **DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATIONS** section for additional application information.

Surface Application

Apply this product as a broadcast spray at 6 to 12 fl oz of formulated product (0.19 to 0.38 lb ai) per acre plus an adjuvant approved for use in aquatics.

This product is a contact herbicide that quickly degrades in the water column so plants that do not initially come in contact with the herbicide will not be controlled. Apply this product in a minimum of 30 gals of water per acre to all areas of the water body where weeds exist. Coverage is essential for effective control as all floating weeds need to be exposed to lethal concentrations in all parts of the water body. Any untreated escapes or re-introductions of plants that were not treated will reestablish in areas where surface weeds had previously been controlled. If a second application is required to provide control, make a treatment once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Application of this product during early morning hours may enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage. Rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in

dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat floating surface weeds in section to avoid a rapid decrease in dissolved oxygen.

This product may be tank mixed with 2,4-D, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds.

Consult a manufacturer's label for specific rate restrictions and weeds controlled. Always follow the most restrictive label restrictions and precautions for all products used when making an application involving tank mixes.

DIRECTIONS FOR USE

TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATION

This product will control submersed and floating weeds listed in table below, *Submersed and Floating Weeds Controlled by Subsurface Application*, when applied subsurface with appropriate equipment.

Submersed and Floating Weeds Controlled by Subsurface Application

Coontail	<i>Ceratophyllum demersum</i>
Duckweed	<i>Lemna</i> spp.
Fanwort	<i>Cabomba caroliniana</i>
Hydrilla	<i>Hydrilla verticillata</i>
Hygrophila	<i>Hygrophila polysperma</i>
Naiad, Southern	<i>Najas guadalupensis</i>
Pondweed, Curlyleaf	<i>Potamogeton crispus</i>
Pondweed, Sago	<i>Potamogeton pectinatus</i>
Pondweed, Variable-Leaf	<i>Potamogeton diversifolius</i>
Water Fern	<i>Salvinia</i> spp.
Water Lettuce	<i>Pistia stratiotes</i>
Watermeal	<i>Wolffia</i> spp.
Watermilfoil, Eurasian	<i>Myriophyllum spicatum</i>
Watermilfoil, Variable-Leaf	<i>Myriophyllum heterophyllum</i>

Subsurface Treatment

Apply this product at a rate that will produce an initial concentration of 200 to 400 ppb (of active ingredient flumioxazin) in the water column.

This product is rapidly absorbed by target plants, but also breaks down quickly in water with a pH greater than 8.5. The pH of water surrounding mats of submersed vegetation can exceed 8.5 by early to mid-day, due to photosynthetic processes. Application of this product under these conditions may provide only partial weed control, and regrowth is likely. For best control, apply this product in a minimum of 30 gals of water per acre in the early morning to actively growing weeds and early in the season before surface matting occurs. Complete coverage and sufficient contact time of submersed weeds with this product is required for optimal performance. Application of this product with subsurface trailing hoses designed to distribute the herbicide within the plant stand will provide more effective and longer-term control of submersed weeds. Use, *Subsurface Application Rates* to determine the amount of this product needed to achieve desired concentration at different water depths. Use higher concentrations when weed biomass is heavy and/or weeds are more mature and topped out. Any untreated plants that are left in the water column can re-infest treated areas that had previously been controlled. If a second application is required to provide control, it is advised that a treatment be made once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

When applying this product to densely packed actively growing submersed weeds, a rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease

in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat submersed weeds in sections to avoid a rapid decrease in dissolved oxygen.

This product may be tank mixed with other registered submersed applied herbicides for enhanced control of submersed and floating weeds.

Application Equipment for Water Column Treatment

To improve distribution in the water column and ensure adequate coverage, when possible apply this product with subsurface trailing hoses in order to place the herbicide under the surface and throughout the biomass of aquatic vegetation. Keep swath width to a minimum in order to maximize contact with submersed aquatic vegetation. In small shallow water bodies, surface sprays may be required to apply this product. Apply by backpack or handgun sprayer or other application equipment that will ensure adequate coverage of target plant.

Information on Hydrilla Control in Florida

Apply this product as a subsurface treatment for hydrilla control. For best control of hydrilla apply during the late Winter/early Spring and/or early to late Fall. Efficacy of this product will be enhanced at these timings due to lower potential biomass present and lower pH of the water. If applied to mature topped out hydrilla, this product will cause some discoloration and loss of growing tips, but regrowth will be rapid.

Tank mixing this product with other registered herbicides is advised, especially if hydrilla is approaching maturity or biomass is heavy.

Subsurface Application Rates

Water Depth (feet)	Pints of This Product Required Per Surface Acre to Achieve Desired Water Concentration		
	200 ppb	300 ppb	400 ppb
1	1.1	1.6	2.1
2	2.1	3.2	4.2
3	3.2	4.8	6.4
4	4.2	6.4	8.5
5	5.3	8.0	10.6

Example: To achieve an initial concentration of 200 ppb of flumioxazin in a 4-foot deep water column, apply 4.2 pints of this product per surface acre.

BARE GROUND NON-CROP AREAS, CONIFER AND POPLAR RE-FORESTATION SITES*

**Alligare FLUMI SC is not approved for use on Conifer and Poplar Re-Forestation Sites in California*

DIRECTIONS FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS

This product, when used as directed, can be used for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed-free. Apply this product only to:

- Bare ground under guard rails, above-ground pipelines, and railroad beds, railroad yards and surrounding areas
- Bare ground in parking and storage areas, plant sites, substations, pumping stations, and tank farms
- Bare ground areas of airports, brick yards, industrial plant sites, lumber yards, military installations, and storage areas
- Bare ground around farm buildings, and along ungrazed fence rows, wind breaks and shelter belts
- Road surfaces, improved roadside areas and gravel shoulders

This product offers residual and postemergence control of susceptible broadleaf and grass weeds as well as additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. The length of residual control is dependent on the rate applied as well as on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply to moist or wet desirable plant foliage.
- Do not apply within 300 feet of non-dormant pome or stone fruit crops.
- Do not re-apply this product within 30 days.

USE PRECAUTIONS

- Treatment of powdery, dry soil or light sandy soil, or light sandy soil when there is little to no likelihood of rainfall soon after may result in off target movement and possible damage to actively growing susceptible crops when soil particles are moved by wind or water. **Do not** apply when these soil and environmental conditions are present.

PREEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of this product to a weed free soil surface. Preemergence applications of this product must be completed prior to weed emergence.

POSTEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 quart per acre crop oil concentrate). The addition of an adjuvant enhances this product's activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product. Emerged weeds are controlled postemergence with this product, however, translocation of this product within a weed is limited, and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with this product occurs when applied in combination with a surfactant to weeds less than 2 inches in height.

†IN CONIFER RE-FORESTATION SITES FOLLOWING TIMBER HARVEST DIRECTIONS FOR USE

This product is a preemergence and postemergence herbicide for control of selected grass and broadleaf weeds in conifer re-forestation sites following timber harvest operations. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. This product may be used as a site preparation treatment prior to transplanting of conifers or as a conifer release treatment after stand establishment.

†Not for use in California

Site Preparation – Application Before Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre. Transplant operations are permitted as soon as 4 weeks after application. To obtain optimal weed control, apply this product before weed emergence or after a burndown herbicide has controlled existing vegetation. If existing weed canopy is less than 40%, this product may be tank mixed with a burndown herbicide to provide preemergence weed control.

Apply this product in at least 10 gallons of water per acre to achieve uniform spray coverage using ground or aerial spray equipment.

Conifer Release Treatments – Application only within 3 years after transplanting.

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre over the top of trees prior to budbreak in the spring or after dormancy in fall. **Do not** apply this product over the top of trees after budbreak or needle spotting and defoliation may occur. This product will not affect new growth of trees. See table below for a list of tolerant conifers for over the top treatments. Do not apply Flumi SC over the top unless trees are more than one year past seedling emergence.

IMPORTANT: When applied as directed, the conifers listed below have shown tolerance to this product. However, this product is a very active herbicide. Exercise responsible judgment until familiarity is gained with this product. If a desired conifer species is not listed below, evaluate the safety of this product on a small number of plants under commercial growing conditions, and monitor plant response for four to six weeks for phytotoxicity. Test this product on a small number of plants to determine if this product can be used safely on a widespread basis. **Do not** apply this product over the top of conifers until trees have been growing in the treated area for at least one year. The use of nylon mesh wraps, commonly used to deter animal browsing, may increase plant injury if placed on plants after over the top application of this product.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply to moist or wet desirable plant foliage.
- Do not apply within 300 feet of non-dormant pome or stone fruit crops.
- Do not re-apply this product within 30 days.

USE PRECAUTIONS

- Treatment of powdery, dry soil or light sandy soil, or light sandy soil when there is little to no likelihood of rainfall soon after may result in off target movement and possible damage to actively growing susceptible crops when soil particles are moved by wind or water. **Do not** apply when these soil and environmental conditions are present.

TOLERANT CONIFER TREE SPECIES

COMMON NAME	SCIENTIFIC NAME
Arborvitae American Oriental	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
Fir Concolor Cork Bark Douglas Fraser Grand Noble Turkish	<i>Abies concolor</i> <i>Abies lasiocarpa</i> <i>Pseudotsuga menziesii</i> <i>Abies fraseri</i> <i>Abies grandis</i> <i>Abies procera</i> <i>Abies bommuelleriana</i>
Hemlock Eastern Western	<i>Tsuga Canadensis</i> <i>Tsuga heterophylla</i>
Juniper Blue Star Creeping Japanese Garden Tamarix	<i>Juniperus scopularum</i> <i>Juniperus horizontalis</i> <i>Juniperus chinensis</i> <i>Juniperus Sabina</i>
Pine Austrian Eastern White Jack Japanese Black Loblolly Lodgepole Longleaf Mugo Ponderosa Sand Scotch Shortleaf Slash Virginia	<i>Pinus nigra</i> <i>Pinus strobes</i> <i>Pinus banksiana</i> <i>Pinus thunbergiana</i> <i>Pinus taeda</i> <i>Pinus contorta</i> <i>Pinus palustris</i> <i>Pinus mugo</i> <i>Pinus ponderosa</i> <i>Pinus clausa</i> <i>Pinus sylvestris</i> <i>Pinus echinata</i> <i>Pinus elliotii</i> <i>Pinus virginiana</i>
Spruce Blue Dwarf Alberta Norway Sitka	<i>Picea pungens</i> <i>Picea glauca conica</i> <i>Picea abies</i> <i>Picea sitchensis</i>
Yew English Japanese	<i>Taxus baccata</i> <i>Taxus cuspidata</i>

†IN POPLAR PLANTATIONS AND TIMBER RE-FORESTATION SITES

DIRECTIONS FOR USE

This product is a preemergence and postemergence herbicide for control of selected grass and broadleaf weeds in poplar plantations and timber re-forestation sites following timber harvest operations. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. This product may be used as a site preparation treatment prior to transplanting of trees or as a release treatment after stand establishment.

†*Not for use in California*

Site Preparation – Application Before Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre. Transplant operations are permitted as soon as 4 weeks after application. To obtain optimal weed control, apply this product before weed emergence or after a burndown herbicide has controlled existing vegetation. If existing weed canopy is less than 40%, this product may be tank mixed with a burndown herbicide to provide preemergence weed control.

Apply this product in at least 10 gallons of water per acre to achieve uniform spray coverage using ground or aerial spray equipment.

Release Treatments – Applications within 3 Years After Transplanting

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per acre over the top of trees prior to budbreak in the spring or after dormancy in fall. **Do not** apply this product over the top of trees after budbreak or leaf spotting and defoliation may occur. This product will not affect new growth of trees of tolerant poplars for over the top treatments. Do not apply Flumi SC over the top unless trees are more than one year past seedling emergence.

TANK MIXING – Poplar Release Treatments

Certain liquid formulations of other pesticides may increase the postemergence activity of this product but may also increase the potential for injury when applied over the top of various plants. Therefore, tank mixtures of these materials with this product may be more injurious than this product applied alone and need to be tested to determine if they can be used safely on a widespread basis.

ADJUVANTS – Poplar Release Treatments

When applying Release Treatments, **do not** mix this product with any adjuvant or fertilizer.

IMPORTANT: When applied as directed, poplars (*Populus balsamifera*, *P. niger* and *P. tremuloides*), hybrid poplars (*P. sp. x sp.*), and cottonwoods (*P. deltoids* and *P. trichocarpa*) have shown tolerance to this product. However, this product is a very active herbicide. Exercise responsible judgment until familiarity is gained with this product. Test this product on a small number of plants to determine if this product can be used safely on a widespread basis. **Do not** apply this product over the top unless trees are more than one year old.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply to moist or wet desirable plant foliage.
- Do not apply within 300 feet of non-dormant pome or stone fruit crops.
- Do not re-apply this product within 30 days.

USE PRECAUTIONS

- Treatment of powdery, dry soil or light sandy soil, or light sandy soil when there is little to no likelihood of rainfall soon after may result in off target movement and possible damage to actively growing susceptible crops when soil particles are moved by wind or water. **Do not** apply when these soil and environmental conditions are present.

TURF AND ORNAMENTAL SITES

DIRECTIONS FOR USE

This product is a preemergence and early postemergence herbicide for control of selected grass and broadleaf weeds in and around ornamental woody shrubs, deciduous trees, and conifers (including Christmas trees) grown outdoors in containers or in the field (in ground), to maintain non-crop areas and dormant Bermudagrass. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses.

This product controls weeds by inhibiting protoporphyrinogen oxidase, an essential enzyme required by plants for chlorophyll biosynthesis. Seedling weeds are controlled preemergence when exposed to sunlight following contact with the soil applied herbicide.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not apply in enclosed greenhouse structures in plants are present.
- Do not move plants for 24 hours into enclosed greenhouses until the area treated with this product has been watered.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not graze treated fields or hay to livestock.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply when plants are under stress from insects, diseases, animals or winter injury, planting shock or any other stresses.
- Only apply to healthy established trees and ornamentals.

IN ESTABLISHED CONTAINER AND FIELD GROWN CONIFERS

(INCLUDING CHRISTMAS TREES)

DIRECTIONS FOR USE

Apply this product as a single or split application to established container and field grown conifers, which includes applications to Christmas tree plantations. The conifers listed below have exhibited tolerance to this product only when the product is applied to dormant or hardened off plant material. If applied over the top of plant foliage, apply this product before spring bud break or after conifers have sufficiently hardened off. During periods of cool, cloudy weather, use caution to ensure conifers have hardened off prior to herbicide application. **Do not** apply to conifers within 1 year of seedling emergence.

PREEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre before weeds emerge. Apply to weed free, established conifers grown in containers or in the field (in ground). If possible, irrigate treated area with 0.5 to 0.75 inch of water immediately following application. This product may be sprayed directly over conifers listed below, provided bud break has not occurred or plants are hardened off. Needle burn may be observed on new flush if plants are actively growing at time of application. However, this product will typically not affect subsequent growth. If conifers are not dormant or hardened off at time of application, and foliar injury cannot be tolerated, apply this product as a directed spray, taking care to minimize direct contact or drift of sprays onto foliage. Mechanically incorporating this product after application will disturb soil surface, which may reduce herbicidal efficacy. When applied before weed germination, this product will control broadleaf and grassy weeds listed in WEEDS CONTROLLED.

POSTEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre after weeds have emerged. This product may be sprayed directly over conifers listed below, provided bud break has not occurred or plants are hardened off. Needle burn may be observed on new flush if plants are actively growing at time of application. However, this product will typically not affect subsequent growth. If conifers are not dormant or hardened off at the time of application, and foliar injury cannot be tolerated, apply this product as a directed spray, taking care to minimize direct contact or drift of sprays onto foliage.

If applied when weeds are actively growing and no larger than 2 inches in height, this product will provide postemergence control of broadleaf weeds and grasses listed in WEEDS CONTROLLED. Postemergence control of this product may be more effective with certain weed species, and may not control mature, stressed or hardened off weeds that are not actively growing at the time of application.

TOLERANT CONIFERS

This product may be applied to the conifer species listed below. If a desired conifer species is not listed below, evaluate the safety of this product on a small number of plants under commercial growing conditions, and monitor plant response for four to six weeks for phytotoxicity. Testing this product on a small number of plants will determine if this product can be used safely on a widespread basis.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.

TOLERANT CONIFER TREE SPECIES

COMMON NAME	SCIENTIFIC NAME
Arborvitae American Oriental	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
Fir Concolor Cork Bark Douglas Fraser Grand Noble Turkish	<i>Abies concolor</i> <i>Abies lasiocarpa</i> <i>Pseudotsuga menziesii</i> <i>Abies fraseri</i> <i>Abies grandis</i> <i>Abies procera</i> <i>Abies bommuelleriana</i>
Hemlock Eastern Western	<i>Tsuga Canadensis</i> <i>Tsuga heterophylla</i>
Juniper Blue Star Creeping Japanese Garden Tamarix	<i>Juniperus scopularum</i> <i>Juniperus horizontalis</i> <i>Juniperus chinensis</i> <i>Juniperus Sabina</i>
Pine Austrian Eastern White Jack Japanese Black Loblolly Lodgepole	<i>Pinus nigra</i> <i>Pinus strobes</i> <i>Pinus banksiana</i> <i>Pinus thunbergiana</i> <i>Pinus taeda</i> <i>Pinus contorta</i>

COMMON NAME	SCIENTIFIC NAME
Longleaf	<i>Pinus palustris</i>
Mugo	<i>Pinus mugo</i>
Ponderosa	<i>Pinus ponderosa</i>
Sand	<i>Pinus clausa</i>
Scotch	<i>Pinus sylvestris</i>
Shortleaf	<i>Pinus echinata</i>
Slash	<i>Pinus elliotii</i>
Virginia	<i>Pinus virginiana</i>
Spruce	
Blue	<i>Picea pungens</i>
Dwarf Alberta	<i>Picea glauca conica</i>
Norway	<i>Picea abies</i>
Sitka	<i>Picea sitchensis</i>
Yew	
English	<i>Taxus baccata</i>
Japanese	<i>Taxus cuspidata</i>

IN CONTAINER AND FIELD DECIDUOUS TREES AND NON-BEARING FRUIT AND NON-BEARING NUT TREES DIRECTIONS FOR USE

This product may be applied as single or split applications to container and field grown deciduous trees with an established root system. The deciduous trees listed have exhibited tolerance to this product only when applied to the soil and base of plants. Application of this product to deciduous foliage or green bark may result in unacceptable injury.

This product may be applied to established (or transplanted) container and field grown deciduous trees. **Do not** apply to trees that are less than one year old or have been transplanted less than one year, unless completely protected by non-porous wraps, grow tubes, waxed protectors, or other forms of protection to young foliage and/or bark. **Do not** harvest fruit or nuts from treated trees within one year of application.

IMPORTANT: Direct of application of this product to the soil surface and away from plant foliage and bark. Avoid direct spray contact on plant surfaces, foliage and green bark or injury may result. Application of this product after bud swell may cause injury if herbicide contact foliage. Avoid application under environmental conditions that favor drift to non-targeted areas.

PREEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre as a preemergence (to weed emergence) application. Apply this product to weed free deciduous trees grown in containers or in the field (in-ground). If possible, irrigate treated area with 0.5 to 0.75 inch of water immediately following application. This product may be applied to the soil surface and base of deciduous trees, provided that direct and indirect (drift) applications to plant foliage, flowers and green bark does not occur. Mechanically incorporating this product will disturb soil surfaces, which may reduce herbicidal efficacy. The use of spray shields that limit exposure of foliage and bark to this product is suggested. When applied before weed germination, this product will control broadleaf and grassy weeds. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses.

POSTEMERGENCE APPLICATION

Apply 8 to 12 fl oz (0.25 to 0.38 lb ai) per acre of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant). Make postemergence (to weed emergence) applications of this product when weeds are actively growing and are no larger than 2 inches in height. The addition of a surfactant enhances this product activity on emerged weeds. Thorough spray coverage is necessary to maximize the

postemergence activity of this product. When applied after weed germination, this product will provide preemergence and postemergence control of broadleaf weeds and grasses. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses.

Postemergence control of this product may be more effective with certain weed species, and may not control mature, stressed or hardened off weeds that are not actively growing at the time of application.

TOLERANT DECIDUOUS TREES, NON-BEARING FRUIT AND NON-BEARING NUT TREES

This product may be applied as a directed spray to the deciduous, non-bearing fruit and non-bearing nut trees species listed in **TOLERANT DECIDUOUS TREE SPECIES**. If a desired tree species is not listed in Table 6, evaluate the safety of this product on a small number of plants under commercial growing conditions and monitor plant response for four to six weeks for phytotoxicity. Testing this product on small number of plants will determine if this product can be used safely on a widespread basis.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.

TOLERANT DECIDUOUS TREE SPECIES

COMMON NAME	SCIENTIFIC NAME
Apricot*	<i>Prunus</i> spp.
Ash	<i>Fraxinus</i> spp.
Birch	<i>Betula</i> spp.
Buckeye	<i>Aesculus</i> spp.
Cherry*	<i>Prunus</i> spp.
Chestnut	<i>Castanea</i> spp.
Citrus*	<i>Citrus</i> spp.
Dogwood	<i>Cornus</i> spp.
Eucalyptus	<i>Eucalyptus</i> spp.
Ginkgo	<i>Ginkgo</i> spp.
Hawthorn	<i>Crataegus</i> spp.
Honeylocust	<i>Gleditsia</i> spp.
Larch	<i>Larix</i> spp.
Lilac	<i>Syringa</i> spp.
Maple**	<i>Acer</i> spp.
Myrtle, Crepe	<i>Lagerstroemia indica</i>
Oak	<i>Quercus</i> spp.
Poplar	<i>Populus</i> spp.
Peach*	<i>Prunus</i> spp.
Plum*	<i>Prunus</i> spp.
Pecan*	<i>Carya</i> spp.
Redbud	<i>Cercis Canadensis</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Plantanus</i> spp.
Walnut, Black	<i>Juglens nigra</i>
Willow	<i>Salix</i> spp.

*Non-bearing trees only.

**Not for use on maple trees used for production of maple sap or syrup.

AROUND ESTABLISHED WOODY LANDSCAPE ORNAMENTALS AND TO MAINTAIN NON-CROP AREAS DIRECTIONS FOR USE

Application of this product in the vicinity of ornamental plants is limited to directed sprays around well-established woody shrubs and trees including azalea, euonymus, holly, and the conifers and deciduous trees listed. This product may also be applied to maintain weed control in non-crop areas in apartment complexes, fence rows, gravel surfaces and driveways, ground mats and pads prior to the addition of containerized plants, golf courses, lumberyards, office complexes, parks, parking areas, recreational sites, schools, sidewalks, storage areas, grass waterways, rain gardens, and other similar industrial sites. **Do not** apply this product within any enclosed structure in residential or commercial landscapes.

This product offers postemergence and residual control of susceptible grasses and broadleaf weeds, as well as additional mode of action to assist in the control of resistant weeds. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. The length of residual control is dependent on the rate applied, rainfall and temperature. Length of residual control will decrease as temperature and precipitation increase.

IMPORTANT: Contact with spray or spray drift of this product may cause severe injury or destruction of certain desirable plants, especially herbaceous species including bedding plants or direct seeded annual and perennial flowers. Therefore, do not apply this product over the top of ornamental plants growing in the landscape, and do not allow spray of this product to contact, drift or splash from soil onto foliage, green stems, exposed roots or fruit of desirable plants. Avoid application of this product under conditions that favor drift of spray onto desired ornamentals or turfgrass. The use of spray shields that limit the plant exposure to this product is directed when applying this product near desirable plants.

Do not apply this product around landscape ornamentals until plants have been actively growing for at least 30 days after transplanting, or for at least two months before ornamentals will be planted into treated areas.

PREEMERGENCE APPLICATION (NO WEEDS ARE PRESENT)

Mix 0.18 to 0.27 fl oz (5.3 to 8.1 ml) of this product per gallon of spray solution, and apply 1 gallon of spray solution to 1,000 square feet (8 to 12 fl oz/A) prior to weed germination (see Backpack Application table for more options and details). Apply this product to weed free soil, mulch or gravel surfaces. Moisture is necessary to activate this product on soil for residual weed control. When applied before weed germination, this product will control the broadleaf weeds and grasses listed in WEEDS CONTROLLED.

Established landscape ornamentals have shown tolerance to this product **only** when applied to the soil at the base of the plant. For maximum plant safety when using around desirable ornamentals, direct applications of this product to the soil, and leave a sufficient untreated buffer to ensure spray solution does not contact desired plants. **Do not** harvest fruit or nuts from treated trees within one year of application.

POSTEMERGENCE APPLICATION (WEEDS ARE PRESENT)

Mix 0.18 to 0.27 fl oz (5.3 to 8.1 ml) of this product per gallon of spray solution (8 to 12 fl oz/A), and apply 1 gallon of spray solution to 1,000 square feet to actively growing weeds (see calibration chart for backpack sprayers). Tank mixing this product with glyphosate will increase the spectrum of postemergence weed control over this product alone, provide faster postemergence weed control than glyphosate alone, and provide preemergence and postemergence control of the broadleaf weeds and grasses listed in WEEDS CONTROLLED.

Established landscape ornamentals have shown tolerance to applications of this product plus glyphosate **only** when applied to the soil at the base of the plant, and sprays **do not** directly contact or drift onto desirable plants. For maximum plant safety when using around desirable ornamentals, direct applications of this product plus glyphosate towards the soil, and leave a sufficient non-treated buffer to ensure spray solution does not contact desired plants.

Thorough spray coverage of weeds is necessary to maximize weed control. Spray coverage must be uniform, but **do not** spray to the point of runoff.

IMPORTANT: Completely read and follow the glyphosate label. When tank mixing this product with other products, always follow the most restrictive use conditions on either label.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per acre per year.
- Do not re-apply this product within 30 days.
- Do not harvest fruit or nuts from treated trees within one year of application.

ON DORMANT BERMUDAGRASS GROWN ON RESIDENTIAL SITES, GOLF COURSES, SOD PRODUCTIONS AND SIMILAR AREAS DIRECTIONS FOR USE

This product may be applied as a single or split application to well established dormant Bermudagrass. This product will provide preemergence and early postemergence control of annual bluegrass, chickweed, henbit and other winter annual weeds. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. This product will also provide preemergence control of crabgrass, goosegrass and other summer annual weeds. This product may be applied to dormant turfgrass in such areas as apartment complexes, golf courses, sod farms, roadsides, sports fields, campgrounds, office complexes, parks, parking areas, recreational sites, schools, residential turf and other similar sites. Bermudagrass exhibits tolerance to this product only when applied to semi-dormant or completely dormant turf in the late fall and before active growth resumes in the late winter/early spring. Application of this product to actively growing turfgrass (warm season and cool season) or during green-up may cause unacceptable injury.

BROADCAST APPLICATIONS

Apply 8 to 12 fl oz of this product (0.25 to 0.38 lb ai) per broadcast acre as a preemergence (to weed emergence) application. If weeds are present at the time of application apply this product plus an adjuvant (0.25% v/v non-ionic surfactant). Make postemergence (to weed emergence) applications of this product when weeds are actively growing and no larger than 2 inches in height. Thorough spray coverage is necessary to maximize the postemergence activity of this product. When applied after weed germination, this product will provide preemergence and postemergence control of broadleaf weeds and grasses. See WEEDS CONTROLLED section for a list of broadleaf weeds and grasses. Postemergence weed control with this product may be more effective on certain weed species, and may not control mature, stressed or hardened off weeds that are not actively growing at the time of application.

This product will provide best control of annual bluegrass when applied in the late fall while plants are small. Control may be less effective when applied in the winter during cold conditions when weeds are not actively growing. A second application of this product may be required to provide adequate season-long annual bluegrass control. This product will provide best control of crabgrass, goosegrass and other summer annual weeds when applied in the late winter before turfgrass resumes active growth.

TANK MIXING WITH OTHER TURFGRASS HERBICIDES

This product may be tank mixed appropriately labeled metsulfuron-methyl products.

USE AROUND BENTGRASS AND POA GREENS

This product has limited potential for lateral movement on level terrain, but can potentially move down slope after excessive rainfall and effect sensitive turf species including bentgrass and *Poa trivialis*. When applied upslope from bentgrass greens or Bermudagrass greens overseeded with *Poa trivialis*, allow an adequate buffer zone between greens and the treated area. If uncertain about the size of the buffer, 15 feet is suggested.

Risk of movement is decreased when this product is applied to soil at less than field capacity. Avoid application when heavy rain is imminent or when the soil is saturated.

USE RESTRICTIONS

- Do not apply more than 12 fl oz (0.38 lb ai) of this product per acre per application.
- Do not apply more than 24 fl oz (0.75 lb ai) of this product per acre per year.
- Do not apply more than 2 applications at 12 fl oz (0.38 lb ai) per acre or 3 applications at 8 fl oz (0.25 lb ai) per year.
- Exercise good judgment when applying to dormant turfgrass until familiarity is gained with this product.
- Do not apply to golf course putting greens.
- Do not apply to warm season turfgrass that has been overseeded with cool season turfgrass (ex. perennial rye, *Poa trivialis*).
- Do not irrigate within 1 hour before or after application.
- Do not apply if rain is expected within 1 hour after application.
- Do not mow turfgrass within 12 hours after application.
- Do not apply within 30 days prior to cutting or lifting sod.
- Do not re-apply this product within 30 days.
- Do not apply in fall before turfgrass has ceased active growth or in late winter/early spring after turfgrass has resumed active growth.

USE PRECAUTIONS

- Allow 8 weeks between application and seeding or sodding of turfgrass.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Store in a cool, dry, secure place. Do not put formulation or dilute spray solution into food or drink containers. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. Not for use or storage in or around the home. For help with any spill, leak, fire or exposure involving this material, call day or night **CHEMTREC (800) 424-9300**.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Nonrefillable Containers 5 gallons or less:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Nonrefillable containers larger than 5 gallons:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Refillable containers larger than 5 gallons:] [Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.]

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

[PROMENADE® is a registered trademark of ADAMA group company.]

[FLUMIGARD® is a registered trademark of ADAMA group company.]

[EPA insert new approval date]

Rev 020926

[OPTIONAL MARKETING/COMPANY GRAPHICS]

[The company graphics depicted below may or may not appear on final product labeling]



[ADAMA]

[A member of the ADAMA Group]

[A member of Adama Consumer and Professional Solutions]

{[LANGUAGE ON LABEL AFFIXED TO CONTAINER]}

FLUMIOXAZIN	GROUP	14	HERBICIDE
-------------	-------	----	-----------

Alligare FLUMI SC

[Alternate Brand Names: Alligare FLUMIGARD® SC Herbicide; PROMENADE® SC Herbicide; PROMENADE® SC Specialty Herbicide]

ACTIVE INGREDIENT:

Flumioxazin*..... 42.0%

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*(2-[7-flouro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isindole-1,3(2H)-dione)

Alligare FLUMI SC contains 4 pounds flumioxazin per gallon.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container. Store in a cool, dry, secure place. Do not put formulation or dilute spray solution into food or drink containers. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. Not for use or storage in or around the home. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC (800) 424-9300.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Nonrefillable Containers 5 gallons or less:] [Nonrefillable container.

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Nonrefillable containers larger than 5 gallons:] [Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure

rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[Refillable containers larger than 5 gallons:] [Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.]

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

EPA Reg. No. 81927-78

EPA Est. No.

Manufactured for:

Alligare, LLC
PO Box 97
Deer Park, TX 77536

[Lot][Batch No.]_____

Net Contents: