

81943-24

12/18/2008

58437

1/23



**U.S. ENVIRONMENTAL PROTECTION AGENCY**  
**Office of Pesticide Programs**  
**Registration Division (7505P)**  
**Ariel Rios Building**  
**1200 Pennsylvania Ave., NW**  
**Washington, D.C. 20460**

**NOTICE OF PESTICIDE:**

☒ Registration  
☐ Reregistration  
 (under FIFRA, as amended)

EPA Reg. Number:

**81943-24**

Date of Issuance:

**DEC 18 2008**Term of Issuance: **Conditional**

Name of Pesticide Product:

**Pegasus 82.5 DF**

Name and Address of Registrant (include ZIP Code):

**Phoenix Environmental Care, LLC**  
**P.O. Box 370**  
**Valdosta, GA 31603-0370**

**Mailed to: Gary Orr**  
**Agent for Phoenix Environmental Care**  
**Rivendell Consulting USA LLC**  
**400 East Jane Street**  
**Valdosta, GA 31601**

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA section 4.
2. You must submit the following conditional data listed below (run under GLP), before the due date of March 1, 2010:
  - a. Storage Stability (830.6317) and Corrosion Characteristics (830.6320) data.

(Page 1 of 3)

Signature of Approving Official:

Date:

**DEC 18 2008**

**Tony Kish**  
**Product Manager 22**  
**Fungicide Branch**  
**Registration Division (7505P)**

3. You must submit two copies of a final printed label within 45 days from the date of this notice which incorporates the following changes:
- a. Change the product registration number to "EPA Reg. No. 81943-24"
  - b. On page 1 in the **FIRST AID** box delete the "•" after IF IN EYES
  - c. On page 1 change the name of the company to "Phoenix Environmental Care LLC" and add the street address.
  - d. On page 2 change the section **HAZARDS TO HUMANS AND DOMESTIC ANIMALS** to read "Corrosive. Causes irreversible eye damage. May be fatal if inhaled. Harmful if absorbed through skin. Do not get in eyes on skin, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Do not breathe dust. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."
  - e. On page 3 before the last sentence in the paragraph at the top of the page (ie. before "Use of this product in areas..."), add "This chemical is known to leach through soil into groundwater under certain conditions as a result of label use."
  - f. Page 6 and 7 in the chemigation section, add spaces between paragraphs for ease of reading.
  - g. On page 8 in the **PHI** column, after **Beans (Dry)** change "14 7" to "14"
  - h. On pages 8 through 12 change the column header from "Acre Rate Per" to "Rate per Acre"
  - i. On page 9 in the **Acre Rate Per** column, after **Corn (sweet)** change "1.82 to 2.73 pounds" to "1.36 to 1.82 pounds"
  - j. Page 9, as per the RED, the maximum cranberries rate is 5 lbs ai or 6.06 lbs product, so change the 6.36 rate to 6.06.
  - k. On page 12 in the 4<sup>th</sup> block on the right hand side of the page change "Pods 1-11/2 inches" to "Pods 1-1/2 inches".
  - l. On page 12, as per the RED, change the soybean maximum rate from 2.27 to 2.18. Similarly for tomatoes, change 2.75 to 2.61.
  - m. On page 14 in the **Stone Fruits** row delete the horizontal line in the **PHI, RATE PER Acre** and **Spray Volume** columns

- n. On page 16 under **TURF GRASSES** change "Use of this product on home lawns in prohibited." to "Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, recreational park athletic fields, athletic fields located on or next to schools (ie. elementary, middle and high schools), campgrounds, churches, and theme parks."
- o. On page 18 in the box for **Foliage Plants** change the first entry from "glaonema" to "Aglaonema"
- p. On page 19 in the third paragraph under **CONDITION OF SALE:**
  - i. Change "...when used in accordance with directions under normal use conditions." to "...when used in accordance with label directions under normal use conditions" in the third and fourth lines.
  - ii. Delete the words "or under abnormal conditions" from the fifth line in the third paragraph.
  - iii. In front of the last sentence of the second paragraph (ie "All such risks", add "To the extent consistent with applicable law".
  - iv. By April 1, 2009, submit a separate notification or amendment, which revises the "Container Disposal" section to be in harmony with PR 2007-4.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with comments" is enclosed for your records.

Sincerely,

Tony Kish,  
Product Manager, Team 22  
Fungicide Branch  
Registration Division (7505P)

Enclosures: Label stamped "Accepted with Comments"  
Similarity Clinic Determination DP351965 dated May 27, 2008  
Product Chemistry Review DP351937 dated April 29, 2008

## Pegasus 82.5 DF

	<b>% BY WT.</b>
<b>ACTIVE INGREDIENT:</b> Chlorothalonil (tetrachloroisophthalonitrile) . . . . .	82.5%
<b>OTHER INGREDIENTS:</b> . . . . .	17.5%
<b>TOTAL:</b> . . . . .	100.0%

### KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth to mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES •</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have affected person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
<b>EMERGENCY PHONE NUMBERS</b>	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center (human health) (800) 345-4735 ASPCA (animal health)
<b>NOTES TO PHYSICIAN:</b> Probable mucosal damage may contraindicate the use of gastric lavage. Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.	

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
DEC 18 2008**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

Net contents: \_\_\_\_\_ pounds

81943-24

Phoenix Environment Care LLC  
P.O. Box 370 • Valdosta, GA 31603-0370

EPA Reg. No. 81043

## **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. DO NOT get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Avoid contact with eyes, skin or clothing. Do not breathe spray mist.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Mixers, loaders, applicators and all other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Protective eye wear;
- Chemical-resistant gloves made of waterproof material, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinyl chloride, or viton; if you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart;
- A NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

### **USER SAFETY REQUIREMENTS**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

### **ENGINEERING CONTROLS:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **USER SAFETY RECOMMENDATIONS**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates and wildlife. DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwater or rinsate. Chlorothalonil can contaminate surface water through spray drift. DO NOT apply when weather conditions favor drift from treated areas. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, and areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain

to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water. Chlorothalonil degradates are known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

## **DIRECTIONS FOR USE**

### **General Precautions and Restrictions**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, or pets, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation. Use of this product on home lawns is prohibited.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- waterproof gloves,
- shoes plus socks,
- and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted-entry interval expires after 12 hours, for the next 6 1/2 days entry is permitted only when the following safety measures are provided:

At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.

Workers must be informed, in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes;
- that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes;
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water; and
- how to operate the eyeflush container.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170): DO NOT enter or allow others to enter into treated areas until spray deposits have dried.

This product must not be applied within 150 feet (for aerial and air-blast applications), or 25 feet (for ground applications) from marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Avoiding spray drift at the application site is the responsibility of the applicator. The interactions of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

## Aerial Drift Advisory Information

### INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable conditions (see Wind, Temperature).

### CONTROLLING DROPLET SIZE

- Volume- Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure- Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces
- larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles- Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation - Orientating nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle type- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

### BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

### APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

8/2

### **SWATH ADJUSTMENT**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, small drops, etc.).

### **WIND**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **TEMPERATURE AND HUMIDITY**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### **TEMPERATURE INVERSIONS**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **Integrated Pest Management**

PEGASUS 82.5 DF is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. PEGASUS 82.5 DF is recommended for use in programs that are compatible with the principles of Integrated Pest Management (IPM), including the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

## **Fungicide Resistance Management**

PEGASUS 82.5 DF is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. PEGASUS 82.5 DF, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of PEGASUS 82.5 DF in programs which seek to minimize the occurrence of disease resistance to other fungicides.

## **Mixing, Loading and Applying**

PEGASUS 82.5 DF is intended to be diluted into water, and then applied to crops by typical agricultural spraying techniques. **Always apply PEGASUS 82.5 DF in sufficient water to obtain thorough, uniform coverage of foliage and crop surfaces intended to be protected from disease.** Spray volume to be used will vary with crop and amount of plant growth. Spray volume should normally range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Slowly invert container several times to assure uniform mixture. Measure the required amount of PEGASUS 82.5 DF and pour into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations. Do not use on greenhouse-grown crops.



### **Tank Mixing**

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not combine PEGASUS 82.5 DF in sprayer tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine PEGASUS 82.5 DF with Dipel 4L, Foil, Triton AG-98, Triton B-1956 or Latron B-1956, as phytotoxicity may result from the combination when applied to the crops on this label. DO NOT tank mix Pegasus 82.5 DF with oil, or with any adjuvants which contain oil as their principal ingredient. Do not use with Copper-Count N in concentrated spray suspensions.

### **Applications through Sprinkler Irrigation Systems (Chemigation)**

Application through sprinkler irrigation systems is recommended only for those specific crops for which the notation "chemigation OK" is listed on this label.

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). DO NOT apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject PEGASUS 82.5 DF into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

PEGASUS 82.5 DF may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, and then refer to the appropriate directions provided for each type.

#### **A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment**

For injection of pesticides, these continuously moving systems must use a metering pump, such as a positive displacement injection pump of either diaphragm or piston type, constructed of materials that are compatible with pesticides, fitted with a system interlock, and capable of

injection at pressures approximately 2 to 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix recommended amount of PEGASUS 82.5 DF for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until PEGASUS 82.5 DF has been cleared from last sprinkler head.

**B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment**

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of PEGASUS 82.5 DF for acreage to be covered with water so that the total mixture of PEGASUS 82.5 DF plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required. PEGASUS 82.5 DF can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until PEGASUS 82.5 DF has been cleared from last sprinkler head.

**Application Rates**

Dosage rates on this label indicate pounds of PEGASUS 82.5 DF per acre, unless otherwise stated. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

For each listed crop, the maximum total amount of chlorothalonil active ingredient (lbs a.i./A) which may be applied per acre of that crop (or crop group) during each growing season is given in bold print within a box beneath the crop name. For each crop use situation listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased.

Crop	PHI (days)	Diseases	Acre Rate Per	Application Directions
Asparagus 9.0 lbs a.i./A	190 (120 in California)	Rust, Purple spot, Cercospora leaf blight	1.82 to 3.64 pounds	Begin applications after harvest of spears, when conditions favor disease development on ferns, generally when leaf wetness occurs. Repeat applications at 2 to 4 week intervals until ferns are no longer productive. Use the high rate and shortest interval when conditions favor disease.
Bean (Snap) 9.0 lbs a.i./A	7	Rust	1.25 to 2.73 pounds	Begin applications during early bloom stage or when disease first threatens and repeat at 7 day intervals or as necessary to maintain control-
		Botrytis blight (gray mold)	2.73 pounds	
Beans (Dry)* 6.0 lbs a.i./A Chemigation OK	14 7	Rust, Anthracnose, Downy mildew, Cercospora leaf spot (blackeye only), Botrytis blight (gray mold) Ascochyta blight	1.25 to 1.82 pounds 2.73 pounds	Begin applications during early bloom stage and repeat at 7 to 10 day intervals. For use only on beans to be harvest-ed dry with pods removed.
Cabbage, Chinese Cabbage (tight-headed varieties only), Cauliflower, Broccoli, Chinese Broccoli, Brussels Sprouts 12.0 lbs a.i./A)	7	Alternaria leaf spot, Downy mildew	1.36 <sup>a</sup> pounds	Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control.
		Ring Spot	1.82 pounds	
Carrot 15.0 lbs a.i./A Chemigation OK	0	Cercospora (Early) blight, Alternaria (Late) blight Ring spot	1.36 to 1.82 pounds	Start applications when disease threatens and repeat at 7 to 10 day intervals or as necessary to maintain control.
Celery 18.0 lbs a.i./A	7	Cercospora (Early) blight, Septoria (Late) blight, Basal stalk rot (Rhizoctonia solani)	Start applications shortly after crop emergence or when transplants are set in the field. For the indicated rates, re-apply at:	
			0.91 to 1.36 pounds	3 to 5 day intervals
			1.82 to 2.73 pounds	7 day intervals

Crop	PHI (days)	Diseases	Acre Rate Per	Application Directions
Corn (sweet), Corn grown for seed 9.0 lbs a.i./A	14	Helminthosporium leaf blights, Rust	1.82 to 2.73 pounds	Begin applications when conditions favor disease development and repeat at 7 day intervals. Do not allow livestock
Cranberry 15.0 lbs a.i./A Chemigation OK; solid set systems only	50	Upright dieback Fruit rots, Lophodermium leaf & twig blight	3.64 to 6.36 pounds	Apply at budbreak to early bloom and repeat at 10 to 14 day intervals. Under severe disease conditions use the high rate on a 10 day schedule. DO NOT apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application.
Cucurbits: Cucumber, Cantaloupe, Muskmelon, Honeydew melon, Watermelon, Squash, Pumpkin 15.75 lbs a.i./A Chemigation OK	0	Anthracnose, Downy mildew, Target spot	1.36 to 1.82 pounds	Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray interval. Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply PEGASUS 82.5 DF to watermelons when any of the following conditions are present: 1. Intense heat and sunlight; 2. Drought conditions; 3. Poor vine canopy; 4 Other crop and environmental conditions which may be conducive to increases natural sunburn DO NOT combine PEGASUS 82.5 DF with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.
		Cercospora leaf spot, Gummy stem blight (black rot), Alternaria leaf blight, Scab, Powdery mildew (Sphaerotheca only)	1.82 to 2.73 pounds	

12/23

Crop	PHI (days)	Diseases	Acre Rate Per	Application Directions			
Grasses Grown for Seed 4.5 lbs a.i./A	14	Stem rust, Leaf rust, Stripe rust, Septoria leaf spot, Glume blotch, Bipolaris and Drechslera leaf spots	0.91 to 1.36 pounds	Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat application at 14 day intervals. DO NOT allow livestock to graze in treated areas. Do not feed straw, seed or seed screenings to livestock			
		Selenophoma eyespot	0.91 to 1.82 pounds				
Mint 3.0 lbs a.i./A	80	Rust, Septoria leaf spot	1.25 pounds	Being applications when emerging plants are 4 to 8 inches high. Repeat applications at 7 to 10 day intervals or as necessary to maintain control. Based on available residue data, use of this product on mint is restricted to Indiana, Michigan and Wisconsin.			
Mushroom beds	5 Do not apply after first break (harvest)	Verticillium brown spot and dry bubble	Rate per 1,000 sq. ft. of bed surface  2.5 to 5 oz.	Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of bed surface. Make two application. Apply the high rate in the first application and the low rate in the second application. The first application should be made within two days after top-dressing the spawn-colonized mushroom compost with a casing layer. The second applications should be made at pinning. Make no more than two applications per cropping cycle. Do not apply more than 0.4 lbs active ingredient chlorothalonil per 1,000 sq. ft. per cropping cycle.			
Onion (dry bulb), Garlic 15.0 lbs a.i./A	7	Botrytis leaf blight or blast, Purple blotch	0.91 to 1.82 pounds	PEGASUS 82.5 DF is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:			
					Low Disease Hazard & Prior to Infection	Low Disease & Some Disease Present	High Disease Hazard
				Rate per Acre	0.91 pounds	1.25 pounds	1.82 pounds
				Frequency	10 days	7 to 10 days	7 days
		Neck rot	1.25 to 1.82 pounds	For suppression of neck rot (Botrytis spp.) during storage, make a minimum of three weekly applications prior to lifting.			

13/2

Crop	PHI (days)	Diseases	Acre Rate Per	Application Directions
Onion (green bunching), Leek, Shallot, Onion grown for seed 6.7 lbs a.i./A	14 (green onion, leek, shallot)	Botrytis leaf blight or blast, Purple blotch, Downy mildew (suppression)	1.36 to 2.73 pounds	Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persists. If additional disease is needed before harvest, use another registered fungicide
Parsnip 6.0 lbs. a.i./A	10	Alternaria leaf spot, Downy mildew, Anthracnose, Botrytis blight (gray mold), Bottom rot (Rhizoctonia)	1.36 to 1.82 pounds	Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule.
Peanut 9.0 lbs a.i./A Chemigation OK	14	Early leafspot (Cercospora)	0.91 to 1.36 pounds	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals. Do not allow livestock to graze in treated areas. Do not feed hay or threshings from treated fields to livestock.
		Late leafspot (Cercosporidium) Rust, Web blotch	1.36 pounds	
Potato 11.25 lbs a.i./A Chemigation OK	7	Late blight, Early Blight, Botrytis vine rot	0.68 pound  THEN 0.91 to 1.36 - pounds	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat application at 7 to 10 day intervals. Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur: --Vines close within the rows; --Late blight forecasting measures 18 disease severity values (DSV); --The crop reaches 300 P-days  Increase water spray volume as canopy density increase. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.

2/1

Crop	PHI (days)	Diseases	Acre Rate Per	Application Directions		
Soybean 4.5 lbs a.i./A Chemigation OK	42	Anthracnose, Diaporth pod and stem blight, Frog eye leaf spot (Cercospora soja), Purple seed stain, Cercopora leaf blight (ercospora kikuchii), Septoria brown spot, Rust (Phakopsora pachyrhizi)	See tables below for rates and timing of applications. Use the three application program in areas having a history of moderate to severe disease intensity. Do not feed soybean hay or threshings from treated fields to livestock.			
					Determinate southern varieties	Indeterminate northern varieties
			1.36 to 2.27 pounds	2-Application Program	Early pod set (R3) Seed Formation (R5)	Pods 1-1 1/2 inches Then 14 days later
			0.91 to 1.82 pounds	3-Application Program	Early flowering (R1) Early pod set (R3) Seed formation (R5)	One week after first flowering, then at 14 day intervals
		Stem canker (Diaporthe phaseolorum var. caulvora)	0.91 pound	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease make a second and third application at 14 day intervals.		
Tomato 15.1 lbs. a.i./A Chemigation OK; solid set or portable wheel move systems only	0	FOLIAGE(apply every 7- 10 days): Early blight, Leaf blight, Gray leaf spot, Gray leaf mold, Septoria leaf spot, Target spot	1.25 to 1.82 pounds	Begin applications when dew or rain occurs and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe.  PEGASUS 82.5 DF may be combined in the spray tank with EPA-registered pesticide products that claim copper as an active ingredient and are labeled for control of bacterial disease of tomatoes. Check the copper manufacturer's label for specific instructions, precautions and limitations prior to mixing with PEGASUS 82.5 DF.		
		FRUIT (apply every 7-14 day beginning at fruit set): Anthracnose, Alternaria fruit rot (black mold), Botrytis gray mold, Late blight fruit rot, Rhizoctonia fruit rot	1.82 to 2.75 pounds			

15/23

## TREE AND ORCHARD CROPS

Apply this product in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, this product may be applied with aircraft using at least 20 gallons of spray per acre. When concentrate sprays are used or when treating non-bearing or immature trees, use the lower rate of this product listed for the crop being treated.

DO NOT allow livestock to graze in treated areas.

**DO NOT apply Pegasus 82.5 DF within one week before or after application of oil or an oil-based pesticide.**

Crop	PHI (days)	Diseases	RATE PER Acre	Spray Volume (gallons/acre)	Application Directions
Almonds 18.75 lbs a.i./A	150	Shothole, Scab, Anthracnose	3.64 pounds	20 (concentrate) to 300 (full dilute)	For best control of shothole apply at leaf fall in late autumn. Apply also at budbreak to protect newly emerging leaves, and at shuck (jacket) split to prevent nut infections and to control scab and anthracnose.
		Brown rot blossom and twig blight			Apply at popcorn (pink bud) and at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
Blueberry 9.0 lbs a.i./A	42	Mummy Berry, Anthracnose	2.73 to 3.64 pounds	20 (concentrate) to 100 (full dilute)	Begin applications at budbreak (green tip). Repeat applications until early bloom at 10 day intervals. DO NOT apply after early bloom; otherwise phytotoxicity may occur to the developing fruit.
Filberts (Hazelnuts) 9.0 lbs a.i./A	120	Eastern filbert blight	3.64 pounds	20 (concentrate) to 400 (full dilute)	Begin applications at leaf bud break and repeat at 2 to 4 week intervals. Based on available residue data, use of this product on filberts is restricted to Oregon.
Mango 24.0 lbs a.i./A	21	Anthracnose	1.82 to 3.18 pounds	100 (full dilute)	Begin applications at early bloom and repeat at 7 to 14 day intervals until early fruit development. Use the high rate and apply weekly when conditions favor disease.
Papaya 6.75 lbs a.i./A	14	Alternaria fruit spot, Anthracnose, Stem end rot	1.82 to 2.73 pounds	20 (concentrate) to 150 (full dilute)	Apply with ground equipment only. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development.

3/2/01



Crop	PHI (days)	Diseases	RATE PER Acre	Spray Volume (gallons/acre)	Application Directions
Passion Fruit (Hawaii only) 7.5 lbs a.i./A	7	Alternaria fruit and leaf spot (brown spot)	1.82 pounds	20 (concentrate) to 100 (full dilute)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications before fruit spots appear (April to July) and re-apply at 14 day intervals until weather conditions no longer favor disease development.
Pistachio 22.5 lbs a.i./A	14	Shoot & panicle blight, Blossom & shoot blight, Late blight, Leaf blight	5.45 pounds	50 (concentrate) to 200 (full dilute)	Apply when trees begin to blossom, and then re-apply at full bloom for optimal protection against shoot and panicle blights. If conditions are favorable for late blight or leaf spot infections, repeat applications at 4 week intervals. Use the high rate when abnormally wet or cloudy weather conditions prevail. NOTE: Use of this product in the manner described may result in specking or reddening of the fruit shell (epicarp). This effect appears to be superficial, and may not result in any change in nut quality.
Stone Fruits: Peach, Nectarine, Apricot, Cherry, Plum, Prune 15.5 lbs a.i./A	Do not apply after shuck split	Leaf curl	2.84 to 3.76 pounds	20 (concentrate) to 300 (full dilute)	For best control apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels use the high rate and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of PEGASUS 82.5 DF for control of leaf curl may be made at any time prior to budswell the following spring.
		Shothole, Brown rot blossom blight, Lacy (russet) scab on prune, Cherry leaf spot, Scab			Make one application at budbreak or popcorn (pink, red or early white bud). If weather conditions favor disease, make a second application 10 days later (full bloom to petal fall). Apply at shuck split to prevent infections on young fruit. If additional disease control is needed after shuck split and before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10-14 days later.

Crop	PHI (days)	Diseases	RATE PER Acre	Spray Volume (gallons/acre)	Application Directions
Conifers 16.5 lbs a.i./A	N/A	Swiss needlecast	2.5 to 5 pounds	5 to 10 (concentrate ground or aircraft) to 100 (dilute)	Single application technique: In Christmas tree plantations or forest stands make one application in the spring when new shoot growth is ½ to 2 inches in length.
		Scleroderris canker (pines), Swiss needlecast	1.36 to 2.5 pounds		Make the first application in spring when new shoot growth is ½ to 2 inches in length. Make additional application at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule
		Sirococcus tip blight	1.82 to 3.18 pounds		
		Rhizosphaera needlecast (spruces), Scirrhia brown spot (pines)	5 pounds		
		Cyclaneusma and Lophodermium needlecasts (pines)	2.5 to 5 pounds		Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, application may be suspended, and then resumed upon next occurrence of needle wetness.
		Rhabdocline needlecast (Douglas-fir)	1.36 to 2.5 pounds		Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
		Botrytis seedling blight, Phoma twig blight	2.5 pounds		Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
		Autoecious needle rust (Weir's cushion rust) (spruces)	5 pounds		Begin applications when 10% of buds have broken and repeat twice thereafter at 7 to 10 day intervals.

## TURFGRASSES

Use of this product on home lawns is prohibited. Sodfarm turf treated with chlorothalonil prior to harvest must be mechanically cut, rolled and harvested. Do not apply more than the following totals of chlorothalonil active ingredient from all registered product sources to the indicated types of turfgrass:

TYPE OF TURFGRASS	TOTAL CHLOROTHALONIL ACTIVE INGREDIENT PER ACRE PER YEAR
Golf Course Greens	73 lbs
Golf Course Tees	52 lbs
Golf Course Fairways	26 lbs
Sod Farms	13 lbs
Other Turf	26 lbs

Apply PEGASUS 82.5 DF in 30 to 100 gallons of water per acre. Apply with ground equipment only.

Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below. DO NOT mow or irrigate after treatment until spray deposit on turfgrass is thoroughly dry. PEGASUS 82.5 DF should always be used in conjunction with good turf management practices.

DISEASES* CONTROLLED	INTERVAL OF APPLICATION	GOLF COURSE GREENS & TEES RATE PER 1,000 SQ.FT.	GOLF COURSE GREENS & TEES RATE PER 1,000 SQ.FT.
1. Dollar spot 2. Brown patch 3. Leaf spot, Melting-out, Brown blight 4. Gray leaf spot	7-14 days	1.82 to 3.25 ounces (4.1 to 7.3 lbs a.i./acre)	5 to 8.9 pounds (4.1 to 7.3 lbs a.i./acre)
5. Red thread 6. Anthracnose 7. Copper spot 8. Stem rust (bluegrass) 9. Dichondra leaf spot	7 days or 14 days	3.25 ounces or 5 ounces (7.3 or 11.3 lbs a.i./acre)	8.9 pounds or 13.7 pounds (7.3 or 11.3 lbs a.i./acre)

\*Diseases listed are caused by fungi, some of which are named as follows:

1. Dollar spot: *Sclerotinia homeocarpa*; *Lanzia* or *Moellerodiscus* spp.
2. Brown patch: *Rhizoctonia solani*, *R. zeae*, *R. cerealis*
3. Leaf spots, Melting-out, Brown blight: *Drechslera* spp. (including *D. poae*, *D. siccans*), *Bipolaris sorokiniana*, *Curvularia* spp.
4. Gray leaf spot: *Pyricularia grisea*, *P. oryzae*
5. Red thread: *Laetisaria fuciformis*
6. Anthracnose: *Colletotrichum graminicola*
7. Copper spot: *Gloeocercospora sorghi*
8. Stem rust: *Puccinia graminis*
9. Dichondra leaf spot: *Alternaria* spp.

**Gray Snow Mold** caused by *Typhula* spp.: Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1,000 square feet). Apply 5 ounces of PEGASUS 82.5 DF per 1,000 square feet of turf area (13.7 pounds per acre). Application must be made before snow cover in autumn. If snow cover is intermittent or lacking during the winter, re-apply PEGASUS 82.5 DF at monthly intervals until Gray Snow Mold conditions no longer prevail. In areas where Pink Snow Mold

20  
2

(Microdochium or Fusarium patch) is likely to occur, apply PEGASUS 82.5 DF at 5 ounces in combination with products containing iprodione at 1.82 ounces active ingredient, per 1,000 square feet of turf area. Read and observe all label directions for products containing these active ingredients.

**Fusarium (Microdochium) Patch:** PEGASUS 82.5 DF is effective against Fusarium patch only in areas where snow cover is intermittent or lacking during the winter. Apply 5 ounces of PEGASUS 82.5 DF per 1,000 square feet of turf area. Begin applications in late autumn and re-apply at 21 to 28 day intervals until conditions favorable for Fusarium patch no longer prevail.

**Algal scum:** Apply PEGASUS 82.5 DF at 1.82 to 3.25 ounces per 1,000 square feet on a 7 to 14 day schedule. When colonies of algae are well established, every attempt should be made to dry out the afflicted area. Once dry, spiking or verticutting should be done to enhance turfgrass recovery in conjunction with the use of PEGASUS 82.5 DF. Several applications of PEGASUS 82.5 DF at the high rate may be necessary for turfgrass recovery. When environmental conditions are favorable for algae growth, a preventive program with PEGASUS 82.5 DF will suppress recolonization of the turf.

### ORNAMENTAL PLANTS

Apply PEGASUS 82.5 DF at a rate of 1.25 pounds per 100 gallons of water unless other directions are given in the tables below. Apply enough diluted spray per acre to provide thorough coverage of all plant parts that are intended to be protected from disease, generally ranging from 20 to 150 gallons per acre. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply PEGASUS 82.5 DF at 7 day intervals. **DO NOT apply more than a total of 36.4 lbs chlorothalonil active ingredient per acre per growing season on field-grown ornamentals.**

Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN.**

This product may be used in greenhouses. DO NOT use mistblowers or high pressure spray equipment when making applications of this product in greenhouses.

PEGASUS 82.5 DF is recommended for control of fungal diseases referred to by numbers in parentheses following each type of ornamental plant. The user should test for possible phytotoxic responses, using recommended rates on each type of ornamental plant on a small area prior to widespread use. Applications made during, bloom may damage flowers and/or fruits.

### ORNAMENTALS RECOMMENDED FOR TREATMENT WITH PEGASUS 82.5 DF

Broadleaf Shrubs and Trees		
Andromeda ( <i>Pieris</i> ) (4)	Flowering almond (1,2)	Oregon-grape ( <i>Mahonia</i> ) (6)
Ash ( <i>Fraxinus</i> ) (1)	Flowering cherry (1,2)	Red-tip ( <i>Photinia</i> ) (1)
Aspen (1)	Flowering peach (1,2)	Poplar (1)
Azalea (1,2,4)	Flowering plum (1,2)	Privet ( <i>Ligustrum</i> ) (1)
Buckeye, Horsechestnut (1)	Flowering quince (1,2)	Rhododendron (1,2,4)
Camellia (2)	Hawthorn (1,6)	Sand cherry (1,2)
Cherry-laurel (1)	Holly (1)	Sequoia (1)
Crabapple (1,6)	Lilac (5)	Spirea (1)
Dogwood (1)	Magnolia (1)	Sycamore, Planetree (1)
Eucalyptus (3)	Maple (1)	Viburnum (5)
Euonymus (1)	Mountain laurel (1)	Walnut ( <i>Juglans</i> ) (1)
Firethorn ( <i>Pyracantha</i> ) (1)	Oak (red group only) (1,7)	

21  
22

Flowering <sup>a</sup> Plants and Bulbs		
Arabian violet (2)	Gladiolus (1,2)	Petunia (1,4)
Begonia (1)	Hollyhock (6)	Phlox (1)
Carnation (1,2)	Hydrangea (foliage only) (1,6)	Poinsettia <sup>b</sup> (1)
Chrysanthemum (1,2)	Iris (1,2)	Rose <sup>c</sup> (1)
Crocus (1)	Lily (l)	Statice (1)
Daffodil (1)	Marigold (1)	Tulip (1)
Daisy (1)	Narcissus (1)	Zinnia (1,5)
Geranium (1,6)	Pansy (1)	

a/ Avoid applications during bloom period on plants where flower injury is unacceptable.

b/ Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.

c/ Use 0.9 pound of PEGASUS 82.5 DF per 100 gallons of water.

Foliage Plants		
glaonema (1)	Ficus (1)	Parlor palm ( <i>Chamaedorea</i> ) (1)
Areca palm (1)	Florida ruffle fern (1)	Peperomia (1)
Artemesia (1)	Leatherleaf fern (1)	Philodendron (1,4)
Boston fern ( <i>Nephrolepis</i> ) (1)	Lipstick plant (1)	Prayer plant ( <i>Maranta</i> ) (1)
Dracaena (1)	Ming aralia (1)	Syngonium (1)
Dumbcane ( <i>Dieffenbachia</i> ) (1)	Oyster plant ( <i>Rhoeo</i> ) (1)	Zebra plant ( <i>Aphelandrá</i> ) (1)
Fatsia ( <i>Aralia</i> ) (1)	Pachysandra (1)	

### DISEASES CONTROLLED WITH PEGASUS 82.5 DF:

1. Leafspots & Foliar Blights:		
Actinopelte leafspot	Corynespora stem & leafspots	Mycosphaerella ray blight
Alternaria leafspot or leaf blight	Curvularia leafspot	Myrothecium leafspot, brown rot
Anthracoise ( <i>Gnomonia</i> , <i>Glomerella</i> , <i>Colletotrichum</i> , <i>Discula</i> ) blights	Dactylaria leafspot	Phyllosticta leafspot
Black spot ( <i>Diplocarpon</i> )	Didymellina leafspot	Ramularia leafspot
Botrytis blights	Drechslera ( <i>Bipolaris</i> ) leafspots, inkspot	Rhizoctonia web blight
Cephalosporium leafspot	Fabraea (Entomosporium) leafspot	Scab ( <i>Venturia</i> )
Cercospora leafspot	Fusarium ( <i>Gibberella</i> ) leafspot	Septoria leafspot
Cercosporidium leafspot	Gloeosporium black leafspot	Sphaeropsis leafspot
Shothole ( <i>Stigmia</i> )	Marssonina leafspot	Stagonospora leaf scorch
	Monilinia blossom blight, twig blight	Tan leafspot ( <i>Guivularia</i> )
		Volutella leaf blight

2. Flower Spots and Blights		
Botrytis flower spot, flower blight	Monilinia blossom blight	Rhizopus blossom blight
Curvularia flower spot	Ovulinia flower blight	Sclerotinia flower blight

3. Cylindrocladium stem canker
--------------------------------

4. Phytophthora leaf blight, dieback
--------------------------------------

5. Powdery mildews:		
<i>Erysiphe cichoracearum</i>	<i>Sphaerotheca fuliginea</i>	<i>Microsphaera</i> spp.

6. Rusts:		
<i>Gymnosporangium</i> spp	<i>Pucciniastrum hydrangeae</i>	<i>Puccinia</i> spp.

7. Taphrina blister
---------------------

22  
2

### STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Pesticide Storage:** Store in a cool place. Protect from excessive heat.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** DO NOT reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable Refillable Container:** If PEGASUS 82.5 DF is packaged in a returnable refillable container, then, after use, do not rinse container. Return container intact to point of purchase. This container must only be refilled with PEGASUS 82.5 DF. DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE. Before refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn threads on closure devices. Check for leaks after refilling and before transport.

### CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Phoenix Environmental Care, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Phoenix Environmental Care, LLC and Seller harmless for any claims relating to such factors.

Phoenix Environmental Care, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Phoenix Environmental Care, LLC, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, PHOENIX ENVIRONMENTAL CARE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Phoenix Environmental Care, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PHOENIX ENVIRONMENTAL CARE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PHOENIX ENVIRONMENTAL CARE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Phoenix Environmental Care, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Phoenix Environmental Care, LLC.

Phoenix design is a registered trademark of Phoenix Environmental Care, LLC  
Pegasus is a trademark of Phoenix Environmental Care, LLC