

81964-2

11/3/2009

10f48

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



NOV 03 2009

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

ChemStarr LLC
c/o Ann Tillman
Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

Dear Ms. Tillman,

Subject: Amendment to delete residential turf applications, revised labeling per Agency letter dated 4/15/09
Acephate 75% SP
EPA Registration No. 81964-2

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

- On page 32, in the box for "Turfgrass" revise the rate for leafhoppers back to 1.0 oz (2 ³/₄ lbs/acre) as on the last stamped label dated April 12, 2007.
- On page 33, in the box for "Turfgrass" revise the special instructions for greenbugs/grasshoppers that currently reads: "Do not mow urfgrass for at least..." to read: "Do not mow turfgrass for at least..."

Please submit one final printed copy of the labeling before releasing the product for shipment. A stamped copy of the labeling is enclosed for your records. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314 or metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Venus Eagle".

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

[Water Soluble Film Label]

Acephate 75% SP
INSECTICIDE

ACTIVE INGREDIENT:	By Wt.
Acephate (O,S-Dimethyl acetylphosphoramidothioate).....	75.0%
OTHER INGREDIENTS:	<u>25.0%</u>
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

For First Aid, Precautionary Statements and Directions for Use, refer to the complete product labeling located on the outer container.

EPA Reg. No. 81964-2

ACCEPTED
With COMMENTS
 In EPA Letter Dated:
 NOV 03 2009
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act, As amended, for the
 pesticide Registered under EPA Reg. No:
81964-2

[Main Label for Product Packaged in Water Soluble Film]

40048

Acephate 75% SP

INSECTICIDE

ACTIVE INGREDIENT:	By Wt.
Acephate (<i>O,S</i> -Dimethyl acetylphosphoramidothioate).....	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
Acephate is an organophosphate, cholinesterase inhibitor	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-949-679-3535 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Acephate is a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure but decisions regarding treatment will usually need to be made before test results are available. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PROTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 81964-2
EPA Est. No.

Manufactured for:
ChemStarr LLC
21 Hubble
Irvine, CA 92618

Net Weight:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove and wash contaminated clothing before reuse. Do not allow children or pets to come into contact with treated areas until sprays have dried.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils and natural rubber ≥ 14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves for all mixers and loaders and for applicators using hand-held application equipment.

In addition, applicators using low-pressure hand wand application equipment must wear:

- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

Water-soluble packets when used correctly qualify as a closed mixing/loading system under the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(4)]. Mixers and loaders using water-soluble packets must:

- Wear the personal protective equipment required above for mixers/loaders, and
- Be provided, must have immediately available and must use in an emergency, such as a broken package, spill, or equipment breakdown the following PPE: coveralls, chemical-resistant footwear and a NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

The use of human flaggers is prohibited.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

60948

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product and its degradate are highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PRODUCT INFORMATION

Acephate 75% SP is a soluble powder used as an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in Acephate 75% SP is absorbed by plants via the roots and foliage to provide systemic control of feeding insects.

Because it is impossible to test this product for every combination of cultivar, soil and horticultural practice, it is the responsibility of the user to ensure that Acephate 75% SP is compatible with the variety or cultivar of the ornamental crop to which you are making an application under your specific conditions. Test the product on a limited scale and observe for phytotoxicity over a two-week period prior to making large-scale applications.

FAILURE TO FOLLOW THE INSTRUCTIONS IN THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made out of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

ADDITIONAL INFORMATION

- Insect pests are generally controlled more effectively by Acephate 75% SP through ingestion than by contact.
- Make repeat applications of Acephate 75% SP to maintain control only as directed.

TANK MIXES

- When tank mixing Acephate 75% SP with other products, read the entire label of each product being added to the mix and follow the most restrictive guidelines applicable to your situation.
- The user, applicator and/or application advisor accepts full responsibility for using this product with any other product or application method not expressly authorized by this label.

CHEMIGATION

- Only cranberries can have this product applied by chemigation (sprinkler irrigation) system.
- Do not apply to cranberries using any other type of irrigation system.

USE RESTRICTIONS

For use on turf, use limited to sod farms and golf courses, except when applying to mound or spot treatment for fire ant and harvester ant control, but not including residential turf.

Acephate 75% SP is NOT for indoor residential use. For greenhouse use, use is limited to commercial greenhouses for use on ornamental, floral and foliage plants and tobacco (floatbed application).

Hand Wand Restrictions

- Application by low-pressure hand wand is not registered for use in California.
- Agricultural Uses - Low-pressure hand wand equipment may *only* be used for mound treatment to control fire ants.
- Non-Agricultural Uses - Low-pressure hand wand equipment may *only* be used for mound treatment to control fire ants and on ornamental trees, shrubs and floral plants only when grown for non-agricultural or non-commercial use.
- Low-pressure hand wand equipment may *not* be used for perimeter control or wasp treatments.

80948

WATER SOLUBLE PACKAGING

NOTE: Refer to the Table of Equivalents below to calculate the number of packets to use.

MULTIPLE PACKAGING

This container holds water-soluble packets of Acephate 75% SP. Do not puncture or rupture the packets or expose them to any moisture, including handling with wet gloves, before placing the packets into the spray tank. To protect any remaining unused packets, be sure to store them in the original outer container.

PREPARATION OF SPRAY SOLUTION

1. Drop the unopened packet(s) of Acephate 75% SP into a spray tank containing at least one-half the total quantity of water required.
2. Start the agitator and continue adding water until the desired volume is reached by spraying directly at the packet(s) in order to dissolve the packet(s) and release the Acephate 75% SP.
3. Do not add any partners (such as liquid fertilizers, micronutrients or adjuvants) to the tank mix until *after* the packet(s) and its / their contents have completely dissolved.

NOTE: The water-soluble packet(s) should completely dissolve in approximately five minutes. The timing may be impacted (slowed) by insufficient agitation, cold water, or water containing high concentrations of sulfur or boron.

TABLES OF EQUIVALENTS

Amount of Acephate 75% SP	1 1/3 lbs.	1 lb.	2/3 lb.	1/3 lb.	4 oz.	2 oz.
Amount Active Contained	1 lb.	3/4 lb.	1/2 lb.	1/4 lb.	1/5 lb.	1/10 lb.

Table of Equivalents for 1/3 lb. Packet

Acephate 75% SP Application Rate (per Acre)	Acres Treated by a 1/3 lb. Water Soluble Packet
2 oz.	2.66
2.7 oz.	2.00
4 oz.	1.33
1/3 lb.	1.00
2/3 lb.	0.50
1 lb.	0.33
1 1/3 lbs.	0.25
2 2/3 lbs.	0.13
2.7 lbs.	0.12
2.75 lbs.	0.12
3 1/4 lbs.	0.10
4.1 lbs.	0.08
5.2 lbs.	0.06

Table of Equivalents for 1/2 lb. Packet

Acephate 75% SP Application Rate (per Acre)	Acres Treated by a 1/2 lb. Water Soluble Packet
2 oz.	4.00
2.7 oz.	3.00
4 oz.	2.00

Acephate 75% SP Application Rate (per Acre)	Acres Treated by a ½ lb. Water Soluble Packet
1/3 lb.	1.50
2/3 lb.	0.75
1 lb.	0.50
1 1/3 lbs.	0.38
2 2/3 lbs.	0.19
2.7 lbs.	0.19
2.75 lbs.	0.18
3 ¼ lbs.	0.15
4.1 lbs.	0.12
5.2 lbs.	0.10

Table of Equivalents for 1 lb. Packet

Acephate 75% SP Application Rate (per Acre)	Acres Treated by a 1 lb. Water Soluble Packet
2 oz.	8.00
2.7 oz.	5.93
4 oz.	4.00
1/3 lb.	3.00
2/3 lb.	1.50
1 lb.	1.00
1 1/3 lbs.	0.75
2 2/3 lbs.	0.38
2.7 lbs.	0.37
2.75 lbs.	0.36
3 ¼ lbs.	0.31
4.1 lbs.	0.24
5.2 lbs.	0.19

Table of Equivalents for 2 ½ lb. Packet

Acephate 75% SP Application Rate (per Acre)	Acres Treated by a 2 ½ lb. Water Soluble Packet
2 oz.	20.00
2.7 oz.	14.81
4 oz.	10.00
1/3 lb.	7.50
2/3 lb.	3.75
1 lb.	2.50
1 1/3 lbs.	1.88
2 2/3 lbs.	0.94
2.7 lbs.	0.93
2.75 lbs.	0.91
3 ¼ lbs.	0.77
4.1 lbs.	0.61
5.2 lbs.	0.50

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

- All aerial, ground and air-assisted/airblast application equipment must be properly maintained and calibrated using water as carrier. Do not apply this product as an ultralow (ULV) spray except as specified in the directions for use on wasteland (non-crop areas), or in any carrier other than water.
- Use the largest droplet size consistent with good pest control. Small droplets are more prone to spray drift and can be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Do not apply at wind speeds greater than 10 mph at the application site.
- Apply as close to target plants as practical to obtain a good spray pattern for adequate coverage.
- For aerial applications, do not apply at heights greater than 10 feet (consistent with flight safety).
- For airblast applications, direct spray above foliage and turn off outward pointing nozzles at row ends and outer rows.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum boom length should be used and must not exceed 75% of wing span or rotor diameter.
- For ground applications, do not apply at heights greater than 4 feet.

BEANS AND LIMA BEANS
Dry and Succulent Forms

USE RESTRICTIONS:

- Do not feed treated vines to livestock.
- Do not apply more than 2 2/3 lbs./acre (2 lbs. AI) per season.
- Do not harvest snapbeans (succulent or dry beans) for at least 14 days after last application.
- Lima beans may be harvested 24 hours after the last application (1 day preharvest interval)

APPLICATION INSTRUCTIONS:

- Apply when eggs or insects first appear.
- Use the higher rates listed for severe infestations.

FOLIAR APPLICATIONS:

By Ground: Apply using 20 to 100 gals/A of spray.
By Air: Apply in a minimum of 2 gals/A of spray.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Fleahoppers Grasshoppers	1/3 to 2/3 lb.	For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary to maintain control.*
Aphids (excluding Black Bean Aphid) Bean Leaf Beetle Bean Leafroller Cabbage Looper Cutworms Green Cloverworms Leafhoppers Mexican Bean Beetle Plantbugs (Lygus) Soybean Looper Thrips Whitefly (except Sweet Potato or Silverleaf Whitefly)	2/3 to 1 1/3 lbs.	To maintain control: When applying 0.5 lbs. of active ingredient per acre, repeat at 3 day spray intervals as necessary.* When applying more than 0.5 lbs. of active ingredient per acre, repeat at 7 day spray intervals as necessary.*
Armyworms (excluding Beet Armyworm) Corn Earworm European Corn Borer	1 to 1 1/3 lbs.	
* Do not apply more than 2 2/3 lbs./acre (2 lbs. AI) per season.		

CELERY

USE RESTRICTIONS:

- Trim and remove all celery tops before shipment for use.
- Do not use trimmed tops for feed or food.
- Do not apply more than 2 2/3 lbs. per acre (2 lbs. active ingredient) per season.
- Do not harvest celery for at least 21 days after last application.

APPLICATION INSTRUCTIONS:

- Apply when eggs or insects first appear.
- Use the higher rates listed for severe infestations.

FOLIAR APPLICATIONS:

Ground: Apply using 50 to 100 gallons / acre of spray.

By Air: Apply in a minimum of 5 gallons / acre of spray.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Green Peach Aphid	2/3 to 1 1/3 lbs.	To maintain control: For applications of 0.5 lbs. active ingredient per acre, make repeat applications at 3 day intervals as necessary.*
Cabbage Looper Fall Armyworms	1 1/3 lbs.	For applications greater than 0.5 lbs. active ingredient per acre, make repeat applications at 7 day intervals as necessary.*
* Do not apply more than 2 2/3 lbs. per acre (2 lbs. active ingredient) per season.		

COLE CROPS
Brussels Sprouts & Cauliflower

USE RESTRICTIONS:

- Do not feed trimmings to livestock or allow animals to graze in treated areas.
- Do not apply more than 2 2/3 lbs./A (2 lbs. a.i.) per season.
- Do not harvest for at least 14 days after last application.

APPLICATION INSTRUCTIONS:

- Apply when eggs or insects first appear.
- Use the higher rates listed for severe infestations.

FOLIAR APPLICATIONS:

By Ground: Apply using 25 to 150 gallons/acre of spray.

By Air: Apply in a minimum of 5 gals/A of spray.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Green Peach Aphid	2/3 to 1 1/3 lbs.	To maintain control: For applications of 0.5 lbs. active ingredient per acre, make repeat applications at 3 day intervals as necessary.*
Cabbage Looper Diamondback Moth Larvae Imported Cabbage Worm	1 1/3 lbs.	For applications greater than 0.5 lbs. active ingredient per acre, make repeat applications at 7 day intervals as necessary.* NOTE: Diamondback Moth Larvae have been found to develop resistance to various insecticides. Consult your local Agricultural Extension Agent for the current recommended best practices for dealing with this insect.
*Do not apply more than 2 2/3 lbs./A (2 lbs. a.i.) per season.		

COTTON

USE RESTRICTIONS:

- Do not feed gin trash or treated forage to livestock.
- Do not allow animals to graze on treated areas.
- Do not harvest cotton for at least 21 days after last application.
- Do not apply more than 4 lbs. a.i. per season. This includes all uses of Acephate products and Acephate 75% SP on cotton.
- For aerial application in California and Arizona, application is limited to 1.0 lb. a.i./acre. For all other areas of the U.S., application is limited to 0.75 lb. a.i./acre by air.

APPLICATION INSTRUCTIONS:

- Use the higher rates listed for severe infestations.
- Apply when eggs or insects first appear.

IN-FURROW (AT-PLANTING) APPLICATION:

By Ground: Apply using 3 to 5 gallons of spray per acre.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Aphids (excluding Cotton Aphids in AZ and CA) Black (Greasy) Cutworm (Except CA) Thrips (including Western Flower Thrips)	2/3 to 1 1/3 lbs. NOTE: For the Blacklands Region of Texas, use the 1 1/3 lbs./acre application rate	Use only flat-fan nozzles. Verify that the nozzles are aligned to maximize spray deposition into the seed furrow. Securely fasten spray-system tubing to furrow opener and check frequently to ensure proper positioning and operation. Acephate 75% SP can be mixed with fungicides that are sprayed in-furrow for disease control.

FOLIAR APPLICATION:

By Ground: Apply using 10 to 25 gallons of spray per acre.

By Air: Apply using 3 to 10 gallons of spray per acre. In Arizona and California, apply in a minimum of 5 gallons of spray per acre.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Thrips (including Western Flower Thrips)	1/3 lb.	To maintain control, repeat applications at 3-day spray intervals as necessary.*
Plantbugs (Lygus)	1/3 to 1 1/3 lbs. (AZ & CA) 1 to 1 1/3 lbs.	Use the highest rate listed for Lygus adults that have migrated into cotton. To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
Fleahopper	1/3 lb.	To maintain control, repeat applications at 3 day intervals as necessary.*

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Cotton Aphids (excluding AZ and CA)	2/3 to 1 1/3 lbs.	Cotton Aphids have been found to develop resistance to various insecticides. Consult your local Agricultural Extension Agent for the current recommended best practices for dealing with this insect. To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
Whitefly (excluding Sweetpotato Whitefly / Silverleaf Whitefly)	2/3 to 1 1/3 lbs.	To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
Armyworms (excluding Beet Armyworm) Cabbage Looper	1 1/3 lbs.	Apply when eggs appear and repeat applications using 7-day spray intervals.*
Bollworm Tobacco Budworm (Adults, Larvae) Eggs/ DEADHATCH®	2/3 to 1 1/3 lbs. (East of Rockies) 1 1/3 lbs. (AZ & CA)	Moths of budworm larvae are controlled by direct contact with spray. Death is most likely to occur when late evening applications are made. Use 2/3 lbs./acre for mild infestations. Use 1 to 1 1/3 lbs./acre for moderate to severe infestations. To maintain control: For applications of 0.5 lbs. active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.* DEADHATCH®: Control of emerging larvae by consumption of treated egg casings.
Stinkbugs	1.0 lb.	Apply when eggs appear and repeat applications using 7-day spray intervals.*
Pink Bollworm (AZ & CA)	1 1/3 lbs.	Apply when insects appear and repeat applications using 7-day spray intervals.*
Cutworms	1.0 lb.	Apply by ground application; aerial applications are less effective, but may be also used. For best results when making ground applications: <ul style="list-style-type: none"> • Apply in the evening • Direct spray to the lower portion and base of plant. Begin applications when the insects first appear or damage is first noted. To maintain control, repeat applications using 7-day spray intervals as necessary.*
*Do not apply more than 4 lbs. a.i. per season. This includes all uses of Acephate products and Acephate 75% SP on cotton.		

COTTON TANK MIXES

USE RESTRICTIONS:

- Do not feed gin trash or treated forage to livestock.
- Do not allow animals to graze on treated areas.
- Do not harvest cotton for at least 21 days after last application.
- Do not apply more than 5 1/3 pounds per acre (4 lbs. of active ingredient) per season. This includes all uses of Acephate 75% SP on cotton.
- For aerial application in California and Arizona, application is limited to 1.0 lbs. of active ingredient per acre. For all other areas of the U.S., application is limited to 0.75 lbs. of active ingredient per acre by air.

COTTON PEST CONTROL RESISTANCE MANAGEMENT

Consult your local agricultural extension agent for current best practices when controlling cotton pests, and in particular, when dealing with silverleaf whitefly control.

APPLICATION INSTRUCTIONS:

- Always read all label directions when using any pesticide alone or in tank mix combinations. Follow the most restrictive labeling when using a tank mix.
- Use the higher rates listed for severe infestations.
- Apply when insects first appear or when damage is first noted.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 25 gallons per acre of spray.

By Air: Apply using 3 to 10 gallons per acre of spray (minimum 5 gallons per acre in CA).

TANK MIXING:

- This product may be tank mixed with registered cotton herbicides under the following conditions:
 - The tank mix partners do not prohibit tank mixes;
 - The most restrictive of label limitations and precautions are observed; and,
 - No label dosage rates are exceeded.
- To determine physical compatibility of potential tank-mix partners with Acephate 75% SP:
 1. In a pesticide-safe container, pour the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank. NOTE: When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides.
 2. Mix thoroughly and allow to stand for five minutes.
 3. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.

PESTS CONTROLLED	ACEPHATE 75% SP AND TANK MIX PARTNER	ACEPHATE 75% SP + TANK MIX PARTNER (PER ACRE)	SPECIFIC INSTRUCTIONS
Sweetpotato Whitefly (Silverleaf Whitefly)	Acephate 75% SP + Danitol® 2.4 EC Spray	2/3 lb. + 10 2/3 to 16 fl. oz.	To maintain control, repeat applications using 3-day intervals as necessary.*

PESTS CONTROLLED	ACEPHATE 75% SP AND TANK MIX PARTNER	ACEPHATE 75% SP + TANK MIX PARTNER (PER ACRE)	SPECIFIC INSTRUCTIONS
Armyworms (excluding Beet Armyworm) Bollworm Cabbage Looper Cotton Aphid Cutworms Fleahopper Grasshoppers Pink Bollworm Plantbugs (Lygus, Mirids) Salt Marsh Caterpillar Thrips (including Western Flower Thrips) Tobacco Budworm Whitefly (excluding Sweetpotato/Silverleaf Whitefly)	Acephate 75% SP + Lorsban® 4E	1/3 to 1 1/3 lbs. + 1 to 2 pts.	To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.* Special Instructions for Cutworms: Use 1.0 lb. / acre of Acephate 75% SP Apply by ground application; aerial applications are less effective, but may be also used. For best results when making ground applications: <ul style="list-style-type: none"> • Apply in the evening • Direct spray to the lower portion and base of plant.
* Do not apply more than 5 1/3 pounds per acre (4 lbs. of active ingredient) per season. This includes all uses of Acephate 75% SP on cotton.			

COTTON TANK MIXES WITH PYRETHROIDS

USE RESTRICTIONS:

- Do not feed gin trash or treated forage to livestock.
- Do not allow animals to graze on treated areas.
- Do not harvest cotton for at least 21 days after last application EXCEPT when applying with Scout X-TRA. When tank mixing with Scout X-TRA, do not harvest cotton for at least 28 days after last application.
- Do not apply more than 5 1/3 lbs./A (4 lbs. a.i.) per season. This includes all uses of Acephate 75% SP on cotton.
- For aerial application in California and Arizona, application is limited to 1.0 lbs. of active ingredient per acre. For all other areas of the U.S., application is limited to 0.75 lbs. of active ingredient per acre by air.

COTTON PEST CONTROL RESISTANCE MANAGEMENT

Consult your local agricultural extension agent for current best practices when controlling cotton pests, and in particular when dealing with silverleaf whitefly control.

APPLICATION INSTRUCTIONS:

- Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions that appear on all product labels. The most restrictive labeling applies when using a tank mix.
- Use the higher rates listed for severe infestations.
- Apply when insects first appear or when damage is first noted.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 25 gallons per acre of spray.

By Air: Apply using 3 to 10 gallons per acre of spray (minimum 5 gallons per acre in CA).

TANK MIXING:

- This product may be tank mixed with registered cotton herbicides under the following conditions:
 - The tank mix partners do not prohibit tank mixes;
 - The most restrictive of label limitations and precautions are observed; and,
 - No label dosage rates are exceeded.
- To determine physical compatibility of potential tank-mix partners with Acephate 75% SP:
 1. In a pesticide-safe container, pour the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank. NOTE: When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides.
 2. Mix thoroughly and allow to stand for five minutes.
 3. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.

PESTS CONTROLLED	ACEPHATE 75% SP AND TANK MIX PARTNER	ACEPHATE 75% SP + TANK MIX PARTNER (PER ACRE)	SPECIFIC INSTRUCTIONS
Aphids Bollworm Cabbage Looper Cotton Leaf Perforator Cutworms Fall Armyworms Fleahoppers Pink Bollworm (AZ & CA) Plantbugs Stinkbugs Sweetpotato/Silverleaf Whitefly Thrips (including Western Flower Thrips) Tobacco Budworm Whitefly	Acephate 75% SP + one of the following:	2/3 to 1 1/3 lbs. + one of the following:	To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.* For Stinkbugs and Cutworms: Use 1.0 lb. / acre of Acephate 75% SP Special Instructions for Cutworms: Apply by ground application; aerial applications are less effective, but may be also used. For best results when making ground applications: <ul style="list-style-type: none"> • Apply in the evening • Direct spray to the lower portion and base of plant.
	Ammo® 2.5 EC*	Refer to the Ammo® 2.5 EC approved label for use instructions.	
	Asana® XL*	Refer to the Asana® XL approved label for use instructions.	
	Baythroid® 2 EC	Refer to the Baythroid® 2 EC approved label for use instructions.	
	Capture® 2 EC	Refer to the Capture® 2 EC approved label for use instructions.	
	Karate® 1E (except CA)	Refer to the Karate® 1E approved label for use instructions.	
	Scout X-TRA® (except CA)	Refer to the Scout X-TRA® approved label for use instructions.	
*Do not apply more than 5 1/3 pounds per acre (4 lbs. of active ingredient) per season. This includes all uses of Acephate 75% SP on cotton.			

CRANBERRIES

USE RESTRICTIONS:

- Limit to one application per growing season.
- Do not apply more than 1 1/3 lbs. / acre (1 lb. a.i.) per season.
- Do not apply from start of bloom until all berries set.
- Do not harvest cranberries for at least 90 days after last application.
- Do not apply when wind speed may cause spray drift beyond the desired treatment area.

FOLIAR APPLICATIONS:

By Ground or Sprinkler: Use sufficient water to give thorough coverage.
 By Air: Minimum of 2 gallons per acre of spray

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)
Cranberry Blossom Worm Cranberry Cutworm Gypsy Moth False Armyworm Fireworms Spanworms Sparganothis	1 1/3 lbs.

USE PRECAUTIONS FOR SPRINKLER IRRIGATION APPLICATION TO CRANBERRIES:

Acephate 75% SP may only be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Solid Set System: Apply specified dosage for the entire length of the irrigation period or for a 30 to 60 minute period at the end of a regular irrigation set or as a 30 to 60 minute injection as a separate application not associated with a regular irrigation. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining pesticide, a dye indicator may be injected into the lines to mark the end of the application period. See **NOTE**.

Center Pivot Systems: Inject the specified dosage per acre continuously for one complete revolution of the system. See **NOTE**.

NOTE: Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Inject the product with a positive displacement pump into the main line ahead of a right turn to ensure adequate mixing.

Application of more than label-specified quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.

HEAD LETTUCE – Crisphead Type Only

USE RESTRICTIONS:

- Do not feed trimmings to livestock or allow animals to graze on treated areas.
- AZ & CA, Desert Areas: Do not apply after first head begins to form in crops that germinate from mid-September through November.
- Do not apply more than 2 2/3 lbs./A (2 lbs. a.i.) per season.
- Do not harvest lettuce for at least 21 days after last application.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 60 gallons per acre of spray.

By Air: Apply using a minimum of 5 gallons per acre of spray

PESTS CONTROLLED	ACEPHATE 75%-SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Aster Leafhopper Green Peach Aphid	2/3 to 1 1/3 lbs.	To maintain control:
Armyworms (excluding Beet Armyworm) Cabbage Looper	1 1/3 lbs.	For applications of 0.5 lbs. active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
*Do not apply more than 2 2/3 lbs./A (2 lbs. a.i.) per season.		

MINT
(PEPPERMINT AND SPEARMINT)

USE RESTRICTIONS:

- Do not graze treated areas.
- Do not apply more than 2 2/3 lbs./A (2 lbs. a.i.) per season.
- Do not harvest mint for at least 14 days after last application.

APPLICATION METHOD	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
<p style="text-align: center;">FOLIAR</p> <p>By Ground: Apply using 20 to 100 gallons per acre</p> <p>By Air: Apply using 5 to 10 gallons per acre</p>	<p style="text-align: center;">Alfalfa Looper Aphids Cutworms</p>	<p style="text-align: center;">1 1/3 lbs. (21.25 oz.)</p>	<p>Apply when eggs or insects first appear.</p> <p>To maintain control: One repeat application can be made. Allow 7 days between applications.</p>
<p style="text-align: center;">FOLIAR</p> <p>By Ground: 40 to 100 gallons per acre</p>	<p style="text-align: center;">Strawberry Root Weevil (Adult) Black Vine Weevil (Adult)</p>	<p style="text-align: center;">1 1/3 lbs. (21.25 oz.)</p>	<p>For control, good coverage and canopy penetration is essential. In areas of tall or dense canopies, increase spray pressure and / or volume to ensure adequate canopy penetration.</p> <p>For best results: Apply after adult emergence is complete but prior to egg laying. Apply at dusk or during the night on a warm still evening. Two applications 7 days apart may be necessary to reduce heavy infestations.</p>

PEANUTS

USE RESTRICTIONS:

- Do not feed treated forage or hay to livestock.
- Do not allow animals to graze on treated areas.
- Do not apply more than 5 1/3 lbs. per acre (4 lbs. a.i.) per crop cycle.
- Do not harvest (dig) peanuts for 14 days after last application.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 50 gallons per acre.

By Air: Apply using 5 to 10 gallons per acre.

TANK MIXING:

- This product may be tank mixed with registered at-cracking and early postemergence peanut herbicide under the following conditions:
 - The tank mix partners do not prohibit tank mixes;
 - The most restrictive of label limitations and precautions are observed; and,
 - No label dosage rates are exceeded.
- To determine physical compatibility of potential tank-mix partners with Acephate 75% SP:
 1. In a pesticide-safe container, pour the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank. NOTE: When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides.
 2. Mix thoroughly and allow to stand for five minutes.
 3. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.
- Keep combinations agitated and apply immediately. Do not allow combinations to set in the supply tank for prolonged periods of time.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Grasshoppers	1/3 to 2/3 lb.	Apply when eggs or insects first appear. To maintain control:
Thrips	1/2 to 1.0 lb.	
Corn Earworm Fall Armyworm Leafhoppers Loopers Velvetbean Caterpillar	1 to 1 1/3 lbs.	For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
*Do not apply more than 5 1/3 lbs. per acre (4 lbs. a.i.) per crop cycle.		

PEPPERS, BELL

USE RESTRICTIONS:

- Do not apply more than 2 2/3 lbs. per acre (2 lbs. of active ingredient) per season.
- Do not harvest peppers for at least 7 days after last application.

APPLICATION INSTRUCTIONS:

- Apply when eggs or insects first appear.

FOLIAR APPLICATIONS:

By Ground: Apply using 25 to 150 gallons per acre of spray.

By Air: Apply using a minimum of 3 gallons per acre of spray (except in CA where a minimum of 5 gallons per acre is required)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Grasshoppers	1/3 to 2/3 lb.	To maintain control: For applications of 0.5 lbs. or less active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
Cabbage Looper Green Peach Aphid Tobacco Hornworm	2/3 to 1 1/3 lbs.	
European Corn Borer	1 to 1 1/3 lbs.	
*Do not apply more than 2 2/3 lbs. per acre (2 lbs. of active ingredient.) per season.		

PEPPERS, NON-BELL

(FOR USE IN MIDWESTERN STATES, EASTERN STATES AND PUERTO RICO ONLY)

USE RESTRICTIONS:

- Do not apply more than 1 1/3 lbs. per acre (1 lb. of active ingredient) per season.
- Do not harvest peppers for at least 7 days after last application.

FOLIAR APPLICATIONS:

By Ground: Apply using 40 to 150 gallons per acre of spray.

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Aphids	2/3 lb.	To maintain control, make repeat applications using 3-day spray intervals.*
*Do not apply more than 1 1/3 lbs. per acre (1 lb. of active ingredient.) per season.		

TOBACCO

(Use on Tobacco is not registered for use in California)

USE RESTRICTIONS:

- Do not apply more than 1.0 lb. Acephate 75% SP per acre as a transplant-water application as phytotoxicity may occur.
- Do not apply more than 5 1/3 lbs of Acephate 75% SP per acre (4 lbs. active ingredient) per season. This includes the use of Acephate 75% SP in transplant water, plant bed, soil, float bed, (greenhouse), and foliar applications.
- Do not harvest tobacco for at least 3 days after last application.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 50 gallons per acre of spray.

By Air: Apply using a minimum of 3 gallons per acre of spray

APPLICATION METHOD	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
FLOATBED	Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	1.0 lb.	Apply evenly to ensure thorough coverage. Allow 7 days between applications.* NOTE: Floatbed water should be disposed of in the transplanted field as transplant water or foliar spray.
PLANT BED	Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	1.0 lb.	Apply evenly to ensure thorough coverage. Allow 7 days between applications.*
TRANSPLANT Apply in a minimum of 100 gallons of transplant water per acre.	Cutworms Flea Beetle Green Peach Aphid Potato Tuberworm Tobacco Aphid Tobacco Thrips	1.0 lb.	Provides control for approximately 3 to 4 weeks after transplanting. For late season control, use foliar sprays. Make transplant water applications using mechanical transplant equipment only. Using such equipment, the insecticide/water mixture is mechanically applied directly into the soil along with the transplanted plants.

APPLICATION METHOD	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
<p>FOLIAR By Ground: Apply using 10 to 50 gallons per acre</p> <p>By Air: Apply using a Minimum of 3 gallons per acre</p>	Grasshoppers	1/3 to 2/3 lb.	To maintain control: Repeat applications at 3-day spray intervals as necessary.*
	Flea Beetle Green Peach Aphid Hornworm Tobacco Thrips	2/3 lb.	To maintain control: Repeat applications at 3-day spray intervals as necessary.*
	Stinkbugs Tobacco Aphid Vegetable Weevils	2/3 to 1 lb.	To maintain control: For applications of 0.5 lbs. active ingredient per acre, repeat applications at 3-day spray intervals as necessary.* For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.*
	Budworm Cabbage Looper Cutworm Japanese Beetle	1.0 lb.	To maintain control: Repeat applications at 7-day spray intervals as necessary.*
<p>SOIL MOUND TREATMENT</p> <p>DRENCH METHOD</p>	<p>Ants (including Imported Fire Ants and Harvester Ants)</p> <p>(excluding Carpenter and Pharaoh ants)</p>	1/3 lb. per 26.6 gallons water	<p>Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet.</p> <p>Treat all areas within four feet of the mound.</p> <p>Restrictions: Treat a maximum of 13 mounds per acre. Do not treat more than once per season.</p> <p>For best results: Apply in the early morning or late afternoon, when ants are most active.</p> <p>Application under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest.</p>
<p>*Do not apply more than 5 1/3 lbs of Acephate 75% SP per acre (4 lbs. active ingredient) per season. This includes the use of Acephate 75% SP in transplant water, plant bed, soil, float bed, (greenhouse), and foliar applications.</p>			

NON-BEARING CITRUS

USE RESTRICTIONS

- Do not graze treated areas.
- Do not harvest citrus for one year after treatment.

APPLICATION INSTRUCTIONS

- Spray individual juvenile or non-bearing trees for coverage with total application not to exceed specified rate in lbs. per acre.
- The length of residual activity will depend upon spray coverage and the amount of precipitation following the application.

APPLICATION METHOD	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
FOLIAR By Ground: Apply using 100 to 200 gallons/acre of spray	Aphids Grasshoppers Katydid Mealybugs Orangedogs Plantbugs Thrips Whiteflies (except Sweetpotato/Silverleaf)	2/3 lb.	To maintain control: Reapply using 3-day spray intervals as necessary.
	Citrus Blackfly (Use on Citrus Blackfly is not registered for use in California)	2/3 to 1.0 lb.	Apply when eggs or insects first appear. To maintain control: For applications of 0.5 lbs. active ingredient per acre, repeat applications at 3-day spray intervals as necessary. For applications greater than 0.5 lbs. active ingredient per acre, repeat applications at 7-day spray intervals as necessary.
	Ants (excluding fire, harvester, carpenter and pharaoh ants)	1.0 lb.	Repeat application at 7-day spray intervals as necessary to maintain control.
SOIL MOUND TREATMENT DRENCH METHOD	Ants (including Imported Fire Ants and Harvester Ants) (excluding Carpenter and Pharaoh ants) Not registered for use on harvester ants in California.	1/3 lb. per 26.6 gallons water	Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet. Treat all areas within four feet of the mound. Repeat applications may be made as necessary to maintain control.

NON-CROP AREAS

FIELD BORDERS, FENCEROWS, ROADSIDES, DITCHBANKS, BORROW PITS**USE RESTRICTIONS:**

- Do not graze or feed vegetation cut from treated areas.

APPLICATION INSTRUCTIONS

- Make applications only early to mid-season.
- Use the higher application rates when applying to dense foliage.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 20 gallons per acre.

By Air: Apply using 1 to 5 gallon per acre (except in CA where a minimum of 5 gallons per acre must be used)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Grasshoppers	1/3 lb.	Use of an approved drift retardant to assist in controlling drift and reducing evaporation is acceptable.

WASTELAND (NON-CROP AREAS)**USE RESTRICTIONS:**

- Do not make more than one application per season.
- Do not graze or feed vegetation cut from treated areas.
- Do not allow animals to graze on treated areas.

APPLICATION INSTRUCTIONS

- Use the higher application rates when applying to dense foliage.

FOLIAR APPLICATIONS:

By Ground: Apply using 10 to 20 gallons per acre.

By Air: Apply using ½ gallon per acre (except in CA where a minimum of 5 gallons per acre must be used)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Black Grass Bugs Grasshoppers Mormon Crickets	1/8 to 1/6 lb. (2.0 to 2.7 oz.)	If necessary/desired, add an approved drift retardant to assist in controlling drift and reducing evaporation.

SPECIALTY USES – NON CROP AREAS, EXCLUDING RESIDENTIAL TURF

USE RESTRICTIONS:

- Do not treat mound more than once per season.

NOTE: Grass in treated area may be injured.

PEST CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Fire ants and harvester ants (Not registered for use on harvester ants in California)	Drench Method: Mix 1/3 lb. in 26.6 gallons of water.	Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet. Treat all areas within four feet of the mound. <i>Restrictions: Treat a maximum of 13 mounds per acre. Do not treat more than once per season.</i> For best results: Apply in the early morning or late afternoon, when ants are most active. Application under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest.

SPECIALTY USES – CONTAINER GROWN ORNAMENTAL NURSERY STOCK

USE RESTRICTIONS:

- When used on greenhouse and outdoor ornamentals, do not apply more than 1.0 pound of active ingredient / acre per 100 gallons of water.
- Do not exceed 0.75 pounds of active ingredient / acre for cut flowers and 1.0 pound of active ingredient / acre for all other ornamentals.
- Allow a minimum of 7 days between applications.

APPLICATION INSTRUCTIONS:

- Apply the specified amount of Acephate 75% SP per 100 gals. of solution so as to thoroughly drench the root system.

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	APPLICATION TIMING
Container Grown Ornamental Nursery Stock (Arborvitae, Azalea, Camellia, Rhododendron, Roses, Viburnum, Yew)	Black Vine Weevil Strawberry Root Weevil	1.0 lb. (16 oz.)	Application should be made by mid-September for greenhouse stock and by mid-October for outdoor stock. Consult your local county extension agent for information on the identification and control of root weevils on ornamentals.
	Ants (excluding fire, harvester, carpenter and pharaoh ants)		Apply as needed for control.

ORNAMENTAL TREES AND SHRUBS

USE RESTRICTIONS:

- Do not exceed 0.75 pounds of active ingredient / acre for cut flowers and 1.0 pound of active ingredient / acre for all other ornamentals.

APPLICATION INSTRUCTIONS:

- When used on nursery crops, spray several sample plants and observe for signs of phytotoxicity for a minimum of two weeks before applying to all plants.
- Unless otherwise noted, apply the specified amount of Acephate 75% SP per 100 gallons of water using a hydraulic sprayer.
- Avoid applying to Huckleberry, Balm of Gilead, Cottonwood, Lombardy Poplar and Viburnum suspensum as phytotoxicity may be observed.

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 GALLONS)	APPLICATION TIMING	SPECIFIC INSTRUCTIONS
Ornamental Trees and Shrubs (Except Flowering Crabapple, see below)	Aphids Bagworms Birch Leafminer Tent Caterpillars* Lace Bugs Leafrollers	1/3 lb. (5.3 oz.)	Upon appearance of the insects.	To maintain control: Repeat applications at 14-day spray intervals as necessary. Use of a compatible sticker will improve control of Gypsy Moth larvae. Mist blower application: Adjust rates to 1 1/3 lbs. per 100 gals. water for Gypsy Moth control and 1.0 lb. per 100 gals. water for Tent Caterpillar control.
	Douglas Fir Tussock Moth Larvae Gypsy Moth Larvae* Webworms	2/3 lb. (10.5 oz.)	Upon appearance of the insects.	
	Scales (Crawlers)	2/3 lb. (10.5 oz.)	As crawlers begin to appear.	
	Ponderosa Pine Needle Miner	2/3 lb. (10.5 oz.)	Consult your local County Extension Agent for optimal application timing.	
	Grasshoppers	2/3 lb. (10.5 oz.)	Upon appearance of grasshoppers.	
	California Oakworm Cankerworms (Spring and Fall)	1/3 to 2/3 lb. (5.3 to 10.5 oz.)	Upon appearance of the insects. Use the higher amounts listed when larger larvae are present.	
	Nantucket Pine Tip Moth Larvae	1.0 lb. (16 oz.)	Consult your local County Extension Agent for optimal application timing.	
	Root Weevil Adults	1.0 lb. (16 oz.)	Apply as soon as damage is observed.	
	Box Elder Bugs Sawflies Budworms Leafhoppers	1.0 lb. (16 oz.)	Upon appearance of the insects.	

320948

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 GALLONS)	APPLICATION TIMING	SPECIFIC INSTRUCTIONS
	Japanese Beetle	1 1/3 lbs. (21.3 oz.)	Apply as soon as insects are observed.	
	Elm Leaf Beetle (larvae)	1 1/3 lbs. (21.3 oz.)	As the larvae begin to appear. Acephate 75% SP will not prevent Elm Leaf Beetle eggs from hatching.	
Douglas Fir (Christmas Trees)	Douglas Fir Needle Midge	2/3 lb. (10.5 oz.) (2/3 lb. per acre)	Application should be made within two weeks of bud burst. For current best practices information, consult your County Extension Service.	Apply the specified amount of Acephate 75% SP in not less than 2 gals. of spray per acre by air or in 100 gals. of spray per acre by ground.
Flowering Crabapples	Aphids Tent Caterpillars Leafrollers	1/3 lb. (5.3 oz.)	Upon appearance of the insects.	Do not apply more often than 3 times in a growing season (using a minimum of 4-week intervals). Phytotoxicity has been documented on the following crabapple varieties: Hopa, Ichonoski, Malusfloribunda, Pink Perfection, Red Wine and Snow Cloud.

GOLF COURSE AND SOD FARM TURFGRASS

(Use on Golf Course and Sod Farm Turfgrass is not registered for use in California)

USE RESTRICTIONS:

- Do not allow livestock to graze in treated areas.
- Do not feed grass clippings to livestock.
- Use is limited to sod farms and golf courses, except when applying by mound or spot treatment for fire ant and harvester ant control (excluding residential turf).
- Aerial application to turf is prohibited.
- Wait at least 3 days following the last Acephate 75% SP application prior to harvesting sod.
- *The maximum application rate is 3 lbs. a.i./acre for sod farms and 4 lbs. a.i./acre for golf courses.*

APPLICATION INSTRUCTIONS:

- Apply as soon as insects or damage caused by the insects first appears.
- To maintain control, repeat applications using a minimum of one-week intervals as necessary.

Turfgrass (Golf Courses and Sod Farms only)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 1000 SQ. FT.)	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Fall Armyworm Yellow Striped Armyworm Southern Armyworm	½ to 1.2 oz. (1 1/3 to 3 ¼ lbs. / acre)	Use a minimum of 5 gals. of water per 1,000 sq. ft. as necessary to obtain good coverage.	To maintain control, repeat applications at 7-day intervals may be necessary.
Cutworm	1.2 to 1.9 oz. (3 ¼ to 5.2 lbs. / acre)		
Chinch Bugs	1.2 to 1.9 oz. (3 ¼ to 5.2 lbs. / acre)	Use 1 to 15 gals. of water per 1,000 sq. ft. as necessary to obtain good coverage.	Use the higher amount when quick knockdown is needed or with heavy infestations. To maintain control, repeat applications at 7-day intervals may be necessary.
Sod Webworm (Crambus spp.)	½ to 1.0 oz. (1 1/3 to 2 ¾ lbs. / acre)		
Leafhopper	(2 oz. 2 ¾ lbs. / acre)		

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 1000 SQ. FT.)	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Mole Crickets (except CA) Spittlebug (except CA)	1.0 to 1.9 oz. (2.66 to 5.2 lbs. / acre)	Use 1 to 15 gals. of water per 1,000 sq. ft. as necessary to obtain good coverage.	For best results, apply after irrigating in the late afternoon or early evening. Do not irrigate after application. A lemon fragrance substance may be added to the spray mix as a flushing agent, potentially providing increased mole cricket contact with Acephate 75% SP. The following products have been shown to be effective flushing agents: Lemon Palmolive, Lemon Joy, and Mighty Myrt Products Manufacture-Base Pure Lemon Fragrance. Use 2-teaspoons per gallon of water for small mix volumes or 6 fl. oz. per 50 gallons of water for large mix volumes. More than one application may be required for knockdown of existing populations. Allow at least 7 days between applications. For heavy infestations, use the higher dosage rate.
Greenbug (<i>Schizaphis graminum</i>) Grasshoppers	½ oz. (1 1/3 lbs. / acre)	Use 4 gals. of water per 1,000 sq. ft.	Do not mow urfgrass for at least 24 hours after application.
Black Turfgrass Ataenius (except CA)	1.5 to 1.9 oz. (4.1 to 5.2 lbs. / acre)	Use a minimum of 5 gallons water per 1,000 sq. ft.	Irrigate lightly after application (not more than ½ inch). Use the higher rate for severe infestations.

Dichondra
(Plant Site for Golf Course and Sod Farm Turfgrass only)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 1000 SQ. FT.)	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Cutworm Flea Beetle Southern Armyworm Yellow Striped Armyworm	1.0 to 1.9 oz. (2.7 to 5.2 lbs. / acre)	Use a minimum of 15 gals. of water per 1,000 sq. ft. to obtain good coverage.	

MOUND TREATMENT OF FIRE ANTS AND HARVESTER ANTS IN TURFGRASS, EXCLUDING RESIDENTIAL TURF

USE RESTRICTIONS:

- Not registered for use on harvester ants in California.
- Do not treat mound more than once per season.

NOTE: Grass in treated areas may be injured. Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest.

ACEPHATE 75% SP APPLICATION RATE	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Mound Treatment Drench Method: Mix 1/3 lb. in 26.6 gals. of water.	For best results, apply when the ants are most active (in the early morning or late afternoon).	Apply 1 gallon of mix to each mound area by sprinkling the mound until it is wet. Treat all areas within four feet of the mound.

OUTDOOR AND PERIMETER SPRAY

USE RESTRICTIONS:

- Not for use on residential turfgrass.

APPLICATION INSTRUCTIONS:

- Apply as soon as insects appear.

PEST CONTROLLED	ACEPHATE 75% SP APPLICATION RATE	SPECIAL INSTRUCTIONS	APPLICATION INSTRUCTIONS
Wasps	1/3 lb. in 3.3 gals. of water	Treat when wasps are less active (early or late in the day).	Apply as a spot treatment to the nest, nest entrance, and surrounding areas where the wasps may land.
Ants (excluding fire, harvester, carpenter and pharaoh ants) Crickets Cockroaches Earwigs Pillbugs	1/3 lb. in 3.3 gals. of water		Apply to the ground within 6 to 10 feet of the structure and to a height of 2 to 3 feet on the foundation. Apply using spray or with a paint brush to surfaces of buildings, window frames, shutters, entryways, screens, eaves, patios, garages, carports, around garage areas and other areas where the insects may occur.

OUTDOOR FLORAL CROPS AND GROUND COVERS

USE RESTRICTIONS:

- When used on greenhouse and outdoor ornamentals, application is limited to 1.0 lb. of active ingredient per 100 gallons of water.
- Do not exceed 0.75 pounds of active ingredient / acre for cut flowers and 1.0 pound of active ingredient / acre for all other ornamentals.

APPLICATION INSTRUCTIONS:

- Apply as insects begin to appear.
- Apply as a full coverage spray using a hydraulic sprayer.
- Multiple applications may cause slight tip burn or marginal leaf necrosis on some varieties. A test should be made on a sample plant to determine if unacceptable damage occurs prior to application to all plants.

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 GAL.)	APPLICATION TIMING
Chrysanthemums Daisies Dahlias Easter Lily Gladioli Gypsophila Pachysandra Pansy Peony Roses Sedum Statice Strawflower Yarrow Zinnia	Aphids Thrips Lygus	2/3 lb. (10.5 oz.)	To maintain control repeat applications may be necessary. Allow a minimum of 3 days between applications.
Roses Boston Ivy	Japanese Beetle	1 1/3 lbs. (21.3 oz.)	To maintain control, repeat applications may be necessary. Allow a minimum of 7 days between applications.

GREENHOUSE FLORAL AND FOLIAGE PLANT CROPS

(Use on Greenhouse Floral and Foliage Plant Crops is not registered for use in California)

USE RESTRICTIONS:

- When used on greenhouse and outdoor ornamentals, application is limited to 1.0 lb. of active ingredient per 100 gallons of water.
- Do not exceed 0.75 pounds of active ingredient / acre for cut flowers and 1.0 pound of active ingredient / acre for all other ornamentals.

APPLICATION INSTRUCTIONS:

- Apply as insects begin to appear.
- Unless otherwise instructed, apply as a full coverage spray using a hydraulic sprayer.
- Multiple applications may cause slight tip burn or marginal leaf necrosis on some varieties. A test should be made on a sample plant to determine if unacceptable damage occurs prior to application to all plants.
- Phytotoxicity has occurred on the following foliage plants:

Bletchum gibbum
 Cissus Antarctica
 Ficus triangularis
 Fittonia verschaffelti
 Maranta leuconeura kerchoveana
 Pachystachya lutea
 Plectranthus australis
 Polypodium aureus
 Polystichum
 Pteris ensiformis
 Tolmiea menziesii

Before treating large plantings, spray a test sample of plants and observe for 2-weeks for varietal phytotoxicity.

- Acephate 75% SP applications on certain varieties of Poinsettias after bract formation may result in phytotoxicity.
- Phytotoxicity has occurred on the following Chrysanthemum varieties:

Albatross
 Bonnie Jean
 Dixie
 Garland
 Gem
 Iceberg
 Pride
 Showoff
 Statesman
 Tally Ho
 Westward Ho
 Wild Honey

Before treating large Chrysanthemum plantings, spray a test sample of plants and observe for 2-weeks for varietal phytotoxicity.

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 GAL.)	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Roses	Leafrollers	2/3 to 1.0 lb. (10.5 to 16 oz.)	Apply with a hydraulic sprayer as a full coverage spray. To maintain control: Repeat applications at a 5-day minimum spray interval.	Use the higher amount listed when large larvae are present.
Foliage Plants Anthuriums Cacti Orchids Poinsettia	Aphids	1/3 lb. (5.3 oz.)	Apply with a hydraulic sprayer as a full coverage spray. Allow a minimum of 5 days between applications.	A wetting agent may be necessary when applying to difficult to wet foliage.
	Mealybugs Thrips Whiteflies	2/3 lb. (10.5 oz.)	To maintain control of mealybugs and whiteflies, repeat applications at a minimum of 5-day spray intervals.	
	Scales (Crawlers)	2/3 lb. (10.5 oz.)	To maintain control when there is continuous crawler production, repeat applications at a minimum of 5-day spray intervals.	
	Sweet Potato Whiteflies (except CA)	1/3 lb. (5.3 oz.) + Tame® 2.4 EC Spray 10 2/3 fl. oz. (0.2 lb. a.i.)	Allow a minimum of 5 days between applications. If a population is well established, make one application of the tank mix and follow in 5 to 7 days with Tame® alone at 16 fl. oz./100 gals. (See Tame® label for instructions).	

PLANTS	PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 GAL.)	APPLICATION INSTRUCTIONS	SPECIFIC INSTRUCTIONS
Roses Carnations Chrysanthemums	Aphids Thrips	2/3 lb. (10.5 oz.)	Apply with a hydraulic sprayer as a full coverage spray. Apply as insects appear or at the tight flower bud stage. To maintain control, repeat applications may be necessary using the following restrictions: A minimum of 5 days between applications to roses. A minimum of 28 days between applications to Carnations and Chrysanthemums.	Application to Chrysanthemums and Roses with open flowers will cause injury to the flowers.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep pesticide in original container. Store in a cool, dry place away from excessive heat. Do not put concentrate or dilute into food or drink containers.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill the outer bag. Dispose of empty outer foil bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the extent consistent with applicable law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

Ammo[®] and Capture[®] are registered trademarks of the FMC Corporation.
Asana[®] is a registered trademark of E.I. du Pont de Nemours and Company.
Baythroid[®] is a registered trademark of Bayer CropScience.
Danitol[®] is a registered trademark of Sumitomo Chemical Company, Ltd.
Karate[®] is a registered trademark of Syngenta Crop Protection, Inc.
Lorsban[®] is a registered trademark of Dow AgroSciences LLC.
Scout X-TRA[®] is a registered trademark of Hoechst Schering Agrovo S.A.
DEADHATCH[®] and Tame[®] are registered trademarks of Valent U.S.A. Corporation.

[Label for Non-WSP Product]
(product not packaged in water soluble packaging)

Acephate 75% SP

INSECTICIDE

ACTIVE INGREDIENT:	By Wt.
Acephate (<i>O,S</i> -Dimethyl acetylphosphoramidothioate).....	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
Acephate is an organophosphate, cholinesterase inhibitor	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-949-679-3535 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Acephate is a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure but decisions regarding treatment will usually need to be made before test results are available. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PROTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone.	

EPA Reg. No. 81964-2
EPA Est. No.

Manufactured for:
ChemStarr LLC
21 Hubble
Irvine, CA 92618

Net Weight:

430f48

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove and wash contaminated clothing before reuse. Do not allow children or pets to come into contact with treated areas until dusts have settled.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils and natural rubber ≥ 14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical-resistant gloves for all mixers and loaders

In addition, mixers and loaders supporting hopper box seed treatments must wear:

- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

This product and its degradate are highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

440948

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made out of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until dust has settled.

PRODUCT INFORMATION

Acephate 75% SP is a soluble powder used as an insecticide for control of pests on selected agricultural crops and in certain non-crop areas. The active ingredient in Acephate 75% SP is absorbed by plants via the roots and foliage to provide systemic control of feeding insects.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

FAILURE TO FOLLOW THE INSTRUCTIONS IN THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

COTTON

COTTON SEED HOPPER BOX TREATMENT – (Except AZ, CA and the Black Lands of TX)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Black (Greasy) Cutworm Cotton Aphids Thrips (including Western Flower Thrips)	3 to 4 oz.	<p>Add dry Acephate 75% SP to seed in the planter box to achieve an application rate of 3 to 4 oz. per acre.</p> <p>Even coverage of seed is required for acceptable insect control. Plants grown from improperly treated seed may not exhibit insect control. To obtain even coverage, fill planter box ½ full of seed, add ½ of the required amount of Acephate 75% SP and mix thoroughly. Add remainder of seed and Acephate 75% SP to the hopper box and thoroughly mix.</p> <p>Improved distribution of Acephate 75% SP on seed can be achieved by premixing both in a separate container and then transferring treated seed to hopper box.</p> <p>Use of Acephate 75% SP with air planter is not recommended since loss of product will occur.</p>
<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not use treated seed for food or feed purposes or process for oil. • Do not feed gin trash to livestock. • Do not allow animals to graze on treated areas. • Do not apply more than 5 1/3 lbs. of Acephate 75% SP per acre per crop cycle. This limitation includes all methods of application; i.e., hopper-box, in-furrow, and foliar. • Do not use treated seed for food or feed purposes or process for oil. • Acephate 75% SP cotton seed hopper box treatment is not permitted in Arizona, California or the Black Lands of Texas. 		

PEANUTS

PLANTER BOX SEED TREATMENT APPLICATION (HOPPER BOX)

PESTS CONTROLLED	ACEPHATE 75% SP APPLICATION RATE (PER 100 LBS. OF PEANUT SEED)	SPECIFIC INSTRUCTIONS
Thrips	<p>4 oz.</p> <p>DO NOT apply more than 4 oz. per 100 lbs. of peanut seed</p>	<p>Apply Acephate 75% SP at a rate of 4 oz. per 100 lbs. of peanut seed in the planter box as a dry powder. Ensure even coverage of the seed for acceptable insect control.</p> <p>Mixing: Follow the procedure below to ensure complete and adequate coverage of peanut seed:</p> <p>Step 1: Fill planter box 1/2 full of seed. Apply 1/3 the total amount of Acephate 75% SP evenly over the surface of the peanuts.</p> <p>Step 2: Fill planter box 2/3 full of seed. Apply 1/3 the total amount of Acephate 75% SP evenly over the surface of the peanuts.</p> <p>Step 3: Fill planter box with remaining seed. Apply 1/3 the total amount of Acephate 75% SP evenly over the surface of the peanuts.</p> <p>DO NOT attempt to mix Acephate 75% SP with the seed peanuts by stirring or other mechanical agitation. Such action will result in unacceptable damage to seed prior to planting.</p>
<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> Do not apply more than 5 1/3 lbs. of Acephate 75% SP per acre per crop cycle. This includes the use of Acephate 75% SP on peanuts as a foliar spray in addition to the peanut planterbox seed treatment use. Do not use treated seed for food for feed. 		

TOBACCO

SOIL APPLICATION

PESTS	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS	DAYS TO HARVEST
Ants (excluding carpenter and pharaoh ants)	Mound Treatment – evenly distribute 1 to 2 teaspoons over the mound	<p>Apply Acephate 75% SP in the early morning or late afternoon when ants are most active.</p> <p>Applications made under prolonged hot and dry conditions may be ineffective due to the location of ants deep within the nest.</p>	3
<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> Treat a maximum of 13 mounds per acre. Do not treat more than once per season. 			

NON CROP AREAS (FIELD BORDERS, FENCEROWS, ROADSIDES, DITCHBANKS, BORROW PITS)

PESTS	ACEPHATE 75% SP APPLICATION RATE (PER ACRE)	SPECIFIC INSTRUCTIONS
Ants (including Imported Fire Ants and Harvester Ants) (excluding carpenter and pharaoh ants)	Mound Treatment – evenly distribute 1 to 2 teaspoons over the mound.	Apply Acephate 75% SP in the early morning or late afternoon when ants are most active. Applications made under prolonged hot and dry conditions may be ineffective due to the location of ants deep within the nest.
USE RESTRICTIONS: <ul style="list-style-type: none"> Do not treat mound more than once per season. 		

MOUND TREATMENT OF FIRE ANTS AND HARVESTER ANTS IN TURFGRASS, EXCLUDING RESIDENTIAL TURF

PLANT	PESTS	AMOUNT OF ACEPHATE 75% SP	TIME OF APPLICATION	APPLICATION DIRECTIONS
Commercial Turf and Sodfarms Only	Imported Fire Ants and Harvester Ants	Mound Treatment – evenly distribute 1 to 2 teaspoons over the mound.	Apply Acephate 75% SP in the early morning or late afternoon when ants are most active. Applications made under prolonged hot and dry conditions may be ineffective due to the location of ants deep within the nest.	Apply the specified amount of Acephate 75% SP as directed. Grass in treated area may be injured.
USE RESTRICTIONS: <ul style="list-style-type: none"> Do not treat more than once per season. 				

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep pesticide in original container. Store in a cool, dry place away from excessive heat. Do not put concentrate or dilute into food or drink containers.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the extent consistent with applicable law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.